

Original Research Article

Malaria and HIV co-infection among COVID-19 Cohort in Selected Healthcare Facilities in Rivers State Nigeria

ABSTRACT

The COVID-19 pandemic began while improvement in malaria control and elimination had reached a peak. COVID-19 affected routine services particularly those for other killer infectious diseases including, HIV/AIDS and Malaria. This resulted in slowing progress in achieving control programs targets. The study evaluated malaria and HIV co-infection among COVID-19 cohort in selected healthcare facilities in Rivers State of Nigeria. The observational study showed a snapshot sampled only patients within the COVID-19 approved centers who consented to participate within the selected study area otherwise excluded. Simple random sampling technique was used to select 400 participants, giving subjects equal opportunity of participation. Data were basically primary sources however; secondary sources were used as well. Malaria testing was performed using microscopic method. Result outputs were presented on tables and charts. Study showed high prevalence of malaria infection among COVID-19 patients -327 (81.8%) but low rate of HIV co-infection- 2 (0.5%). Much consideration should be channeled towards malaria endemicity and co-infection with COVID-19 unless it confers protective advantage as postulated by some studies however, a large scale and more in-depth study is suggested.

Keywords: Malaria, HIV, co-infection, COVID-19 Cohort, Healthcare Facilities, Rivers State, Nigeria

Introduction

Malaria is amongst the diseases that have threatening to humanity, it is mostly spread through the bite of infected female *Anopheles* mosquito caused by *Plasmodium* parasites. About 241 million malaria cases and 627 000 malaria deaths was estimated worldwide in 2020, African is said to have the highest cases (WHO, 2021).

HIV infection usually result to reduced number of CD4 T cells, leading to AIDS, AIDS refers to a CD4 T-cell count < 200 cells/ μ L (Jordan *et al.*, 2020). HIV (Human Immunodeficiency virus) is a virus that destroys the CD4. HIV is a major public health issue globally, by the end of 2020 about 37.7 million lived with HIV over two thirds of whom (25.4 million) are in the WHO African Region. HIV virus has two main means of transmission horizontal and vertical transmission. Horizontal transmission is as a result of unprotected physical contact and may include unprotected sex with the affected victim, blood transfusion of infected blood. However, vertical transmission is direct transfer of virus from mother to child during pregnancy, breastfeeding or at delivery (Manish *et al.*, 2020). HIV disease can be managed by treatment regimens that comprise the combination of three or more antiretroviral (ARV) drugs, 28.2 million individuals living with HIV were receiving ART in 2021 globally (WHO, 2021).

منسّق: الخط: مائل، خط اللغة العربية
وغيرها: مائل

منسّق: الخط: مائل، خط اللغة العربية
وغيرها: مائل

[DS21] التعليق: In my opinion, it was better to focus on the prevalence in Rivers State Nigeria

[DS22] التعليق: No need

It is estimated that COVID-19 pandemic has spread to over 52 million persons and caused 1.2 million deaths from the time it was discovered in 2019 at Wuhan China (WHO, 2020). The first country in African where COVID-19 was first reported is Egypt in February 14, 2020 and has since spread across the region causing about 1.9 million infections and 46,000 deaths, (CDC, 2020). The population affected with COVID-19 in Sub-Saharan Africa is relatively low compared to global proportion of COVID-19 cases and deaths in other continents.

Though Africa has a relatively low proportion of COVID-19 cases and deaths, compare with other continents (WHO, 2021). It has an excessively higher burden of other infectious diseases, like malaria and HIV/AIDS. The possible effects of COVID-19 in controlling these diseases and any possible effect of clinical interactions between COVID-19 and these diseases is still a major public health concern in Africa, hence the need for this study. Also, this study would help in the understanding of the epidemiology and clinical course of COVID-19 with Malaria and HIV infection (WHO, 2019).

COVID-19 pandemic has caused disruptions in the prevention of HIV and malaria including treatment. Global fund report states that the COVID-19 pandemic threatens to reverse global progress in combating HIV and malaria. Specifically, malaria had halted progress with about 4% decrease in testing and a 0.5% treatment reduction. However, some prevention efforts heightened like the distribution of bed nets rose by 17% and use of mosquito spray increased by 3% in addition to 1% increase prophylactics administered to pregnant women who received preventive treatment for malaria (Kuehn, 2021; Peter, 2021).

The impact of COVID-19 has been massive to HIV prevention and control. Notwithstanding the progressing rate of people living with HIV/AIDS taking antiretroviral therapy, the decline in prevention services and testing in the COVID-19 pandemic period specifically the heat of it in 2020 were disturbing. Services such as public health HIV prevention outreach programs and services reduced by 11%. Also, HIV testing programs accessed decreased by 22% compared to the preceding pandemic year (2019). This disrupting consequence of the COVID-19 pandemic has placed those at risk of infection to have minimal access to the information and requisite tools for protection (Peter, 2021).

COVID-19 was first reported in Wuhan, China, on 8 December 2019 and declared a pandemic by the World Health Organization (WHO) on 11 March 2020 affecting about 185 countries (WHO, 2020). 45 countries out of these are in the WHO African region with endemic record of malaria and human immunodeficiency virus (HIV) with approximately 8767 confirmed COVID-19 cases and 413 deaths attributed to COVID-19 as of 13 April 2020. However, Africa has lower number of cases and deaths attributed to COVID-19 (Baud *et al.*, 2020; Dong *et al.*, 2020; WHO, 2020) although, cross country variation in the case fatality rate in the WHO Africa region exists (Dong *et al.*, 2020). The low rate of COVID-19 in the region of this study can be explained by age demography of the region comprising mainly of young people (Verity *et al.*, 2020; Gilbert *et al.*, 2020; Nuwagira & Muzoora, 2020; Paintsil, 2020).

Amimo *et al.* (2020) also attributed that to low testing in the area as most testing were done only following symptom presentation. Maze *et al.* (2018) suggested accurate diagnosis and prompt treatment as a measure to control malaria morbidity and mortality. Remarkably, COVID-19, malaria, and HIV share similar symptoms in some cases including cough, fever and others. The similarity of these symptoms poses significant concerns to the current measures to control, not only COVID-19, but the other diseases too (Amimo *et al.*, 2020).

The challenges posed by COVID-19 pandemic are enormous, the scarcity of published data on triple co-infection of malaria, HIV and COVID-19 heightened the researchers' interest to put it into consideration and investigate the rate of malaria and HIV co-infection among COVID-19 patients. The burden of single infection is big and much overwhelming in co-infection hence; information from the study is crucial and useful for clinicians for clinical management of patients.

Materials and Methods

Study Area:

The study was carried out in Rivers state, Nigeria. Rivers state is one of the states in the Niger Delta region found in South-South Nigeria. The state comprises of twenty-three local government areas and Port Harcourt is the capital of the state. The state has two tertiary hospitals and referral centers and other secondary hospitals and twenty-three primary healthcare centers managed by the government.

Study Design:

The study is a descriptive observational design which investigated the prevalence rate of malaria and HIV among COVID-19 patients in selected healthcare facilities in Rivers State of Nigeria.:-

Ethical Consideration

Ethical approval was obtained from the authorities of the selected healthcare facilities. Also, the laboratory heads gave approval prior to the use of the various laboratory sections. Informed written consent was obtained per participants. Researchers remained committed to privacy and confidentiality.

Inclusion Criteria

Conditions for recruitment and inclusion into this study were based on certain criteria such as; participants' approval and consent to participate in this study. Informed written consent was obtained from each subjects otherwise exclude. Also, only COVID-19 patients on admission within the designated healthcare facilities were included. Both males and females had equal chances of participation provided they were found within the selected facilities.

Exclusion Criteria

التعليق [DS23]: The method is very long and has many paragraphs. Its preferable to combine some them together (study area and design, inclusion and exclusion Criteria, sample Size and technique.

Exclusion from this study was on the bases of lack of consent by the patients, patients on unconscious state, COVID-19 patients found outside the selected accredited healthcare facilities, and those currently or recently treated for malaria parasite infections.

Sample Size

Sample size of 400 was obtained following Cochran sample size calculation for descriptive study. This was the sample size used and involved recruitment of four hundred (400) patients from the selected healthcare facilities.

Sampling Technique

Simple random sampling technique was used as sampling technique. The choice provided subjects equal opportunity of participating in the study. Random selection of the available healthcare facilities accredited for COVID-19 and thereafter, random selection of patients using the hospital register as the sample frame. All patients within the selected health facilities who met the inclusion criteria and gave consent participated in the study.

Laboratory Procedure

Conventional microscopy was used to carry out the malaria test following the procedure of thin and thick film making and smear. The smear was passed through staining using GeimsaGemisha stain which was latter viewed using the light binocular microscope. HIV testing was conducted using sequential approach. Serology testing was done using rapid detection kit from Alere and statpack for confirmatory. All laboratory procedures followed the standard clinical laboratory practices according Chessbrough (2010).

التعليق [DS24]: What was the approved test for diagnosing covid?

Validity/Reliability

Study instruments were validated and equipment precision obtained. The laboratory protocol adhered to all quality assurance measures. Quality control was used during the analysis stage. Standardization and calibration of equipments were performed. Standard clinical laboratory practice and WHO recommended guidelines were followed including all safety practices. All samples were run in duplicates.

التعليق [DS25]: No need

Statistical Analysis

Statistical analysis was performed with the use of Microsoft excel spread sheath for descriptive statistics. Percentage and frequency distribution were obtained, also, prevalence rate calculated. Results were presented on tables and charts.

Results

Table 1, figures 1 ~~&-and~~ 2 illustrate malaria and HIV co_infection in a cohort of four hundred (400) COVID-19 study participants showing malaria infection rate of 327 (81.8%) and

undetectable malaria infection rate of 73 (18.2%) among COVID-19 patients in the study population. The study result assumed marked malaria detection among the study population which implies a high malaria parasitaemia in the area of study.

On the other hand, the prevalence rate of HIV infection among the study population – cohort of COVID-19 patients appeared small 2 (0.5%) while 398 (99.5%) were negative to HIV infection. Comparatively, malaria parasitaemia among COVID-19 was high compared to HIV co-infection with COVID-19 as revealed in the study.

Table 1: Malaria and HIV co-infection among COVID-19 study participants

Study Participants	Number Examined	Malaria Infection (%)		HIV Infection (%)	
		Number Infected	Number Uninfected	Number Infected	Number Uninfected
COVID – 19 Patients	400	327 (81.8)	73 (18.2)	2 (0.5)	398 (99.5)




Figure 1: Prevalence of Co-infection with Malaria and HIV among COVID-19 Patients

Figure 2: Detection of Malaria and HIV among COVID-19 Patients

Discussion

التعليق [DS26]: It's the same as table 1

التعليق [DS27]: ???

The COVID-19 pandemic began while improvement in malaria control and elimination had reached high level (Amimo et al., 2020). COVID-19 affected routine services particularly those for other killer infectious diseases including, HIV/AIDS and Malaria. This resulted in slowing progress in achieving control programs targets (Ivers & Walton, 2020). The study investigated the prevalence of malaria, HIV co-infection among COVID-19 patients in selected health facilities in Nigeria.

This study recorded high detection of malaria infection among the study population (COVID-19 patients). Also, a high prevalence rate of 81.80% malaria co-infection with Covid-19 was recorded. The outcome of a high prevalence in malaria co-infection with COVID-19 maybe due to high malaria parasitaemia in the study area. The result is consistent with some research previously carried out.

World Health Organization estimated that 241 million malaria cases and 627,000 malaria deaths were recorded worldwide as at 2020, a rise 14 million cases and 69,000 deaths compared to 2019, with 47,000 deaths specifically due to COVID-19 related disruptions of malaria prevention, diagnosis, and treatment services (WHO, 2021b). This is evident in the high rate observed in this study despite the low rate of triple co-infections of COVID-19, malaria and HIV. The high rate of malaria in this study can be explained by these pandemic disruptions in addition to climatic change as well as mobility (Cella et al., 2019). Furthermore, poor and late health seeking behavior as a result of lack of access as well as pandemic restriction is a contributory factor (Bannister-Tyrrella, 2017).

Malaria and COVID-19 co-infection is said to be complex since malaria and COVID-19 have common symptoms, even though COVID-19 is more aggressive in adults while malaria affects more children (Micheal, 2021).

On the other hand, the occurrence of malaria and COVID-19 co-infection has been found to be of advantage for COVID-19 patients due to the immune response and chloroquine and its derivatives. Research has revealed possibility of cross immunity, where a pathogen can provide immunity against another pathogen (Teo *et al.*, 2018). Studies also have discovered that SARS-CoV-2 uses the angiotensin-converting enzyme 2 (ACE2) receptor to enter the host cells (Hoffmann *et al.*, 2019). More so, it has also been discovered that ACE1 and ACE2 polymorphisms protects the host from susceptibility. Additionally, studies have reported that interferons generated by lymphocytes which serve as an immune response to malaria have in vitro and in vivo efficacy against the coronavirus responsible for COVID-19.

The prevalence rate of HIV infection among the study population – cohort of COVID-19 patients which appeared low 2 (0.5%) in this study. This is similar to a case report by Toombs et al among 3 people living with HIV who were infected with coronavirus (Toombs *et al.*, 2020). They reported low prevalence (0.34%) of human immunodeficiency virus (HIV) in their studied population of 500 000 people suffering from COVID-19 in their hospital. This is also consistent with the report of Okoh et al and other studies on HIV and COVID-19 co-infection, there was no risk of co-infection (Okoh *et al.*, 2020).

Furthermore, Toombs and colleagues reported a patient living with HIV who had a high viral load of greater than one million copies per mL and a low CD4 count of 50 cells per μL . Notwithstanding, the adverse immunologic profile, the patient recovered from COVID-19 and was discharged. Their observation is consistent with other studies. The above observation is consistent with the clinical cure of at least four patients with COVID-19, who were newly diagnosed with HIV (Wu *et al.*, 2020)

Consequently, the study done by Thirumalaisamy and colleague reported that there is an indirect effect of the COVID-19 on HIV and Malaria infections in Africa. They explained that since the emergence of COVID-19, the availability of treatment and prevention materials is low and could cause increased mortality rate. (Thirumalaisamy *et al.*, 2021). This is similar to the research done by Vaughan who postulated that people with HIV and malaria may not go to healthcare facilities because of the fear of being infected with coronavirus would suffer the severity of both HIV and malaria (Vaughan, 2020).

Although the prevalence of COVID-19 and HIV co-infection showed low in this study, HIV/AIDS prevention and control have suffered drawback. Jewell and colleagues (2020) model estimate predicted disruption which has impacted negatively. About 50% distraction of condoms distribution and use in the population was predicted to raise incidence rates by 1.19 times, and a fold increase in HIV mortality. This prediction affected the service delivery which might have influenced the number of co-infection observed in this study as these disruptions could be the reason for low turnout and in-turn low detection rate. Furthermore, Holtzman *et al.* (2022) confirmed disruption. “ Information from simulation models also predicted an additional 0.002 - 0.15 COVID-19 deaths per 10,000 clients if HIV services were maintained while averting 19-146 HIV deaths per 10,000 clients, showing that it was beneficial to continue HIV services (Stover *et al.*, 2021)”.

Conclusion/recommendation

Malaria and HIV have been endemic in the region of this study with numerous efforts to combat the threat which have yielded success at a point. The COVID-19 pandemic started at a time at the peak of prevention and control of malaria and HIV attained marked improvement thereby affecting the achieved progress prior to the pandemic.

The study evaluated co-infections of malaria and HIV among COVID-19 patients with a resultant account of low prevalence rate within the population tested however, high rate of malaria prevalence was reported. The study suggests scale up of malaria prevention, control and elimination programme [program](#) in malaria endemic area.

Low rate of co-infections of malaria, HIV and COVID-19 could possibly be fear of being tested restricting people from getting involved therefore, campaigns encouraging people to overcome those fears will be one way to mitigation. More studies are required to evaluate triple co-

infections of malaria, HIV and COVID-19 as well as the clinical effect or relationship of this co-morbidity.

References

Amimo , F. Lambert, B. & Magit, A (2020). What does the COVID-19 pandemic mean for HIV, tuberculosis, and malaria control? *Tropical Medicine and Health* (2020) 48:32 | <https://doi.org/10.1186/s41182-020-00219-6>

Baud D, Qi X, Nielsen-Saines K, Musso D, Pomar L, Favre G. Real estimates of mortality following COVID-19 infection. *Lancet infect dis*. 2020

Center for Disease Control and Prevention (Africa) test updates on the COVID-19 crisis from Africa CDC. Accessed June 1, 2022. Available at <https://africacdc.org/covid-19/>

Cheesbrough M., part 1 . *District Laboratory Practice in Tropical Countries*. 2nd Edition. Cambridge: Cambridge University Press; 2010

Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis*. 2020 May;20(5):533–44.

Gilbert M, Pullano G, Pinotti F, Valdano E, Poletto C, Boëlle P-Y, et al. Preparedness and vulnerability of African countries against importations of COVID-19: a modelling study. *Lancet*. 2020;395(10227):871–7.

Hellewell J, Abbott S, Gimma A, Bosse NI, Jarvis CI, Russell TW, et al. Feasibility of controlling COVID-19 outbreaks by isolation of cases and contacts. *Lancet Glob Health*. 2020;8(4):e488–e96.

Hoffmann, M., Kleine-Weber, H., Kruger, N., Muller, M., Drosten, C., Pohlmann, S., (2020). The novel coronavirus uses the SARS-coronavirus receptor ACE2 and the cellular protease TMPRSS2 for entry into target cells. bioRxiv.

Jordan, R. E., Adab, P., Cheng, K. K., (2020). Covid-19 risk factors for severe disease and death. *British Medical Journal*. 368:1198.

Kuehn BM. (2021). COVID-19 Rolls Back Progress Against HIV, TB, and Malaria. *JAMA*. 2021;326(15):1471. doi:10.1001/jama.2021.18252

Manish, G., Haci, M., Baskonus, A., (2020) Regarding new positive, bounded and convergent numerical solution of nonlinear time fractional HIV/AIDS transmission model 139:0960-0779, <https://doi.org/10.1016/j.chaos.2020.110096>.

منسّق: لون الخط: أحمر

منسّق: لون الخط: أحمر

منسّق: لون الخط: أحمر

منسّق: لون الخط: أحمر

منسّق: لون الخط: أحمر

منسّق: لون الخط: أحمر

Maze MJ, Bassat Q, Feasey NA, Mandomando I, Musicha P, Crump JA. The epidemiology of febrile illness in sub-Saharan Africa: implications for diagnosis and management. *Clin Microbiol Infect.* 2018;24(8):808–14.

منسّق: لون الخط: أحمر

Michael, U., Anyanwu (2021). The association between malaria prevalence and COVID-19 mortality. *BMC Infectious Disease* 21:975 doi: 10.1186/s12879-021-06701-8

Nuwagira E, Muzoora C. Is Sub-Saharan Africa prepared for COVID-19? *Tropical Medicine and Health.* 2020;48(1):1–3.

Paintsil E. COVID-19 threatens health systems in sub-Saharan Africa: the eye of the crocodile. *J Clin Invest.* 2020.

منسّق: لون الخط: أحمر

Peter, S. (2021). COVID-19 Hit HIV, TB and Malaria Programs Hard, We're Fighting Back. Global Fund. Retrieved from; <https://www.theglobalfund.org/en/blog/2021-09-17-covid-19-hit-hiv-tb-and-malaria-programs-hard-we-re-fighting-back/> Accessed 6th June, 2022.

Thirumalaisamy, P., Velavan, Christian, G., Meyer, Meral, E. Peter, G., Francine, N. (2021) COVID-19 and syndemic challenges in 'Battling the Big Three': HIV, TB and malaria, *International Journal of Infectious Diseases* 106: 29-32, <https://doi.org/10.1016/j.ijid.2021.03.071>

منسّق: لون الخط: أحمر

Toombs, J., M., Van den, Abbeele, K., Democratis, J., Merricks, R., Akj, Mandal, Missouri C., G., (2020) COVID-19 in 3 people living with HIV in the United Kingdom. *Journal Medical Virology.*

منسّق: لون الخط: أحمر

منسّق: لون الخط: أحمر

Vaughan A. (2020). Covid-19 could have disastrous impacts on HIV, TB and malaria. *New scientist* (1971), 247(3291), 8. [https://doi.org/10.1016/S0262-4079\(20\)31225-2](https://doi.org/10.1016/S0262-4079(20)31225-2)

Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis.* 2020

منسّق: لون الخط: أحمر

World Health Organization Coronavirus disease 2019 (COVID-19): situation reports <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports> (2020), Accessed 31st May 2022

منسّق: لون الخط: أحمر

منسّق: لون الخط: أحمر

World Health Organization (2021). HIV and AIDS Global health sector strategy on 2021, Accessed 31st May 2022

منسّق: لون الخط: أحمر

World Health Organization (2021b) Malaria. Accessed May 31 2022

World Health Organization. World malaria report World Health Organization, Geneva (2020) Accessed 1st June 2022

منسّق: لون الخط: أحمر

World Health Organization, (2019). Coronavirus disease 2019 (COVID-19): situation reports. World Health Organization; <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>.

منسق: لون الخط: أحمر

Wu, Q., Chen, T., Zhang, H. (2020) Recovery from COVID-19 in two patients with coexisted | HIV infection. *Journal of Medical Virology*. 10.1002/jmv.26006

UNDER PEER REVIEW