

Effect of seaweed extracts on growth, yield parameters in Chickpea (*Cicer arietinum.L*)

ABSTRACT

This field experiment entitled “Effect of seaweed extracts on growth, ~~yield parameters~~ yield parameters in Chickpea (*Cicer arietinum.L*)” was conducted during rabi at Field Experimentation Centre of the Department of Genetics and Plant Breeding, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj, Uttar Pradesh, India during 2019 & 2020. The experiment was consisted of 2 varieties and 14 treatments comprising of seaweed extracts (i.e., *Ascophyllum nodosum*, red and brown algae) which was laid in RBD (Randomized Block Design) with kabuli & desi chick pea, 14 treatments, 03 replications.

The results shows that among all the treatments, the genotypes kabuli & desi chickpea treated with 2.0% *Ascophyllum nodosum* for 6 hours recorded the maximum value in growth parameters such as percent field germination after four (10.74 & 10.74), seven (49.26 & 47.41), and ten (77.41 & 70.74) DAS, Days to 50% flowering (88.67 & 86.00), days to maturity (154.00 & 156.33), number of branches 25 (17.93 & 16.60) and 50 DAS (36.70 & 33.57) and height at 25 (4.47 & 3.93) and 50 (7.27 & 7.00) DAS.

Similar results were also recorded in different yield parameters in both kabuli & desi chickpea genotypes such as number of pods plant (61.52 & 60.24), number of pods per plot (1025.67 & 922.00), number of seeds per plant (102.5 & 100.4), number of pods per plant (14.25 & 15.25), seed yield per plant (28.30 & 25.35), seed yield per plot (358.44 & 328.86), biological yield (3187.42 & 3061.24), economical yield (1341.67 & 1231.33), and harvest index showing most superior values when treated with treatment 2.0% *Ascophyllum nodosum* for 6 hours.

Remaining treatments i.e, treatment with 0.5% *Ascophyllum nodosum* solution, 1.0% *Ascophyllum nodosum* solution, 0.5% red and brown algae solution, 1.0% red and brown algae solution, 2.0% red and brown algae solution for 6 hours recorded the second most effective treatments observed significantly superior than untreated checks.

Keywords: *Ascophyllum nodosum*, Chickpea, red and brown algae.

Introduction

Chickpea (*Cicer arietinum* L.) belongs to Fabaceae family. It is self pollinated and true diploid ($2n=2x=16$) in nature. In the world, chickpea stands third most important legume. In the world, India is one among the pulse producing countries. 62-67 percent of total production in globe was contributed by India. Among the countries in world, though India is one of the largest country in pulse production, to meet annual domestic consumption, 2-3 million tons of pulses are imported annually. The current per capita availability of pulses as 80g/day, recommended by FAO is very low that can't meet the per capita requirement. Chickpea contains low cholesterol, sodium, fat and high fibre contents. Chickpea also a rich source of protein and minerals. 100g of boiled chickpea contains energy- 164 calories, fat- 2.6g (in which only 0.27g fat is saturated), dietary fibre-7.9g, protein-8.9g and dietary calcium-49 to 53mg/100g. Chickpea used in both human and animal consumption. Seaweeds are marine algae, that are macroscopic and found attached at the bottom of shallow coastal waters. Marine algae that grows upto 180 meters depth in intertidal, deep and shallow waters and also grows on solid substrates like rocks, pebbles and dead corals in back waters. The extracts of seaweed has beneficial effects on germination of seeds and on growth of the plant (Bhosle *et al.*, (1975); Kumar *et al.*,

Comment [sm1]: remove the paragraph in abstract

Comment [sm2]: mention the scientific name for both algae

Comment [sm3]: mention like *A. nodosum*, full genus name required only first time in a manuscript. similarly correct in other portions of manuscript.

(1994). Verkleij (1992) reported that, on application of liquid seaweed fertilizer to soil enhances the water retention capacity. Extracts of seaweed enhance antioxidant properties (Verkleij, 1992), contains amino acids, vitamins, cytokinins, auxins, major & minor nutrients and abscisic acid, that acts as growth promoting substances that promotes growth, yield and develops tolerance against environmental stress in plants (Zhang *et al.*, 2003), increase soil nutrient uptake (Turan and Kose, 2004). From marine algae, bioactive substances are extracted, used in agricultural and horticultural crops and many beneficial effects on yield and quality enhancement can be achieved. Now-a-days, ~~in many~~ in many crops including pulses, cereals and vegetable crops, using liquid seaweed extracts as foliar sprays has gained importance. Than chemical fertilizers, seaweed extracts are free from toxins, biodegradable, pollution free and safe to human, animals and birds (Dhargalkar and Pereira, 2005).

Comment [sm4]: no clarity in sentence

Formatted: Highlight

Formatted: Highlight

MATERIALS AND METHODS:

This Research work was carried out at experimental field during rabi season 2019-2020. Department of Genetics and plant breeding, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P). The experiment was consisted of 2 varieties and 14 treatments comprising of seaweed extracts (i.e., *Ascophyllum nodosum*, red and brown algae).

Preparation of Sea Weed Extract Solution: For preparing Sea weed extract solution, measure 5ml, 10ml, 20ml Sea weed extracts in a beaker and add 1000 ml- distilled water by constant stirring. ~~The final~~ The final volume ~~constitute~~ made up to one litre, and then it became 0.5%, 1.0%, 2.0% stock solution of Sea weed extracts.

The treatments were represented as T0- Desi Chickpea Control (Untreated), T1- Desi seeds treated with 0.5% *Ascophyllum nodosum* solution @ 6 hours, T2-Desi seeds treated with 1.0% *Ascophyllum nodosum* solution @ 6 hours, T3-Desi seeds treated with 2.0% *Ascophyllum nodosum* solution @ 6 hours, T4-Desi seeds treated with 0.5% red and brown algae solution @ 6 hours, T5-Desi seeds treated with 1.0% red and brown algae solution @ 6 hours, T6-Desi seeds treated with 2.0% red and brown algae solution @ 6 hours, T7-Kabuli Chickpea Control (Untreated), T8-kabuli seeds treated with 0.5% *Ascophyllum nodosum* solution @ 6 hours, T9-kabuli seeds treated with 1.0% *Ascophyllum nodosum* solution @ 6 hours, T10-kabuli seeds treated with 2.0% *Ascophyllum nodosum* solution @ 6 hours, T11-kabuli seeds treated with 0.5% red and brown algae solution @ 6 hours, T12-kabuli seeds treated with 1.0% red and brown algae solution @ 6 hours, T13-kabuli seeds treated with 2.0% red and brown algae solution @ 6 hours.

Comment [sm5]: treatment detail put in table for clarity

The data recorded during the course of the investigation will be subjected to Statistical Analysis by 1 X 1 RBD, as per the method "Analysis of Variance (ANOVA) technique". Experiment will be laid out in RBD and the treatments will be replicated three times. The significant and non-significant effect was judged with the help of "F" (variance ratio) table. The significant difference between the means was tested against the critical difference of 5% level. For testing the hypothesis, ANOVA table was used.

Comment [sm6]: correct the sentence grammar

RESULTS AND DISCUSSION:

Comment [sm7]: 2 subheadings required similar to table format

Result is not discussed in manuscript.

Highest field germination percentage after four, seven and ten days of treatment application in kabuli genotype was recorded in treatment T10 (2.0% *Ascophyllum nodosum* solution) with 10.74, 49.26 & 77.41 percent and was found statistically at par with treatment T9 (1.0% *Ascophyllum nodosum* solution) with 10.37, 44.44 & 72.22 percent and T8 (2.0% red and brown algae solution) with 9.26, 43.33 & 70.00 percent respectively.

The highest days to 50% flowering and days to maturity in kabuli genotype was recorded in treatment T10 (2.0% *Ascophyllum nodosum* solution) i.e., 88.67 & 156.33 days and was found at par with treatment T8 (0.5% *Ascophyllum nodosum* solution) and T9 (1.0% *Ascophyllum nodosum* solution) with 85.00 & 151.67 and 84.33 & 149 days respectively whereas in desi chickpea genotypes the application of treatment T2 (1.0% *Ascophyllum nodosum* solution) & T3 (2% *Ascophyllum nodosum* solution) recorded highest days to 50% flowering & maturity with 87.00 & 154.00 days.

The maximum plant height after 25 & 50 days of sowing and treatment application in kabuli genotype was recorded in treatment T10 (2.0% *Ascophyllum nodosum* solution) with 17.93 & 36.70 and was found statistically at par with treatment T9 (1.0% *Ascophyllum nodosum* solution) with 16.87 & 34.73 whereas, in desi chickpea genotypes the application of treatment T3 (2% *Ascophyllum nodosum* solution) recorded highest plant height with 16.60 & 33.57 and was found statistically at par with treatment T2 (1% *Ascophyllum nodosum* solution) with 13.87 & 26.60.

The maximum number of branches after 25 & 50 days of sowing and treatment application in kabuli genotype was recorded in treatment T10 (2.0% *Ascophyllum nodosum* solution) with 4.47 & 7.27 branches and was found statistically at par with treatment T9 (1.0% *Ascophyllum nodosum* solution) with 4.07 & 6.80 branches whereas, in desi chickpea genotypes the application of treatment T3 (2% *Ascophyllum nodosum* solution) recorded highest number of branches with 3.93 & 7.00 branches and was found statistically at par with treatment T2 (1.0% *Ascophyllum nodosum* solution) with 3.27 & 5.60 branches.

The maximum numbers of pods per plant & pods per plot after sowing and treatment application in kabuli genotype was observed in treatment T10 (2.0% *Ascophyllum nodosum* solution) with 61.55 pods/ plant & 1025.67 pods/ plot and was found statistically at par with treatment T9 (1.0% *Ascophyllum nodosum* solution) with 56.20 pods/ plant & 860.33 pods/ plant respectively whereas, in desi chickpea genotypes the application of treatment T3 (2% *Ascophyllum nodosum* solution) recorded highest number of pods with 60.24 pods/ plant & 922.00 pods/ plot and was found statistically at par with treatment T2 (1.0% *Ascophyllum nodosum* solution) 57.31 pods/ plant & 901.33 pods/ plot.

The maximum number of seeds per plant & seeds per plot after sowing and treatment application in kabuli genotype was recorded in treatment T10 (2.0% *Ascophyllum nodosum* solution) with 102.5 seeds/ plant & 1525 seeds/ plot and was found statistically at par with treatment T8 (0.5% *Ascophyllum nodosum* solution) with 93.07 seeds/ plant & 1393 seeds/ plot whereas, in desi chickpea genotypes the application of treatment T3 (2% *Ascophyllum nodosum* solution) recorded highest number of seeds with 100.4 seeds/ plant & 1425 seeds/ plot and was found statistically at par with treatment T2 (1.0% *Ascophyllum nodosum* solution) with 84.4 seeds/ plant & 1410 seeds/ plot.

The maximum seed yield per plant and per plot after sowing and treatment application in kabuli genotype was recorded in treatment T10 (2.0% *Ascophyllum nodosum* solution) 28.30 gm/ & 358.44 gm/ plot plant and was found statistically at par with treatment T9 (1.0% *Ascophyllum nodosum* solution) with 26.78 gm/ & 333.18 gm/ plot plant whereas, in desi chickpea genotypes the application of treatment T3 (2% *Ascophyllum nodosum* solution) recorded highest seed yield per plant & per plot with 25.35 gm/ plant & 328.86 gm/ plot and was found statistically at par with treatment T2 (1.0% *Ascophyllum nodosum* solution) 23.82 gm/ plant & 319.59 gm/ plot.

The maximum biological & yield per hectore after sowing and treatment application in kabuli genotype was recorded in treatment T10 (2.0% *Ascophyllum nodosum* solution) with 3187.42 & 1341.67 kg/ ha and was found statistically at par with treatment T9 (1.0% *Ascophyllum nodosum* solution) with 3092.49 & 1252.00 kg/ ha whereas, in desi chickpea genotypes the application of treatment T3 (2% *Ascophyllum nodosum* solution) recorded highest biological yield per hectore with 3061.24 & 1231.33 /kg and was found statistically at par with treatment T2 (1.0% *Ascophyllum nodosum* solution) 2987.73 & 1217.00/kg.

The maximum harvest index after sowing and treatment application in kabuli genotype was recorded in treatment T7 (Untreated check) i.e., 44.08 and was followed by treatment T12 (1.0% red and brown algae solution) and T13 (2.0% red and brown algae solution) with 42.63 and 42.43 respectively whereas, in desi chickpea genotype, the application of treatment T0 (Untreated check) recorded the highest harvest index i.e., 43.67 and was followed by treatment T1 (0.5% *Ascophyllum nodosum* solution) and T5 (1.0% red and brown algae solution) with 42.39 and 41.94 respectively.

TABLE 2 : Effect of seaweed extracts on different yield parameters:

Tr. No.		Number of pods per plant	Number of pods per plot	Seed yield gm plant ⁻¹	Seed yield gm plot ⁻¹	Number of seeds per plant	Number of seeds per plot	Biological yield (Kg ha ⁻¹)	Economical yield (Kg ha ⁻¹)	Harvest index
T0	Deysi Chickpea Control (Untreated)	41.56	626.67	16.42	272.70	69.3	1044	2313.05	1010.00	43
T1	Deysi seeds treated with 0.5% <i>Ascophyllum nodosum</i> solution	49.68	769.67	22.41	316.35	82.8	1282	2716.33	1151.33	42
T2	Deysi seeds treated with 1.0% <i>Ascophyllum nodosum</i> solution	57.31	901.33	23.82	319.59	84.4	1410	2987.73	1217.00	40
T3	Deysi seeds treated with 2.0% <i>Ascophyllum nodosum</i> solution	60.24	922.00	25.35	328.86	100.4	1425	3061.24	1231.33	40
T4	Deysi seeds treated with 0.5% red and brown algae solution	50.44	767.67	20.84	305.37	84.1	1278	2837.07	1171.67	41
T5	Deysi seeds treated with 1.0% red and brown algae solution	48.68	768.33	20.58	310.86	81.1	1279	2648.16	1110.67	41
T6	Deysi seeds treated with 2.0% red and brown algae solution	51.32	784.67	19.89	300.01	85.5	1307	2847.74	1152.00	40
T7	Kabuli Chickpea Control (Untreated)	44.68	817.67	17.56	291.15	74.5	1167	2446.55	1078.33	44
T8	Kabuli seeds treated with 0.5% <i>Ascophyllum nodosum</i> solution	55.91	976.00	22.54	311.04	82.1	1393	3004.98	1207.00	40
T9	Kabuli seeds treated with 1.0% <i>Ascophyllum nodosum</i> solution	56.20	860.33	26.78	333.18	93.7	1228	3092.49	1252.00	40
T10	Kabuli seeds treated with 2.0% <i>Ascophyllum nodosum</i> solution	61.52	1025.67	28.30	358.44	102.5	1525	3187.42	1341.67	42
T11	Kabuli seeds treated with 0.5% red and brown algae solution	52.28	929.33	21.78	325.89	87.1	1327	2888.33	1183.67	40
T12	Kabuli seeds treated with 1.0% red and brown algae solution	51.97	955.67	21.06	329.04	86.4	1317	2653.20	1131.00	42
T13	Kabuli seeds treated with 2.0% red and brown algae solution	54.16	874.00	19.41	328.59	90.3	1248	2896.00	1228.67	42
	SE(m) ±	3.09	36.49	1.27	12.01	3.51	48.41	101.99	41.71	
	CD at 5%	8.99	106.07	3.70	34.9	10.19	140.72	144.23	121.24	
	CV %	10.20	7.39	10.06	6.57	7.06	6.44	6.25	6.14	

TABLE 3 : Analysis of variance for quantitative characters in chickpea.

S.No.	Characters	Replications (df=2)	Mean sum of square Treatments (df=13)	Error (df=26)
1.	Percent Field Emergence at 4 DAS	46.77	5.22*	37.33
2.	Percent Field Emergence at 7 DAS	918.90	55.65*	15.80
3.	Percent Field Emergence at10DAS	160.54	1673.48*	56.89
4.	Days to maturity	1152.93	104.68*	41.24
5.	Plant height at 25 DAS	12.41	27.65*	55.73
6.	Plant height at 50 DAS	1.20	135.77*	10.78
7.	No. of branches at 25 DAS	2.56	4.06*	0.12
8.	No. of branches at 50 DAS	11.56	7.68*	0.49
9.	No. of pods per plant	277.40	92.74*	28.72
10.	No. of pods per plot	6195.07	34002.33*	3994.43
11.	Seed yield per plot	9.43	33.18*	4.86
12.	No. of seed per plant	995.13	237.25*	36.89
13.	No. of seeds per plot	19016.74	41627.27*	7029.92
14.	Biological yield	61204.50	185800.68*	31204.64
15.	Economical yield	14229.74	19926.65*	5218.15

* Significant at 5% level of significance.

CONCLUSION:

Among the different seaweed extracts used during the experiment the most superior values are recorded from the treatment with ~~2% 0%~~ *Ascophyllum nodosum* solution in both kabuli as well as desi chickpea genotype. An agronomical parameters such as field germination percentage, days to 50% flowering, days to maturity, plant height and number of branches was recorded maximum in treatment with 2.0% *Ascophyllum nodosum*. Similarly, the same treatment with ~~2-0%~~ *Ascophyllum nodosum* recorded maximum numbers of pods per plant & number pods per plot, number seeds per plant & number seeds per plot and yield per plant & yield per plot. Yield parameters such as biological yield per ~~hectare or~~, economical yield per hector and harvesting index was recorded was found highest in treatment 2.0% *Ascophyllum nodosum* and was found at par with treatment 1.0% *Ascophyllum nodosum* followed by treatment 0.5% *Ascophyllum nodosum*. The remaining treatment 2.0% red and brown algae, 1.0% red and brown algae and 0.5% red brown algae recorded the second best treatment. The maximum growth and yield parameters can be ~~arehived-achieved~~ through the treatment with 2.0% *Ascophyllum nodosum* solution in both kabuli as well as desi chickpea genotype.

Comment [sm9]: no need to mention point zero

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES:

- Abou El-Yazied A., El-Gizawy A.M., Ragab M.I. and E.S. Hamed (2012) Effect of Seaweed Extract and Compost Treatments on Growth, Yield and Quality of Snap Bean, *Journal of American Science* 2012;8(6)
- Anisimov M.M. Chaikina E.L. Klykov A.G. and Rasskazov V.A., (2013) Effect of Seaweeds Extracts on the Growth of Seedling Roots of Buckwheat (*Fagopyrum esculentum Moench*) is depended on the Season of Algae Collection. *Agriculture Science Developments*, 2 (8): 67-75.

Comment [sm10]: References should present in uniform format. please correct the same in manuscript and reference section also.
so many references not covered in this manuscript.

- Anonymous. (2016). 5th National seminar on coarse cereals development, challenges and opportunities in the country. Sponsored by Directorate of millets development, Jaipur (India). March 20-21, 2016
- Arthur G.D., Stirk W.A. and Van Staden J. (2003) Effect of seaweed concentrates on the growth and yield of three varieties of *Capsicum annuum*. *S. Afr. J. Bot.* **69**:207-211.
- Basavaraja P.K., Yogendra N.D., Zodape S.T., Ravi Prakash and Ghosh A. (2018) Effect of seaweed sap as foliar spray on growth and yield of hybrid maize, *Journal of Plant Nutrition*, DOI: 10.1080/01904167.2018.1463381
- Bhosle N.B., Dhargalkar V.K. and Untawale A.G. (1975) Effect of seaweed liquid fertilizer on the yield of *Zizphus rugosa* (Bores), *International symposium on MarineAlgae*. India Ocean eg., India, Abstracts:27.
- Crouch I.J. and Van Staden J. (1994) Commercial seaweed products as biostimulants in horticulture. *J. Home Consumer Horticult.* **1**:19-76.
- Dhargalkar V.K. and Pereira N. (2005) Seaweed: promising plant of the millennium. *Sci. Cul.* **71**:60-66.
- Ferreira M.I. and Lourens A.F. (2002) The efficacy of liquid seaweed extract on the yield of Canola plants. *S. Afr. J. Plant Soil.* **19**:159-161.
- Gehan A. Elsharkawy, Hanaa S. Hassan and Hassan A.H. Ibrahim (2017) Effect of promoting diastrophic bacteria and seaweed extract formula on growth yield and quality of pea (*Pisum sativum* L.) plants, *Alexandria science exchange journal* vol-**40**,no .1.
- Hagharast M., Melaki F.S., Sinaki J.M. and Jerei G.H. (2012) Mitigation of drought stress in chickpea through application of humic acid and seaweed extract, *Crop production in environmental stress*, vol.**4**(8): pp. 59-71.
- Hamed E.S. (2012) Effect of seaweed extract and compost treatments on growth, and quality of snap bean. Ph.D. Agric. Sc. Ain Shams University.
- Hanan E.O. and Olfat M.A. Salem (2011) Effect of seaweed extracts as foliar spray on sunflower yield and oil content, *Egyptian J. of Phycol.* Vol.**12**, (ISSN: 1110-8649)
- Hussain H.I., Naga Kasinadhuni N. and Tony Ariolil T. (2021) The effect of seaweed extract on tomato plant growth, productivity and soil, *Journal of Applied Phycology* <https://doi.org/10.1007/s10811-021-02387-2>

- Iswarya S., Latha K.R. and Srinivasan K. (2019) Evaluation of seaweed extract on growth determinants, yield and biochemical parameters of greengram (*Vigna radiata*), *Journal of Pharmacognosy and Phytochemistry* vol.8(3): 1861-1864.
- Kalaivanan C. and Venkatesalu V. (2012) Utilization of seaweed *Sargassum myriocystum* extracts as a stimulant of seedlings of *Vigna mungo* (L.) Hepper, *Spanish Journal of Agricultural Research* vol. 10(2), 466-470 ISSN: 1695-971-X
- Kalaiyany V., Sutharsan S. and Sikrishnah S. (2019) Effect of natural and commercially available seaweed liquid extracts on growth and yield of *vigna unguiculata* L. *Asian journal of biological sciences*.
- Karthikeyan K. and Shanmugam M. (2015) Yield and oil content of peanut (var. TMV-7) and sunflower (var. Co-2) applied with bio-stimulant AQUASAP manufactured from seaweed, *African Journal of Agricultural Research*, Vol. 10(25), pp. 2537-2543.
- Kavipriya R., Dhanalakshmi P. K., Jayashree S. and Thangaraju N. (2011) Seaweed extract as a biostimulant for legume crop, greengram, *Journal of Ecobiotechnology* 2011, 3(8): 16-19 ISSN: 2077-0464
- Kumar, Venkatranaman, Mohan V.R., Murugewari M. and Muthuswami S. (1994). Effect of crude and commercial seaweed extract on seed germination and seedling growth in green gram and black gram. *Seaweed Res. & Utilization*, 16:23-28.
- Laura G.C.B., Fernando S.R., Gustavo H.C., Ernesto R.B. and Rosalba M.H.H. (2017) Effect of seaweed liquid extracts from *Ulva lactuca* on seedling growth of mungbean (*Vigna radiata*), *J Appl Phycol* DOI 10.1007/s10811-017-1082-x
- Layek J. & Das A., Idapuganti R.G., Sarkar D., Ghosh A., Zodape S.T., Lal R., Yadav G., Panwar A., Ngachan S. and Meena R.W. (2017) Seaweed extract as organic bio-stimulant improves productivity and quality of rice in eastern Himalayas, *J Appl Phycol* DOI 10.1007/s10811-017-1225-0
- Narasimha Rao G. M. and Chatterjee R (2014) Effect of seaweed liquid fertilizer from *Gracilaria textorii* and *Hypnea musciformis* on seed germination and productivity of some vegetable crops, *Universal Journal of Plant Science* 2(7): 115-120.
- Pittaway J. K., Roberston I. K. and Madeleine J. B (2008). Chickpea may influence fatty acid and fiber intake in an Libitum Diet, leading to small improvements in serum lipid profile and glycemic control. *Journal of the Academy of Nutrition and Dietetics*, 108(6): 1006- 1013
- Pramanick B., Brahmachari K. and Ghosh A. (2013) Effect of seaweed saps on growth and yield improvement of greengram, Vol. 8(13), pp. 1180-1186, DOI: 10.5897/AJAR12.1894

- Rathore S.S., Chaudhary D.R., Boricha G.N., Ghosh A., Bhatt B.P., Zodape S.T. and Patolia J.S. (2009) Effect of seaweed extract on the growth, yield and nutrient uptake of soybean (*Glycine max*) under rainfed conditions, *South African Journal of Botany* **75** (2009) 351–355.
- Rawheya A. Salah El Din., Elbakry A.A., Ghazi S.M. and Olfat Hamid M.A. (2008) Effect of seaweed extract on the growth and yield of faba bean (*vicia faba* l.), *Egyptian J. of Phycol.* Vol. **9**, ISSN: 1110-8649
- Russo R.O. and Beryln G.P. (1990) The use of organic biostimulants to help low inputs. *J. Sustain. Agric.* **1**:9-42.
- Sarhan T.Z. and Ismael S.F. (2014) Effect of Low Temperature and Seaweed Extracts on Flowering and Yield of Two Cucumber Cultivars (*Cucumis sativus* L.), *International Journal of Agricultural and Food Research* ISSN 1929-0969 | Vol. **3** No. 1, pp. 41-54 (2014)
- Sasikala M., Indumathi E., Radhika S. and Sasireka R. (2016) Effect of Seaweed Extract (*Sargassum tenerrimum*) on seed germination and growth of tomato plant (*Solanum lycopersicum*), *International Journal of ChemTech Research*, Vol.**9**, No.09 pp 285-293, ISSN: 0974-4290.
- Selvam G.G. and Sivakumar K. (2014) Influence of seaweed extract as an organic fertilizer on the growth and yield of *Arachis hypogea* L. and their elemental composition using SEM– Energy Dispersive Spectroscopic analysis, *Asian Pacific Journal of Reproduction*, vol.**3**(1): 18-22
- Shahbazi F., Nejad M.S., Salimi A. and Gilani A. (2015) Effect of seaweed on the growth and biochemical constituents of wheat, *International journal of agriculture and crop science*, vol.**8**(3):pp283-287:ref.23
- Sivasangari Ramya S., Vijayanand N. and Rathinavel S. (2012) Influence of seaweed liquid fertilizers on growth, biochemical and yield parameters of cluster bean plant, *J. Green Bioenergy* **1** (1): 19 – 32
- Sivasankari S., Venkatesalu V., Anantharaj M. and Chandrasekaran M. (2006) Effect of seaweed extracts on the growth and biochemical constituents of *Vigna sinensis* . *Bioresource Technol.*, **97**: 1745–1751.
- Sridhar S. and Rengasamy R. (2011) Influence of Seaweed Liquid Fertilizer on Growth and Biochemical Characteristics of *Arachis hypogea* L. under field trial, *Journal of Ecobiotechnology*, vol.**3**(12): 18-22 ISSN: 2077-0464
- Subbulakshmi S., Saravanan N., Sivaprakash M. and Harisudan C. (2009) Nutrition management for pulses. *J. Pulses Research*, **7** (8) : 48.
- Swaminathan M.S. (1981). Improving pulse production and productivity challenges ahead's. *Pulse Crops Newsletter*,(2): 25.

- Thirumaran G., Arumugam M., Arumugam R. and Anantharaman P. (2009) Effect of seaweed liquid fertilizer on growth and pigment concentration of *Abelmoschus esculentus* (l) medikus, *American-Eurasian Journal of Agronomy* vol.2 (2): 57-66, ISSN 1995-896X.
- Thirumaran G., Arumugam M., Arumugam R. and Anantharaman P. (2009) Effect of seaweed liquid fertilizer on growth and pigment concentration of *Abelmoschus esculentus* (I) Medikus. *Am Eurasian J. Agron.*, **2**: 5766.
- Turan M. and Kose C. (2004) Seaweed extracts improve copper uptake of grapevine. *Acta. Agric. Scand.* B-S P. **54**:213-220.
- Verkleij F.N. (1992). Seaweed extract in agriculture and horticulture: A review. *Biol. Agric. & Hort.*, **8** : 309-324.
- Zewail, R. M. Y (2014) effect of seaweed extract and amino acids on growth and productivity and some biocostituents of common bean (*Phaseolus vulgaris* L) plants, *J. Plant Production*, Mansoura Univ., Vol. **5** (8): 1441 – 1453.

UNDER PEER REVIEW