

# **Original Research Article**

## **Impact of Storage Period on Fertility and Mortality in Chabro Breeds of Poultry**

### **Abstract**

An investigation was conducted to study the effect of storage period on fertility and hatchability in large, medium and small sized eggs of Chabro breed. The study was conducted at poultry farm of SKN College of Agriculture, Jobner. For the present study, a total of 297 eggs were collected for this study. These eggs were grouped into three egg size categories i.e. Small (38-44 g), medium (45-52 g) and large (53-59 g). To study the impact of incubation period, eggs were stored at 21 °C temperature for three different time periods (fresh, 3 day and 6 day). It was observed that egg fertility was increased on 3<sup>rd</sup> day and decreased on 6<sup>th</sup> day as compared to 0 day. For small egg class, egg fertility was 95.55, 96.02 and 92.13% respectively, on 0, 3 and 6 day of storage. Similarly, for medium-sized egg class, fertility was 95.17, 95.64 and 90.28% and for large sized egg class, it was 91.11, 92.15 and 87.41% respectively, on 0, 3 and 6 day of storage. Embryonic mortality increased with increase in storage duration. For small egg class, egg mortality was 20.95, 17.56 and 24.87% respectively, on 0, 3 and 6 day of storage. Similarly, for medium-sized eggs class, mortality was 10.15, 8.45 and 12.63% and for large sized egg class, it was 29.30, 20.14 and 33.41% respectively, on 0, 3 and 6 day of storage.

**Formatted:** Superscript

**Comment [Ma1]:** What is the conclusion and recommendation of the finding???

Key words: Fertility, mortality, breed, egg weight, Chabro

### **1. Introduction**

Poultry is currently India's fastest-growing agricultural sector (put the reference). The primary goal of modern hatcheries around the world is to produce high-quality chicks. To grow this sector, basic information on the influence of storage period and egg weight on fertility and mortality is required. Fertility in poultry refers to the percentage of incubated eggs that are fertile. Fertility tests are used to determine a bird's genetic and reproductive fitness, as well as to assess the economic efficiency of parent stocks. Fertility and mortality are influenced by a variety of environmental factors. Improper temperature and humidity,

male/female ratio, and prolonged storage period are the most common factors. only a few of them. Many research (King'ori et al., 2007, Yadav et al. 2021) have found that the length of storage has a significant impact on fertility and mortality. Egg storage for longer than 7 days results in a delay in embryo growth, a change in metabolic rate, and a decrease in fertility (Petek and Dikmen, 2006 and Yassin et al., 2008). As a consequence, precise knowledge of fertility in relation to egg weight and storage period can help in the selection of fertile eggs in a logical manner. As a result, the purpose of this study was to see how the storage duration affected the fertility of large, medium, and small sized eggs from the chabro breed of poultry.

**Comment [Ma2]:** Better to describe the breed in some sentence to be clear for the readers

## 2 Material and Method

### Description of the Study Area

The experiment was carried out in the Poultry Farm of the S.K.N. College of Agriculture at Jobner, Jaipur (Rajasthan, India). This region's climate is typically semi-arid, with temperature extremes in both summer and winter.

**Comment [Ma3]:** Should be clearly describe the area by stating longitude and latitude including the temperature and rainfall.....

### Experimental procedure

After collection of eggs, all the eggs were sampled and eggs with visible external abnormalities were screened out. In this process extra-large or very small eggs and abnormal eggs will be discarded. A total of 207 eggs were collected for this study. At commencement of the study, the eggs were weighed individually using sensitive weighing balance and later grouped into three egg size categories as follows: Small (38-44 g), medium (45-52 g) and large (53-59 g) eggs of different sizes. All experimental eggs were ~~To study the impact of incubation period, eggs were~~ stored at 21 °C temperature for three different time periods (fresh (0), 3 days and 6 days) and ~~-All experimental eggs were~~ incubated in an automated sanitized electrical incubator at 99.50 F- 99.75 F (37.5 °C ) with 60-65% relative humidity and turning hourly. Candling was done on 18<sup>th</sup> -18<sup>th</sup> days to determine infertile eggs and dead in germs. respectively. By using these data, fertility and hatchability were calculated for each strain as follow

**Comment [Ma4]:** What is the standard to say where the eggs are normal and abnormal???

**Comment [Ma5]:** This is different you stated in abstract 297 sample, why???

**Comment [Ma6]:** State the scale of the sensitive balance

**Formatted:** Superscript

**Comment [Ma7]:** Use mathematical equations

$$\text{Fertility rate} = \frac{\text{number of fertile eggs}}{\text{number of incubated eggs}} \times 100$$

### Hatchability=.

### Fertility of Eggs

The percentage of fertility was calculated on the basis of total eggs set:

$$\text{Fertility rate (\%)} = \frac{\text{No. of fertile eggs}}{\text{No. of egg incubated}} \times 100$$

*Mortality of Eggs*

Mortality rate was determined in each replicate by dividing the number of unfertilized egg to the total number of fertile eggs.

$$\text{Mortality} = \frac{\text{No. of unfertilized eggs}}{\text{Total no. of fertile eggs}} \times 100$$

*Statistical analysis*

Data obtained were subjected to statistical analysis using Completely Randomized Design (CRD). Consequently, a level of (P<0.01) was used as the criterion for statistical significance.

### 3 Result and Discussion

#### 3.1 Effect of storage period on fertility and hatchability

The effect of the storage period on egg fertility and hatchability is presented in Fig 1 and fig.2, respectively. It was observed that egg fertility was higher on 3<sup>rd</sup> day as compared to fresh 0<sup>th</sup> day for all egg weight classes but the percent increase was small (0.48-1.12%). In contrast to this, fertility was decreased on 6<sup>rd</sup> day as compared to 0<sup>th</sup> day for all egg weight classes and percent decrease was 3.7-5.4%. For small egg class, egg fertility was 95.55, 96.02 and 92.13% respectively, on 0, 3 and 6 day of storage. Similarly, for medium-sized egg class, fertility was 95.17, 95.64 and 90.28% and for large sized egg class, it was 91.11, 92.15 and 87.41% respectively, on 0, 3 and 6 days of storage. The fertility and hatchability were found to be decreased as storage duration increased. Petek et al. (2005) reported fertility of quail eggs at 1, 3, 5 and 7 days were 89.52, 92.93, 89.34 and 86.71, respectively, which show a decrease in fertility of eggs with long storage. A similar result was observed by Yadav et al (2021) in the Kuroiler breed of chicken.

**Comment [Ma8]:** Use mathematical equation

**Comment [Ma9]:** Poor way of writing and then should be re-write all M&M properly

**Comment [Ma10]:** Should be properly discussed, shallow comparison???




Fig 1: Effect of storage period on egg fertility in Chabro breed

### 3.2 Effect of storage period on mortality

The effect of storage period on mortality is given in Fig 2 and data revealed that embryonic mortality increased with increase in storage duration. Embryonic mortality was found higher on 6<sup>th</sup> day of storage as compared to 0 day and percent decrease was 12.3-19.6%. For small egg class, egg mortality was 20.95, 17.56 and 24.87% respectively, on 0, 3 and 6 day of storage. Similarly, for medium-sized egg class, mortality was 10.15, 8.45 and 12.63% and for large sized egg class, it was 29.30, 20.14 and 33.41% respectively, on 0, 3 and 6 day of storage. Fresh eggs had a significant rate of embryonic mortality, which could be attributed to internal egg causes. Alsobayel et al. (2017) found that embryonic mortality of Baladi eggs held for 5, 10, and 15 days was 16.40 %, 30.40 %, and 33.87 %, respectively, in their study. Reijrink et al. (2009) and Khan et al. (2014) found that embryonic mortality increases as storage time increases.




Fig 2: Effect of storage period on embryo mortality in Chabro breed

The regression equations and  $R^2$  of exponential, linear, polynomial, Logarithmic, polynomial and power models are represented in Table 1. The relationship between egg fertility and embryo mortality was highest for polynomial equation ( $R^2=0.47$ ) followed by Linear and logarithmic. Least correlation was observed for power equation with  $R^2=0.35$ .

Table 1: The regression equation between egg fertility and embryo mortality

Trend line name	Regression equation	$R^2$
Exponential	$y = 121730e^{-0.095x}$	0.36
Linear	$y = -1.8959x + 195.71$	0.44
Logarithmic	$y = -175\ln(x) + 812.49$	0.44
Polynomial (2 <sup>nd</sup> order)	$y = 0.1136x^2 - 22.828x + 1159.4$	0.47
Power	$y = 3E+18x^{-8.755}$	0.35

#### 4. Conclusion

An investigation was conducted to study the effect of storage period on fertility and mortality in large, medium and small sized eggs of Chabro breed. The egg fertility was higher on 3<sup>rd</sup> day of storage as compared to 0 day and decreased on 6 day of storage for all the egg sized classes. Similarly, mortality was also increased with the increase of storage duration. Overall mortality was lowest for medium sized egg class compared to small and large sized classes.

**Comment [Ma11]:** Please write properly the conclusion based on the main finding of the result including the your recommendation

**Formatted:** Highlight

**Formatted:** Highlight

## Reference

Ahmad, Mohd., Singh, B., Chauhan, S.S. and Singh, K.S. 2000. Effect of breed, egg size and shape on fertility, late embryonic mortality and hatchability. *Ind. J. Poult. Sci.* **35**(2). [ppp](#).

**Formatted:** Highlight

Alsobayel, A. A., & Albadry, M. A. (2012). Effect of age and sex ratio on fertility and hatchability of Baladi and Leghorn laying hens. *J. Anim. Plant Sci*, 22(1), 15-19.

**Comment [Ma12]:** Follow the author guide line of the journal, inconsistency way of writing and be sure that all references are sited and vis-

Khan, M. J., Khan, S. H., Bukhsh, A., & Amin, M. (2014). The effect of storage time on egg quality and hatchability characteristics of Rhode Island Red (RIR) hens. *Vet. arhiv*, 84(3), 291-303.

Kingori, A. M., Tuitoek, J. K., Muiruri, H. K., Wachira, A. M., & Birech, E. K. (2007). Protein intake of growing indigenous chickens on free-range and their response to supplementation. *International Journal of Poultry Science*, 6(9), 617-621.

Petek, M., & Dikmen, S. (2006). The effects of prestorage incubation and length of storage of broiler breeder eggs on hatchability and subsequent growth performance of progeny.

Petek, M., Başpınar, H., Oğan, M. M., & Balcı, F. M. (2005). Effects of egg weight and length of storage period on hatchability and subsequent laying performance of quail.

Reijrink, I. A. M., Meijerhof, R., Kemp, B., Graat, E. A. M., & Van den Brand, H. (2009). Influence of prestorage incubation on embryonic development, hatchability, and chick quality. *Poultry Science*, 88(12), 2649-2660.

Yadav, S., Datt, M., Choudhary, O. P., & Kumar, L. (2021). Effect of storage temperature and duration on fertility of Kuroiler breed of chicken.

Yassin, H. A. G. J., Velthuis, A. G., Boerjan, M., van Riel, J., & Huirne, R. B. (2008). Field study on broiler eggs hatchability. *Poultry Science*, 87(11), 2408-2417.