

STUDY OF THE PHYSICAL-CHEMICAL, FUNCTIONAL, GRANULOMETRIC, MINERAL AND ANTIOXIDANT PROPERTIES OF THREE (3) FLOURS FS, F40, F50 OF YOUNG SHOOTS OF ROAN (*BORASSUS AETHIOPUM* MART.) ACCORDING TO THE DRYING METHOD

Comment [h1]: Physicochemical

Comment [h2]: Marketed

Comment [h3]: No need

ABSTRACT :

The young shoot of roan (*Borassus aethiopum* Mart.) is cultivated and consumed in the cities of central Côte d'Ivoire such as Dimbokro. However, it has conservation problems, its nutritional value is not well known by the population and it is little processed by local industries. This study is a valorization of the young shoots of roan. In this study, the young shoots were dried respectively in the sun, at 40°C, at 50°C and then crushed to give respectively the FS, F40, F50 flours. The study of the physico-chemical properties gave, according to the flours, 6.47 to 7.17% for moisture; 5.71 to 6.07 for pH; 130.39 mg/100g on average for the reducing sugar content; 2.42 to 3.94% for total sugars; 0.53 to 1.07% for lipids; 6.23 to 7.57% for protein content; 2.19% on average for ash; 2.72 to 3.21% for fibre; 82.56 to 84.07% for total carbohydrates; 365.65 to 369.69 Kcal/100g for the energy value. The study of functional properties revealed 60.55 to 63.63% for dispersibility; 0.70 to 0.75 g/ml for tapped density; 143.08% on average for water absorption capacity; 118.67 to 128% for oil absorption capacity; 18.17 to 25.66% for foaming capacity. The granulometry showed that all three (3) flours are composed of grains whose size is lower than 1mm. In addition 62.7%, 52.9%, 47.8% of the grains of the FS, F40, F50 flours have sizes in the interval]250 µm, 125 µm]. The young shoots flours contain 243.45 to 280.44 mg/100g polyphenols; 68.46 to 83.03 mg/100g flavonoids; 45.47 to 59 mg/100g tannins. The mineral contents are 0.04 to 0.11 ppm for sodium; 0.02 ppm on average for phosphorus; 0.07 to 0.18 ppm for potassium; 0.11 ppm on average for calcium; 0.04 to 0.07 ppm for iron. The three (3) flours FS, F40, F50 of young shoots contain macronutrients, high energy value, fibre, interesting functional and granulometric properties, antioxidants, minerals that are beneficial for local populations and can be used by the food industries.

Comment [h4]: xxxxx

Comment [h5]: Please Rewrite the abstract according to guideline

Keywords: Young shoots, flour, *Borassus aethiopum*, physicochemical properties, functional properties

INTRODUCTION

The roan palm is a dioecious tree plant (Mollet, 1999) [1]. In Africa, it is called the sentinel of the savannah because of its size. It is a tall palm tree that can grow to an average height of 20 m or even 30 m, with a diameter of up to 1 m (Mollet, 1999) [1]. The natural range of the roan palm (*Borassus aethiopum* Mart.) is the semi-arid or sub-humid part of Africa (Mollet, 1999; Ali et al., 2010) [1,2]. It grows naturally from Senegal to the Central African Republic (Ali et al., 2010) [2]. In the central regions of Côte d'Ivoire (Toumodi, Dimbokro, Didiévi, ...), roan is very important for the population. In general, more than 88% of roan is used for the well-being of the population (Ouinsavi et al., 2011) [3]. It is used as food (fruit, sap, young shoots) (Waziri et al., 2010) [4], as building material (stipe, foliage, etc.) (Akinniyi et al., 2000; Sanon et al., 2007) [5,6]. It is also used in traditional African medicine (roots, male inflorescence) (Sanon et al., 2007; Waziri et al., 2011; Sakande et al., 2013) [6,7,8]. The fibres are used to make nets for fishing. The leaves are used to make a variety of objects: brushes, baskets, fences and roofs (Mollet, 1999) [1].

Comment [h6]: Referencing should follow in unique style

The female plants of the roan palm produce so-called fleshy fruits with stones, derived from the transformation of the sclerenchyma of the endocarp. These fruits are commonly called

drupes. A mature tree can produce about 200-300 drupes annually (Davis and Johnson, 1987) [9]. Some of the naturally fallen fruits remain in the forest for the regeneration of the tree and some are collected by the rural population for food and seedlings. Each fruit gives an average of three young shoots. They are obtained after 6 to 8 months by the rural population putting the ripe fruits in the ground (Ajanvon, 2019) [10]. The young shoot roan is a perishable commodity because it contains 52% moisture (Niamke, 2014) [11] which makes its consumption periodic and a considerable loss rate hence the need to transform it into flour. This alternative facilitates its storage and allows people to keep it longer and think of other uses.

Moreover, in central Côte d'Ivoire, the young shoots are sold on the local market at low prices to discerning consumers for food, which does not improve the living conditions of the farmers. Transforming them into flour with nutritional potential and for use in biscuit, pastry and bakery products would add value to the young shoots, which would benefit the farmers. In addition, farmers will turn away from the production of roan wine, which brings them money but destroys roan, which is an ecological tragedy. However, the method of drying the young shoots of roan is a very important operation for the good quality of the flour produced. This study is therefore a valorisation of the young shoots of roan. It will consist in determining for the three (3) flours:

- Physicochemical properties
- Functional properties
- Granulometry
- Antioxidants
- Minerals

Comment [h7]: spacing

1 MATERIALS AND METHODS

1.1 Materials

1.1.1. Young shoots of roan

The young shoots of roan studied come from Dimbokro in central Côte d'Ivoire. These young shoots are 6 to 8 months old.

1.2 Methods

1.2.1. Sampling

Samples of young shoots of roan were purchased in three (3) markets in the town of Dimbokro. For each market, 20 kg of saplings were purchased from three (3) women sellers, which makes 60 kg of saplings purchased per market. The total was 180 kg for all three markets. These samples of young shoots were taken to the laboratory for further analysis.

1.2.2. Production of young shoots flour

The young shoots flour was obtained in several steps. First, the tubers of young shoots were peeled, washed and cut into small pieces. These pieces were then divided into three (3) batches which were dried respectively in an oven at 40°C, at 50°C and in the sun for 4 days. Finally, the dried pieces of tubers from each batch were crushed and the crushed material was sieved using a 2mm diameter sieve. Thus, three (3) types of young shoots flour were obtained: F40, F50, and FS, respectively for oven drying at 40°C, 50°C and sun drying

1.2.3. Determination of physico-chemical properties

1.2.3.1 Moisture content

The method of moisture determination is that proposed by **AOAC (1990) [12]**. Moisture was assessed by drying 5 g of FS, F40, F50 flours in an oven (MEMMERT) at 105 °C for 24 h.

1.2.3.2 Determination of pH

The pH is determined according to the **AOAC (1990) method [12]**. Ten (10) grams of sample are diluted in 100 mL of distilled water for 1h. The solution obtained is filtered through filter paper (Whatman). The pH is measured directly by immersing the electrode of the pH meter (HANNA), previously calibrated, in the filtrate obtained.

1.2.3.3 Total and reducing sugar content

a. Extraction of ethanolsoluble sugars

Ethanol-soluble sugars are extracted according to the method of **Agbo et al (1986) [13]** as follows.

One (1) gram of flour is weighed and then diluted in 10 mL of ethanol (80%; v/v). To the resulting mixture are added 2 mL of zinc acetate (10%; w/v) and 2 mL of oxalic acid (10%, w/v). The mixture is then centrifuged at 3000 rpm for 10 min. The pellet is taken up with 10 mL ethanol (80%; v/v) and centrifuged again at 3000 rpm for 10 min. The supernatants are transferred to a 50 mL flask and the excess ethanol is evaporated in a sand bath for 10 min. The resulting solution is made up to 50 mL with distilled water.

b. Determination of total sugars

The total sugar content is determined according to the phenol-sulphuric method as described by **Dubois et al (1956) [14]**.

One hundred (100) µL of ethanol-soluble sugar extract is introduced into a test tube, then 0.9 mL of distilled water, 1 mL of 5% (w/v) phenol and 5 mL of concentrated sulphuric acid are added successively. After shaking and cooling the tube, the absorbance is read with a spectrophotometer (PG INSTRUMENTS) at 490 nm against a blank. The determination of the quantity of total sugars is carried out using a standard range of glucose stock solution at 1 mg/mL carried out under the same conditions as the test.

c. Determination of reducing sugars

The quantification of reducing sugars is performed according to the method of **Bernfeld (1955) [15]**.

One (1) mL of ethanosoluble extract of sugars is introduced into a test tube. To the contents of this tube, 0.5 mL of distilled water and 0.5 mL of DNS solution are added. The whole mixture is heated in a boiling water bath for 5 min. After cooling, 2 mL of distilled water is added and the absorbance of the solution is read with a spectrophotometer (PG INSTRUMENTS) at 540 nm against a blank. A standard range established under the same conditions as the test from a stock solution of glucose (1 mg/mL) is used to determine the quantity of reducing sugars.

1.2.3.4. Lipid content

Lipids were quantified from 10 g of each flour by Soxhlet extraction using 300 mL n-hexane for 7 h (AFNOR, 1986) [16]. The resulting hexane-oil mixture was recovered and separated with a rotavapor (Heidolph). The flask, initially tared and containing the oil, was weighed to determine the mass of oil extracted.

1.2.3.5. Protein content

Crude protein was determined by total nitrogen determination according to the Kjeldhal method (AOAC, 1990) [12]. Thus 1g of each flour was mineralised in the presence of Kjeldahl catalysts (potassium sulphate K₂SO₄ and copper sulphate and concentrated sulphuric acid H₂SO₄). The mineralisate was purified by distillation. Nitrogen was then quantified by titration with 0.1 N H₂SO₄. The crude protein content of the flours was deduced from the nitrogen content using 6.25 as a conversion factor.

1.2.3.6. Ash content

The ash content (total mineral matter) was determined according to the method described by AOAC (1990) [12] by incinerating five (5) grams of each flour in a muffle furnace (PYROLABO) at 550°C for 12 h.

1.2.3.7. Fibers content

The determination of the fibers content was carried out according to the method of Wolff (1968) [17]. The determination of the crude fibers content consisted in treatment of 2 g of FS, F40, F50 flours sample with 50 mL of 0.25 N sulfuric acid and 50 mL of 0.31 N sodium hydroxide and filtration of the resulting solution upon Whatman paper. The residue was dried for 8 h at 105 °C then incinerated at 550 °C for 3 h into ovens. The final residue was weighed as crude fibers and expressed in percentage.

1.2.3.8. Total carbohydrates content and energy value

Total carbohydrates and energy values were determined using calculation formulas recommended by FAO (2002) [18] accounting the moisture, fat, protein, ash contents and the energy coefficients for macromolecules. $TCC (\%) = 100 - [P(\%) + M(\%) + F(\%) + A(\%)]$
 $CEV \text{ (kcal/100g)} = [(4 \times P) + (9 \times F) + (4 \times C)]$
With: TCC, total carbohydrates content; CEV, caloric energy value; P, protein content; M, moisture content; F, fat content; A, ash content; C, total carbohydrates content

1.2.4. Determination of functional properties

1.2.4.1. Dispersibility of the flours

The dispersibility of the flours was determined according to the (modified) technique described by **Kulkarni et al (1991) [19]**. A volume of 10 mL of distilled water was added to 1 g of flour in a graduated cylinder. The mixture was stirred thoroughly with a rod for 2 min. The dispersibility of the flour is the difference between the total volume (V_0) of the particles just after manual agitation and the volume (V_t) of the deposited particles recorded at time t (min).

$$\text{Dispersibility (\%)} = \frac{(V_0 - V_t)}{V_0} \times 100$$

1.2.4.2. Tapped density

The tapped density (DT) of the flours was determined using the (modified) technique of **Oladele and Aina (2007) [20]**. A quantity of 50 g of flour (ME) was placed in a 100 mL graduated cylinder. The test tube was then gently tapped on the bench until a constant volume V_t

$$\text{DT (g/ml)} = \frac{\text{ME}}{V_t}$$

1.2.4.3 Water absorption capacity (WAC)

The water absorption capacity (WAC) of the flours was determined using the (modified) techniques of **Phillips et al. (1998) [21]**. Exactly 1 g of each flour (M_0) was dissolved in 10 mL of distilled water in a centrifuge tube. This mixture was stirred for 30 min by a shaker and then kept in a water bath at 37 °C for 30 min. It was then centrifuged at 14674 xg for 15 min in an ORTO ALRESAR centrifuge. The resulting pellet (M_2) was weighed and then dried at 105 °C to a constant mass (M_1). The WAC was calculated from the following relationships:

$$\text{WAC (\%)} = \frac{(M_2 - M_1)}{M_1} \times 100$$

1.2.4.4. Oil absorption capacity

The oil absorption capacity of the flours was determined using the (modified) technique of **Eke and Akobundu (1993) [22]**. A 1 g sample (M_0) of flour was dissolved in 10 mL of oil. The mixture was stirred for 30 min at room temperature using a magnetic stirrer and then

centrifuged at 11886 xg for 10 min in an ORTO ALRESAR centrifuge. The recovered pellet was weighed (M1). The oil absorption capacity (OAC) was calculated from the following formula:

$$\text{OAC (\%)} = \frac{(M_1 - M_0)}{M_0} \times 100$$

1.2.4.5. Foaming capacity

The foaming capacity (FC) of the flours was determined using the (modified) technique of **Coffman and Garcia (1977) [23]**. Three grams of flour were placed in a 50 mL graduated cylinder that had been oven dried at 50 °C. Then 30 mL of distilled water was added to the sample to facilitate dispersion of the flour in the test tube and the volume was noted (volume before homogenisation). The test tube was then vigorously shaken by hand and the new volume was read off the test tube (volume after homogenisation). The volume of the foam obtained was calculated as the difference between the volume after homogenisation (V after) and the volume before homogenisation (V before). The foaming capacity (FC) was calculated from the following formula:

$$\text{FC (\%)} = \frac{V_{\text{after}} - V_{\text{before}}}{V_{\text{before}}} \times 100$$

1.2.5. Determination of flour grain size

The particle size of the flours is estimated by fractionating the total mass of flours through a series of sieves of decreasing mesh size (2mm, 1mm, 500um, 250um, 125um and 63um). Underneath the 63um sieve is a collection lid.

The sample is placed in a sieve with different mesh sizes and closed with a lid. The sieve is stirred for 30 minutes.

The grains are deposited one after the other according to their diameter size and the rejects and passings of each sieve are weighed with a technical balance (Denver instrument SI-4002) of precision.

1.2.6. Determination of antioxidants

1.2.6.1. Extraction of phenolic compounds

Phenolic compounds are extracted with methanol by the method of **Singleton et al (1999) [24]**.

One (1) gram of sample of FS, F40, F50 flours is homogenised in 10mL of 70% (v/v) methanol. The resulting mixture is centrifuged at 1000 rpm for 10 min. The pellet is recovered in 10 mL of 70% (v/v) methanol and centrifuged again. The supernatants are collected in a 50 mL flask and made up to the mark with distilled water.

1.2.6.2. Determination of total phenols

The method of **Singleton et al (1999) [24]** is used for the determination of total phenols. One (1) mL of methanolic extract is introduced into a test tube. To the contents of the tube is added 1mL of Folin-ciocalteu reagent. The tube is left to stand for 3min and then 1mL of 20% (w/v) sodium carbonate solution is added. The contents of the tube are made up to 10 mL with distilled water. The tube is placed in the dark for 30 min and the OD reading is taken at 745 nm against a blank. The amount of phenol in the sample is determined by a standard range using a stock solution of gallic acid (1 mg/mL) under the same conditions as the test.

1.2.6.3. Determination of tannins

The determination of tannins was carried out according to the method described by **Bainbridge et al. (1996) [25]**.

One (1) mL of methanolic extract is introduced into a test tube. To the contents of the tube is added 5 mL of vanillin reagent. The tube is left to stand for 20 min in the dark and the optical density (OD) is read at 500 nm against a blank. The amount of tannin in the samples is determined using a standard solution of tannic acid (2 mg/mL) under the same conditions as the test.

1.2.6.4. Determination of flavonoids

The flavonoid assay was performed as described by **Meda et al. (2005) [26]**.

A volume of 0.5 mL of methanolic extract was introduced into a test tube. To the contents of the tube are successively added 0.5 mL of distilled water, 0.5 mL of 10% aluminium chloride, 0.5 mL of 1M potassium acetate and 2 mL of distilled water. The tube is allowed to stand for 20 min in the dark and the optical density (OD) is read at 415 nm against a blank. The amount of flavonoids in the sample is determined using a standard solution of quercetin (0.1 mg/mL) under the same conditions as the test.

1.2.7. Determination of minerals

The determination of the minerals was carried out by atomic absorption with an air-acetylene flame AAS 20 type VARIAN.

The FS, F40, F50 flours were ground to a particle size of 0.1mm. A mass of 0.3 g of each flour is calcined at 600°C for 5h in an oven until a white ash is obtained. After cooling, 5 mL of 1N nitric acid was added and evaporated to dryness on a sand bath. To the residue is added 5mL of 1N hydrochloric acid and the whole is fired again at 400°C for 30 min. Once the calcined product is recovered from the furnace, 10mL of 0.1N hydrochloric acid is added to the crucible to recover the product. The resulting mixture is poured directly into a 50 ml

volumetric flask. The operation (washing the crucible with 10 ml of 0.1 ml HCL) is repeated three times and the flask is filled to the mark. Allow to decant and take the supernatant for filtration with 0.45 µm wattman paper or with a 0.36 syringe filter. The elements contained in the solution are then determined by AAS.

NB: To avoid interference from the elements Ca, K, 5 ml of lanthane chloride is added.

1.2.8. Statistical study

All statistical analyses were carried out with R software. The physicochemical and functional properties, the determination of antioxidants and minerals of the three flours were compared using the ANOVA test. The null hypothesis (H0) is that all the means are equal and that the properties are equal between the 3 flours and for the alternative hypothesis (H1) there is a significant difference between the 3 flours. The significance level was set at 0.05 for all analyses.

To perform the ANOVA test, the Shapiro-Wilk and Bartlett tests were performed a posteriori to test normality and equality of variances respectively. When the conditions of normality and/or equality of variances of certain parameters were not met, the non-parametric Kruskal-Wallis test was applied.

3. RESULTS AND DISCUSSION

3.1. Physico-chemical properties (Table I)

The results of the physicochemical analyses of the different flours, show that the moisture content of FS flour (7.17%) is higher than that of F40 flour (7.1%) and F50 flour (6.47%). These levels are lower than that of flour from the uncooked pulp of breadfruit (*Artocarpus altilis*) which is $10.7\% \pm 0.36$ (Oulai, 2014) [27]. These moisture contents are below the maximum level of 15.5% defined by the Codex Alimentarius Commission (CAC, 1985) [28]. Indeed, when the moisture content is high, aggregation of the flour particles occurs, thus reducing its quality and functionality (Aguilera et al., 1995) [29]. Another advantage of having a low moisture content lies in the technological uses. Indeed, with a moisture content below 12%, both untreated and treated flours are favourable for long-term preservation. Microbiologically, these low moisture levels limit the growth of microorganisms, with the exception of moulds (Aryee et al., 2006) [30].

FS flour has a pH of 5.71. It is more acidic than F40 flour with a pH of 6.07 and F50 flour with a pH of 5.95. The pH varies very little and remains below 7. This result is comparable with Yetunde et al, (2009) [31] and Alozie et al, (2009) [32]. The pH is a sign of the acidity or alkalinity of the flour and greatly affects its performance during its use in the food system. When a flour has a pH < 4 this flour is said to be very acidic which denotes high fermentation and consequently high degradation of the starch present, so this type of flour would not be suitable for bread making (Tortoe et al., 2017) [33].

The three (3) flours have a total sugar content that varies from 2.42% to 3.94% and a reducing sugar content that varies from 117.35 mg/100g to 152 mg/100g. These levels are higher than those observed by Gutap and Nagar (2008) [34]. These authors obtained total sugar levels between 0.05 and 0.4% in uncooked and cooked soybean meal and also reducing sugar levels between 0.03 and 0.04%. This difference would be due to the fact that the young shoot of the roast tree belongs to the starch group and is therefore rich in starch (Niamke, 2014) [11] and consequently rich in sugars. It should be noted that flours rich in sugars are useful for the manufacture of certain foods such as cakes, biscuits, cakes.

The lipid content of F50 flour (1.07%) is higher than that of F40 (0.60%) and FS (0.53%) flours. These levels are close to those obtained by **Oulai, 2014 [27]** (0.84% to 0.48%) when cooking *Artocarpus altilis* pulp. According to **Anses in 2021[35]**, lipids play two (2) major roles: an energy storage role and a structural role (enter into the composition of cells).

The protein percentage of F50 flour (7.57%) is higher than that of F40 (6.65%) and FS (6.23%) flours. These contents are close to that of unripe banana flour (6.57%) (**Anuonye et al., 2012 [36]**). Proteins are essential to the body, they play a structural role (in muscles and even skin) but are also involved in a large number of processes such as the immune response (antibodies), oxygen transport in the body (haemoglobin) or digestion (digestive enzymes) (**Anses, 2021 [35]**).

The fibre content of F50 flour (3.21%) is higher than that of F40 (2.96%) and FS (2.72%). These three (3) flours could be a significant source of dietary fibre which is eliminated more slowly from the stomach and thus improves intestinal transit. These dietary fibres are absolutely essential for the balance of the digestive tract and the body. It is a factor in good health. Studies have shown an inverse correlation between dietary fibre consumption and colon cancer. This is because fibre has the ability to complex with carcinogenic molecules, thus preventing their contact with the colon and facilitating their excretion (**Jansen et al., 1999; Chene, 2003 [37,38]**). Consumption of prepared flours could therefore increase gastric volume and provide a post-ingestive state to reach a state of satiety more quickly (**Chene, 2003; Al-Dobaib, 2009 [38,39]**). Fibre generally reduces blood glucose, HDL-cholesterol, LDL-cholesterol and thus contributes to the reduction of coronary heart disease (**Jalili et al., 2000 [40]**).

The percentage of total carbohydrates in FS flour (84.07%) is higher than in F40 (83.41%) and F50 (82.56%) flours. These total carbohydrate levels are higher than those of millet (71.35% to 77.13%) (**Ocheme et al., 2010 [41]**). FS, F40, F50 flours are rich in carbohydrates, which are compounds that provide energy for the functioning and maintenance of muscle cells, brain, red blood cells and other organs, etc. (**Martin, 2000; Folin, 2005; Fredot, 2009 [42, 43, 44]**). These flours could be an important alternative source of calories in rural areas in case of shortage of high energy foods (Attiéké, Yam, banana etc.) commonly consumed by the population. Moreover, the transformation of young shoots into flour offers a definite advantage because of its better conservation (for the population), as this avoids its rapid degradation.

The energy values of FS, F40, F50 flours vary from 366 to 369.69Kcal/100g. This energy value is due to the high total carbohydrate content. This value is close to that of *Dioscorea alata* yam raw and cooked for 90 min (357.65 and 370.01 Kcal/100g respectively) (**Ezeocha and Ojmelukwe, 2012 [45]**). FS, F40, F50 flours could be used partly as energy flour in porridges for infants and children whose energy requirements vary from 547 to 1092 Kcal/day (**Butte, 1996 [46]**).

Table I : Physico-chemical properties of FS, F40, F50 flours

Parameters	FS	F40	F50	General Average	P-value
Moisture (%)	7.17±0.15 ^c	7.1±0.17 ^b	6.47±0.15 ^a		0.003
pH	5.71±0.01 ^a	6.07±0.01 ^c	5.95±0.01 ^b		0.03
Reducing sugars (mg/100g)	117.35±6.2 ^a	121.19±1.4 ^a	152.63±21.57 ^a	130.39	0.06
Total sugars (%)	2.42±0.22 ^a	3.36±0.44 ^b	3.94±0.32 ^c		0.004

Lipids (%)	0.53±0.11 ^a	0.60±0.2 ^b	1.07±0.11 ^c		0.009
Proteins (%)	6.23±0.04 ^a	6.65±0.35 ^b	7.57±0.02 ^c		0.03
Ash (%)	2±0.2 ^a	2.24±0.06 ^a	2.34±0.15 ^a	2.19	0.08
Fibers (%)	2.72±0.05 ^a	2.96±0.1 ^b	3.21±0.01 ^c		0.0004
Total carbohydrate (%)	84.07±0.2 ^c	83.41±0.6 ^b	82.56±0.4 ^a		0.01
Energy content(Kcal/100g)	366±0.35 ^a	365.65±1.14 ^a	369.69±0.1 ^a	367.11	0.06

Per line, values followed by different superscript letters are statistically different at 5%. P-value: value of the statistical probability test. With a < b<c; P value < 0.05 (5%) so the difference is significant

3.2. Functional properties of FS, F40, F50 flours (Table II)

The dispersibility (after 30 min) of F50 flour (63.63%) is higher than that of F40 (61.83%) and FS (60.55%) flours. The dispersibility of a flour, which is an indicator of its reconstitution power in water, is a useful functional parameter in the formulation of various food products (Mora-Escobedo *et al.*, 1991) [47]. The dispersibility percentage of the three flours is similar to that of local Nigerian rice (56-66%) (Eke-Ojiofor *et al.*, 2011) [48]. The higher the dispersibility percentage, the greater the ability of the flour to reconstitute in water to give a fine, coherent paste

The tapped (bulk) density of F50 flour (0.75 g/ml) is higher than that of F40 (0.73g/ml) and FS (0.70g/ml) flours.) All these values are close to that of wheat flour (0.80 g/cm³) (Ijarotimi, 2012) [49]. Density is a very important parameter in that it determines the packaging and transport conditions of the food product (Shittu *et al.*, 2005) [50]. Nutritionally, a low packed density promotes the digestibility of food products, especially in children because of their immature digestive systems (Nelson-Quartey *et al.*, 2007) [51]

The water absorption capacity of F50 flour (146.58%) is higher than that of F40 (143.70%) and FS (138.95%) flours. These values are lower than those of uncooked and cooked rice (225% and 250% respectively) (Abulude, 2004) [52]. Furthermore, the use of flours as food ingredients depends, to a large extent, on their interaction with water. The water absorption capacity of flours plays an important role in the food preparation process as it predicts the ability of the flour to absorb water under conditions where water is in short supply. A high capacity allows more water to be added to the dough, thus improving its workability. Furthermore, water absorption capacity is an essential property of doughs and bakery products as it allows for thickening and increasing the viscosity of foods (Falade and Okafor, 2015) [53]. The high water absorption capacity of seedling flour could reflect the presence of high amounts of hydrophilic substances capable of improving the viscosity of various food products (Grah *et al.*, 2014) [54]. This could also reflect a greater interaction between proteins and water in the formed system. Furthermore, the type of protein such as polar proteins would also increase this ability (Grah *et al.*, 2014; Tortoe *et al.*, 2017) [54,33].

Oil is more absorbed by F50 flour with a percentage of oil absorption of 128% higher than F40 (119.67%) and FS (118.67%) flours. These oil absorption capacities are lower than that of yam flour (190%) (Ukpabi, 2010) [55]. Oil absorption capacity is an important characteristic in fatty food formulations as it is believed to act as a flavour retention device and mouthfeel enhancer (Yadahally, 2011) [56]. This oil absorption capacity of flour is due to the existing interactions between the side chain of non-polar amino acids and the hydrocarbon chains of lipids (Shindhu *et al.*, 2016) [57]. Since these flours have a high oil absorption capacity, they could be a good lipophilic component and therefore suitable for the preparation of sausages, soups and cakes (Aremu *et al.*, 2006) [58].

The foaming capacity of F50 flour (25.66%) is higher than that of F40 (23.43%) and FS (18.17%) flours. The foaming capacity improves the texture, uniformity and appearance of the food (Akubor, 2007) [59]. According to Yasumatsu et al. 1972 [60], foam formation depends on pH, viscosity, protein and processing methods. The foaming capacity of these three flours is higher than that of cassava flour (13.70%) (Ubbor and Akobundu, 2009) [61].

Table II : Functional properties of FS, F40, F50 flours

Parameters	FS	F40	F50	General Average	P-value
Dispersibility of flours after 30 min (%)	60.55±3.04 ^a	61.83±0.11 ^a	63.63±0.006 ^a	62	0.06
Tapped density (g/ml)	0.70 ^a	0.73 ^b	0.75 ^c		0.02
water absorption capacity (%)	138.95±0.09 ^a	143.70±4.8 ^a	146.58±0.37 ^a	143.08	0.2
Oil absorption capacity (%)	118.67±4.72 ^a	119.67±3.05 ^b	128±2.64 ^c		0.03
The foaming capacity (%)	18.17±0.01 ^a	23.43±0.76 ^b	25.66±1.52 ^c		0.03

Per line, values followed by different superscript letters are statistically different at 5%. P-value: value of the statistical probability test. With a < b<c; P value < 0.05 (5%) so the difference is significant

3.3. Particle size of FS, F40, F50 flours (Table III)

The results of the particle size classification made it possible to separate seven characteristic fractions of the three flours expressed as a percentage in Table III. All three (3) flours are composed of grains smaller than 1mm. In addition, 98.8% of the grains of the FS and F40 flours and 98.9% of the grains of the F50 flour are smaller than 500 µm. These results are almost identical to those of Djidohokpin (2012) [62] for fonio-soya composite flours. Our flour could be recommended after formulation as infant flour because according to the standard established on the granulometry of infant flours by the Advisory Committee of the PAAN project (Projet d'Appui aux Activités de Nutrition) which stipulates that infant flour should not contain particles larger than 500 micrometers (Tchibindat and Trèche, 1995) [63]. Also, 62.7%; 52.9%; 47.8% of the grains of FS, F40, F50 flours have sizes in the range [250 µm, 125µm]. For Djidohokpin (2012) [62], the highest retention rate for fonio-soya composite flours is obtained at the 180µm sieve level while for the three flours FS, F40, F50, the highest retention rate is obtained at the 125µm sieve level; this means that the roast flours are finer than the fonio-soya composite flours. The granulometry of flours is of major importance for their analysis and use. Indeed, it allows to detect the presence of foreign particles and to pronounce on milling problems. It plays a fundamental role in hydration, which is the operation of bread-making and the preparation of the dough. It also makes it possible to predict its behaviour during hydration. In baking, the amount of water absorbed during dough fermentation, as well as the rate of water absorption increases with the fineness of the flour particles (Feillet, 2000) [64].

Table III : Particle size of FS, F40, F50 flours

Flours Particle size	FS	F40	F50
< 2mm	100%	100%	100%
<] 2mm ,1mm]	100%	100%	100%
] 1mm, 500µm]	1,2%	1,2%	1,1%
] 500 µm, 250µm]	18,9%	15,3%	18,6%
] 250 µm, 125µm]	62,7%	52,9%	47,8%
] 125 µm, 63µm]	9,7%	15,2%	15,8%
<63µm	7,5%	15,4%	16,7%

3.4. Antioxidants (Table IV)

The presence of antioxidants can reflect a response to stress (scarcity of rainfall, unfavourable soil quality, which are associated with an increase in tannin levels) (Mebirouk-Boudechiche *et al.*, 2014) [65]. Thus, depending on the efforts made by the plant to adapt to environmental conditions, the amount of antioxidant decreases or increases. Flavonoids can neutralise free radicals and reduce cancer risk by stopping cell growth in tumours (Wang *et al.*, 2014) [66].

Current literature suggests that long-term consumption of a polyphenol-rich diet protects against certain cancers, cardiovascular disease, type 2 diabetes, osteoporosis, pancreatitis, gastrointestinal problems, lung damage and neurodegenerative diseases (Fuujiki *et al.*, 2015; Xiao *et al.*, 2015; Martin- Pelaez *et al.*, 2013; Fraga *et al.*, 2010) [67,68,69,70].

Table IV : Antioxidants of FS, F40, F50 flours

Parameters		FS	F40	F50	P-value
Antioxidants	Polyphenols (mg/100g)	243,45±0,68 ^a	267,25±0.67 ^b	280,44±0.93 ^c	0.02
	Flavonoids (mg/100g)	68,46±0.48 ^a	75,15±0.086 ^b	83,03±0.16 ^c	4.49*10 ⁻⁹

	Tannins (mg/100g)	45,47±0.69 ^a	54,13±0.26 ^b	59±0.73 ^c	4.56*10 ⁻⁷
--	----------------------	-------------------------	-------------------------	----------------------	-----------------------

Per line, values followed by different superscript letters are statistically different at 5%. P-value: value of the statistical probability test. With a < b<c; P value < 0.05 (5%) so the difference is significant

3.5. Minerals (Table V)

The results of this study showed the presence of minerals in young shoots flours. Some of them were affected by the drying temperature. Indeed, the mineral contents increased with temperature. This increase in the mineral content of the young shoots flours would be due to the effect of the heat. This can be explained by the fact that certain anti-nutritional factors interfere with the availability of minerals by complexing them in their structure, as suggested by **Alonso et al., (2001) [71]** and **Anigo et al., (2009) [72]**. The degradation of these anti-nutritional factors by heat releases these minerals into the matrix (**Ewald et al., 1999) [73]**. The sodium content of roasted sapling flours increases with temperature (0.04 to 0.11 ppm). This result is lower than that reported for breadfruit (*Artocarpus altilis*) from Ghana (690ppm DM) (**Polycarp et al., 2012) [74]**. Sodium plays an important physiological role in humans at several levels as it participates in the control of the volume of the extracellular medium (which refers to the fluid balance) of the body, maintenance of the electrochemical gradient of cells, transmission of nerve impulses, muscle contraction, intestinal absorption of some nutrients (**Alexandra Retion, 2022) [75]**. With its low sodium content, the flour of young roasted shoots could be used as food without fear of health risks for people with high blood pressure.

Potassium is a mineral that increases cardiovascular well-being and is recommended for the prevention of certain complications of myocardial infarction (**Kannel, 1997; Chow, 2009) [76,77]**. Its content varies from 0.07 to 0.18 ppm. This content is much lower than that of *Artocarpus altilis* flour (7045 to 8706 ppm) (**Oulai, 2014) [27]**. Regarding phosphorus, the results showed that the content which was 0.02 ppm for FS flour did not increase for F40 (0.02 ppm) and F50 (0.02 ppm) flours. These levels are lower than that of uncooked *Colocasia esculenta* (208 ppm DM) (**Lewu et al., 2010) [78]**. The calcium content of the three flours also did not vary with temperature (0.11 to 0.12 ppm). These levels are lower than those of cooked *Hibiscus sabdariffa* seeds (740 ppm) (**Yagoub et al., 2008) [79]**. A diet rich in calcium and phosphorus is a factor in the prevention of osteoporosis and also a factor in reducing the risk of high blood pressure, colon and prostate cancer (**Bonithon-Kopp et al., 2000) [80]**. Iron plays an important role in the human body. The haemoglobin present in the red blood cells absorbs the 70% of iron consumed. This allows oxygen to function properly. This oxygen is then transmitted to the cells. Iron is also found in the myoglobin of the muscles, which enables air to be stored. The remaining 30% of iron plays a role in activating the body's metabolisms. It contributes greatly to the production of energy and the activation of the immune system (**Eric Favre, 2022) [81]**. Iron deficiency anaemia affects one third of the world's population. However, excessive iron intake causes colorectal cancer (**Senesse et al., 2004) [82]**. The iron contents of our flours (0.04 to 0.06 ppm) are lower than that of sweet potato flour (10.97 ppm) revealed by **Ofori et al., 2016) [83]**. The iron values of the three flours are below the limit of 15 ppm set by WHO as the limit of iron in food (**WHO 1982) [84]**.

Table V : Minerals of FS, F40, F50 flours

Parameters	FS	F40	F50	General average	P-value
Sodium (ppm)	0.04±0.001 ^a	0.07±0.004 ^b	0.11±0.004 ^c		0,000
Phosphorus (ppm)	0.02±0.005 ^a	0.02±0.006 ^a	0.02±0.003 ^a	0.02	0,892
Potassium (ppm)	0.07±0.009 ^a	0.12±0.004 ^b	0.18±0.001 ^c		0,000
Calcium (ppm)	0.11±0.007 ^a	0.11±0.006 ^a	0.12±0.003 ^a	0.11	0,082
Iron (ppm)	0.04±0.006^a	0.06±0.006^b	0.07±0.003^b		0,002

Per line, values followed by different superscript letters are statistically different at 5%. P-value: value of the statistical probability test. With a < b<c; P value < 0.05 (5%) so the difference is significant

CONCLUSION AND PERSPECTIVES

With the exception of moisture, which is high for FS flour, and pH, which is higher for F40 flour, all other parameters increase in the flour with increasing temperature. This is because sun-drying exposes the flax to moisture in the air; the higher the temperature, the more concentrated the various parameters are in the flour. The young shoot of roan is a tuber, which makes it a potential staple food that could be used in the diet to combat hunger and to provide food security. The results of this study prove that the three (3) flours are a good source of carbohydrate and could therefore be useful for energy purposes. The low moisture content of the three (3) flours (< 8%) would allow the flours to be stored for a long time. All flours contain fibre and minerals that are beneficial to the health of the population. The functional properties of the flours, i.e. dispersibility, packed density, water absorption capacity, oil absorption capacity and foaming capacity, suggest that the flours from young shoots would be suitable for use in food formulations (infant porridges, pastries, cakes, etc.) for which these different properties are required

In perspective, it would be important to carry out the following research

- Study the different properties of flours made from roasted sapling flour with other flours.
- To propose a feed formulation

REFERENCES

- [1]Mollet M. (1999). L'utilisation durable des palmiers *Borassus aethiopum*, *Elaeis guineensis* et *Raphia hookeri* pour l'extraction de vin de palme en Côte d'Ivoire. Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ) GmbH Postfach 5180, D-65726 Eschborn République fédérale d'Allemagne.
- [2] Ali A., Alhadji D., Tchiegang C. & Saïdou C., 2010. Physico-chemical properties of palmyra palm (*Borassus aethiopum* mart.) fruits from Northern Cameroon. *Afri. J. Foods Sci.* 4,115-119.
- [3] Ouinsavie C., Gbemevo C., Sepko N., (2011). Ecological structure and fruit production of Africa fan palm (of *Borassus aethiopium*) population. *Amer. J. plant sci.* 2. pp 733-743.
- [4] Waziri, M.; Akinniyi, J.A. ; Salako, A.A., 2010. Toxicity of Acetone Extract of Muruchi, the shoot of *Borassus aethiopum* mart. *Eur. J. Sci. Res.* 4 1, 6-12.
- [5] Akinniyi, JA., Waziri,M. and Usman, H.S. (2000). study of the androgen contenty of barassusaethiopum mart. Unimaid research report. pp.20-28.

Comment [h8]: Please give a concrete message to consumer on the basis of presence of mineral

- [6] **Sanon, M.D. & Sakande M. (2007)**. *Borassus aethiopicum* Mart. Release Seed leaflet no. 120. Forest and Landscape, Horsholm, Denmark. Available at [http:// en.sl.life.ku.dk / upload / 120net.pdf](http://en.sl.life.ku.dk/upload/120net.pdf)
- [7] **Waziri, M.; Akinniyi, J.A. & Chide, M.A., 2011**. Elemental composition of Dalang . A food condiment from evaporated extract of *Borassus aethiopicum* fruit ash. *Am. J. Food. Nutr.*, 1 : 123-125.
- [8] **Sakande J., kaba E., lampo M., Pale E., Niamké J.B., Noucouлма D.G., sawadogo M., et Guissore J.P. (2013)**. Anty-inflammatory and oxidant activities of a fraction 11 of male inflorescence of *borassus aethiopicum* Mart (Arecacea), *Amer. J. biochem. Mol. Bio*, 3. pp 101-109.
- [9] **Davis T.A. et Johnson D. V (1987)**. Current utilization and further development of the palmyrah palm (*Borassus flabellifer* L) in tadu nade state indai *Eco. Bot.* 41(2) 247-266
- [10] **Ajavon A. Y. C. (2019)**. Importances socio-economiques de la production des hypocotyles du ronier (*borassus aethiopicum* mart.) dans la commune de save au centre du benin (afrique de l'ouest) statisticien economiste. *Revue ivoirienne de geographie numeros special janvier 2019 ; ISSN*, pp. 2125-2125.
- [11] **Niamke A.M. (2014)**. Contribution de la valorisation de la jeune pousse de ronier (*Borassus Aethiopicum* Mart.) de Didievi : Propriété physico-chimiques, hydrolyse enzymatique et acide de la farine, de la pâte fraîche et de l'amidon. Thèse de doctorat, Cote D'Ivoire. 168p
- [12] **AOAC (1990)**. Official methods of analysis of the AOAC, 15th Edition, Methods 932.06, 925.09, 985.29, 923.03. Association of official analytical chemists. Arlington, VA, USA
- [13] **Agbo NG, Uebersax M, Hosfield G. (1986)**. An efficient extraction technique of sugars from dry edible beans (*Phaseolus vulgaris*) and estimation in HPLC, National University, Côte d'Ivoire. *Annals series C (sciences) Tome XXI*; pp.169-184
- [14] **Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. (1956)**. Colorimetric method for determination of sugar and related substances. *Anal. Chem.*; 28: pp.350-356.
- [15] **Bernfeld P. (1955)**. Amylase bêta and alpha (assay method), in *methods in enzymology*, Ed., Academic Press, New-York, USA.; pp.149-154.
- [16] **AFNOR. (1986)**. Collection of French standards. Quality control of dairy products. AFNOR, Paris – la defense; 1986
- [17] **Wolff JP. (1968)** Fat analysis manual. Azoulayed. Paris (France).;519.
- [18] **FAO (2002)**. Report of the International Rice commission. Vingtième session, 23-26 Juillet 2002, Bangkok, Thaïlande.
- [19] **Kulkani K. D., Kulkari D. N. et Ingle U.M. (1991)**. Sorghum malt-based weaning formulation. Preparation, functional properties, and nutritive value. *Food and nutrition bulletin* 13 : pp. 322-327.
- [20] **Oladele A.K. et Aina J.O. (2007)**. Chemical composition and functional properties of winged bean flours produced from two varieties of tigernut (*Cyperus exculentus*). *African journal of biotechnology* 6: pp.2473-2476.

- [21] **Phillips R.D., Chinman M.S, Branch A.I., Miller J. et Mewatters K.H. (1998).** Effect of pre-treatment on functional and nutritional properties of cowpea meal journal of food sciences 53: 805-809
- [22] **Eke O.S. et Okobundu E.N.T. (1993).** Functional properties of african yam bean (*sphenostylis stenocarpa*) seed flour as affected by processing. Food chemical 48: pp.337-340
- [23] **Coffman C.W.L., et Garcia V.C. (1977).** Rheology of soy protein and amino acid of a protein isolate from mungbean gelation. Cereal chemistry 41: pp.473-484
- [24] **Singleton VL, Orthofer R & Lamuala-RRM (1999).** Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. Methods Enzymol. 299: 152-178.
- [25] **Bainbridge Z, Tomlins K, Willings K & Westby A (1996).** Methods for assessing quality characteristics of non-grain starch-staple. Part 4 Advanced methods. National resources institute, University of Greenwich, UK ISBN 0-85954- 400-, 43-79.
- [26] **Meda A, Lamien CE, Romito M, Millogo J & Nacoulma OG (2005).** Determination of the total phenolic, flavonoid and proline contents in Burkina fasan honey, as well as their radical scavenging activity. Food Chemistry. 91: 571-577.
- [27] **Oulaï S.F., 2014.** Paramètres biochimiques et physic-fonctionnels des farines et des amidons du fruit à pain (*Artocarpus altilis*). Thèse. Université Nangui Abrogoua.160p
- [28] **CAC, (1985).** Codex Standard for Wheat Flour: Codex Stan 152-1985. Codex Alimentarius Commission. Italy: CAC Press.
- [29] **Aguilera J.M., Del Valle J.M. and Karel M. (1995).** Caking phenomenon in amorphous food powders. Trends in Food Sciences Technology 6: pp.149-155.
- [30] **Aryee F.N.A., Oduro I., Ellis W.O. et afuakwa J.J., (2006).** The physicochemical properties of flour samples from the roots of 31 varieties of cassava. Food control 17: pp. 916-922.
- [31] **Yetunde Ezinwanyi Alozie, Mary Arikpo Iyam, Olajumoke Lawal Ukpong Udofia et Ime Franklin Ani, (2009).** Utilisation of Bambara Groundnut flour blends in bread production, *J Food Technol* 7 (4) pp111-114.
- [32] **Alozie Ye, Lyam Ma, Lawal O., Udofia U., Ani If. (2009).** Utilization of Bambara ground flour blends in bread production. In *J Food Technol.* 7(4), 2009, pp.111-114.
- [33] **Tortoe, C., Akonor, PT, Koch, K., Menzel, C. and Adofo, K. (2017).** Physicochemical and functional properties of flour from twelve varieties of Ghanaian sweet potatoes. *Inter Food Res J.* 24(6): pp.2549-2556
- [34] **Gutap V. et Nagar R. (2008).** Physicochemical and acceptability of rabadi (a fermented soya flour product) as affected by cooking and fermentation time. *International Journal of Food and Food Sciences and Technology* 43 : pp.939-943.
- [35] **Anses, 2021.** Rôle des lipides et des proteines. Anses.fr
- [36] **Anuonye J.C., Jigam A.A. et Ndeceko G.M. (2012).** Effects of Extrusion-Cooking of the Nutrient and Anti-Nutrient Composition of Pigeon Pea and Unripe Plantain Blends. *Journal of Applied Pharmaceutical Science* 02 : pp. 158-162.

- [37] Jansen MC, Buenode MHB, Buzina R, Fidenza F, Menotti A, Blackburn H, Nissinen AM, Kok FJ & Kromhout D (1999). Dietary fiber and plant foods in relation to colorectal cancer mortality: the seven countries study. *Int. J. cancer*. 81: 174-179.
- [38] Chene C (2003). *Journal de l'ADRIANOR Agro-jonction n°33*. Septembre-octobre 2003. Les fibres alimentaires pp.1-8
- [39] Al Dobaib N (2009). Effect of diets on growth, digestibility, carcass and meat quality characteristics of four rabbits breeds. *Saudi J. Biol. Sci.* 17: 83-93.
- [40] **Jalili T., Wildamn R.F.C., Medeiros D.M. (2000)**. Nutraceutical roles of dietary fiber. *J.Nutri.Funct.and Med.Foods*. 2, pp.19-34.
- [41] **Ocheme O.B., Oludamilola O.O. et Gladys M.E. (2010)**. Effect of lime soaking and cooking (nixtamalization) on the proximate, functional and some anti-nutritional properties of millet flour. *Assumption University Journal of Technology* 14 : pp.210-216.
- [42] **Martin, A. (2000)**. Apports nutritionnels conseillés pour la population française. Tec et Doc Lavoisier. CNERNA-CNRS. 3e édition, 658p.
- [43] **Folin, J. (2005)**. L'encyclopédie visuelle des aliments. Edition du Chariot d'Or, 6 8 8 p.
- [44] **Fredot, E. (2009)**. Connaissance des aliments. Bases alimentaires et nutritionnelles de la diététique. Coll. BTS Diététique. Tec et Doc.
- [45] **Ezeocha V.C et Ojmelukwe P.C. (2012)**. The impact of cooking on the proximate composition and anti_nutritional factors of water yam (*Dioscorea alata*). *Journal of shored Products and Postharvest Research* 3 : pp.172-176.
- [46] **Butte N. F. (1996)**. Energy requirement of ints European journal of clinical nutrition 50 24.36
- [47] **Mora-escobedo R., Lopez O.P. et Lopez G.F.G. (1991)**. Effect of germination on the rheological and functional properties of amaranth sedes. *Lebensmittel wissenschaft und Technologie* 24 (3): pp 241-244
- [48] **Eke-Ojiofor J., Barber I.I. et Kiin –Kabari D.B. (2011)**. Effect of pre-boiling on the chemical, functional, and pasting properties of rice. *Journal of Agriculture and Biological Sciences* 2 : pp.214-219.
- [49] **Ijarotimi O.S. (2012)**. Influence of germination and fermentation on chemical composition, protein quality and physical properties of wheat flour (*Triticum aestivum*). *Journal of cereals and oil seeds* 3 : pp.35-47.
- [50] **Shittu T.A, Sanni I.O., Awonorin S.O., Maziya-Dixon B. et Dixon A. (2005)**. Use of multivariate techniques in studing flour making characteristics of some cassava Mosaic Disease resistant cassava clone. African Crop Science Conference Proccedings 7 : pp.621-630.
- [51] **Nelson-Quartey F.C., Amagloh F.K, Oduro I. et Ellis WO. (2007)**. Formulation of an infant food based on breadfruit (*Artocarpus altilis*) and brednut (*Artocarpus camansi*). *Acta Horticultura (ISHS)* 757 : pp.212-224.
- [52] **Abulude F.O. (2004)**. Effcets of processing on nutritional composition, phytate and functional properties of rice (*Oriza sativa*) four. *Nigerian Food Journal* 22 : pp. 97-104.

- [53] **Falade, Kolawole O. et Okafor, Chidinma A (2015)**. Physical, functional, and pasting properties of flours from corms of two Cocoyam (*Colocasia esculenta* and *Xanthosoma sagittifolium*) cultivars. *J Food Sci Technol* 2015 Jun; 52(6): pp.3440–3448.
- [54] **Grah Avit Maxwell Beugre1, Beda Marcel Yapo, Sika Hortense Blei, Dago Gnakri. (2014)**. Effect of Fermentation Time on the Physico-Chemical Properties of Maize Flour. *Inter J Res Studies Bios (IJRSB)* 2(8): pp.30-38.
- [55] **Ukpabi U.J. (2010)**. Farmstead Bread making potential of lesser yam (*Dioscorea esculenta*) flour in Nigeria. *Australian Journal of Crop Science* 4 : pp.68-73.
- [56] **Yadahally N. Sreerama ↑, Vadakkoot B. Sashikala, Vishwas M. Pratape, Vasudeva Singh. (2011)**. Nutrients and antinutrients in cowpea and horse gram flours in comparison to chickpea flour: Evaluation of their flour functionality. *Food Chem.* 131 (2012) 462–468.
- [57] **Shindhu Ritu, Bhupendar Singh Khatkar, 2016**. Physicochemical and functional Properties of Starch of flour of Tartary Buckweat (*F. Tataricum*) Grains. *International Journal of Engineering Research &Technology (IJERT)*. Volume 05, Issue 06 (June 2016), <http://dx.doi.org/10.17577/IJERTV5IS060432>
- [58] **Aremu M.O., Olonisakin A., Atolaye B.O. et Ogbu C.E. (2006)**. Some nutritional and functional studies of *Prosopis africana*. *Electronic Journal of Environmental, Agricultural and Food Chemistry* 5: pp.1640-1648.
- [59] **Akubor P.I. (2007)**. Chemical, Functional and cookie baking properties of soybean/maize flour blends. *Journal Food Sciences and Technology* 44 : pp. 619-622.
- [60] **Yasumatsu K., Sawada K., Moritaka S., Mikasi M., Toda J., Wada T., Ishi K. (1972)**. Whipping and emulsifying properties of soybean products. *Agricultural and Biological Chemistry* 36 :PP. 719-727.
- [61] **Ubbor S.C. et Akobundu E.N.T. (2009)**. Quality Characteristics of cookies from Composite flours of Watermelon Seed . Cassava and Wheat. *Pakistan Journal of Nutrition* 8 : pp.1097-1102.
- [62] **Djidohokpin M. E., 2012**. Aptitude des deux variétés de Fonio à la formulation d'aliment de complément. Mémoire. Université d'bomey-Calavi. 85p
- [63] **Tchibindat F., Trèche S., 1995**. Vitafort : Une farine infantile de haute densité énergétique au Congo. In, Trèche S., de Benoit B., Benbouzid D. et Delpeuch F. L'alimentation de complément du jeune enfant. ORSTOM. Paris, France, 177-188.
- [64] **Feillet P. (2000)**. Les grains de blé, composition et utilisation. INRA. Paris. 308p.
- [65] **Mebirouk-Boudechiche L., Cherif M., Boudechiche L. et Sammar F. 2014**. Teneurs en composés primaires et secondaires des feuilles d'arbustes fourragers de la région humide d'Algérie. *Revue de Médecine Vétérinaire*, 165, 11-12, 344-352.
- [66] **Wang S, Moustaid-Moussa N, Chen L, Mo H, Shastri A, Su R. Nouveaux aperçus des polyphénols alimentaires et de l'obésité. J Nutr Biochem. (2014) 25 :1-18.**
- [67] **Fujiki H, Sueoka E, Watanabe T, Suganuma M. Prévention primaire du cancer par le thé vert et prévention tertiaire du cancer par la combinaison de catéchines de thé vert et de composés anticancéreux . J Cancer Préc. (2015) 20 :1-4.**

- [68] **Xiao JB, Hogger P.** Polyphénols alimentaires et diabète de type 2 : perspectives actuelles et perspectives d'avenir . *Curr Med Chem* . (2015) 22 :23-38
- [69] **Martin-Pelaez S, Covas M, Fito M, Kusar A, Pravst I.** Effets sur la santé des polyphénols de l'huile d'olive : avancées récentes et possibilités d'utilisation des allégations de santé . *Mol Nutr Food Res* . (2013) 57 : 760-771.
- [70] **Fraga CG, Galleano M, Verstraeten SV, Oteiza I.** Mécanismes biochimiques de base derrière les bienfaits des polyphénols pour la santé . *Mol Aspects Med.* (2010) 31 :435–445.
- [71] **Alonso R., Rubio L.A., Muzquiz M. et Marzo F., 2001.**The effect of extrusion cooking on mineral bioavailability in pea and kidney bean seed meals. *Animal Feed Sciences and Technology* 94: 1-13.
- [72] **Anigo L.A., Ameh D. A., Ibrahim S. et Danbauchi S.S. (2009).** Nutrient composition of commonly used complementary foods in north western Nigeria. *African Journal of Biotechnology* 8 : 4211-4216
- [73] **Ewald C., Fjelkner-Modig S., Johansson K., Sjöholm L. et Akesson B. (1999).** Effect of processing on major flavonoids in processed onions, greenbeans and peas. *Food Chemistry* 64 : 231-235.
- [74] **PolycarpD., Afoakwa E.O., Budu A.S. et Otoo E. (2012).**Characterization of chemical composition and anti-nutritional factors in seven species within the Ghanaian yam (*Dioscorea*) germplasm. *International Food Research Journal* 19: 985-992
- [75] **Alexandra Retion, 2022.** <https://www.femmeactuelle.fr/sante/alimentation-equilibree/sodium-quel-est-son-role-dans-lorganisme-2097588> Consulté le 18/05/2022
- [76] **Kannel W.B., (1997).** Hazards risks and threat of heart disease from the early stages to symptomatic coronary heart disease and cardiac failure. *Cardiovasc. Drugs and ther.* 11 : 199-212
- [77] **Chow, R., 2009.** Potassium-sodium ratios is crucial for heart health. http://www.naturalnews.com/025820_sodium_potassium_health.html / 05/05/2013
- [78] **Lewu M.N., Adebola P.O. et Afolayan P.J. (2010).** Effect of cooking on the mineral contents and anti-nutritional factors in seven accessions of colocasia esculenta (L.)Shott growing in South Africa. *Journal of Food Composition and Analysis* 23 : 389-393
- [79] **Yagoub A.E.G.A., Mohammed A.M. et Asma A.A.B. (2008).** Effect of Soaking, Sprouting and Cooking on Chemical composition. Bioavailability of minerals and in vitro Protein Digestibility of Roselle (*Hibiscus sabdariffa L.*) Seed. *Pakistan Journal of Nutrition* 7:50-56.
- [80] **Bonithon-Kopp C, Kronborg O, Giacosa A, Rath U, Faivre J. 2000.**Calcium and fiber supplementation in prevention of colorectal adenoma recurrence: a randomized intervention trial. *The Lancet.* 356,1300-1306.
- [81] **Eric Favre, 2022.** Nutrition expert. <https://www.ericfavre.com/lifestyle/fer/> consulté le 18/05/2022
- [82] **Senesse P, Meance S, Cottet V, Faivre J, Boutron-Ruault MC (2004).** High dietary iron and copper and risk of colorectal cancer: a case –control study in Burgundy, France. *Nutr Cancer.* 2004;49:66–71.

Comment [h9]: Please follow proper referencing guideline

- [83] **Ofori* Hayford , Charles Tortoe, Paa Toah Akonor and Jonathan Ampah, 2016.** Trace metal and aflatoxin concentrations in some processed cereal and root and tuber flour. . International Journal of Food Contamination (2016) 3:15 DOI 10.1186/s40550-016-0038-2
- [84] **Organisation Mondiale de la Santé (OMS), 1982.** Evaluation of Certain Foods Additives and Contaminants (Twenty-Six Report of the Joint FAO/WHO Expert Committee on Food Additives). WHO Technical Report series, No. 683, Geneva; 1982.

UNDER PEER REVIEW