

Assessment of Proximal radio-median cubital/radio-cephalic Arterio-venous Fistula

Abstract

The present study aims to conducted the Assessment of Proximal radio-median cubital/radio-cephalic Arterio-venous Fistula. Arterio-venous Fistula is life line for long-term hemodialysis for end stage renal disease patients. The order of preference as per National Kidney Foundation/ Kidney Disease Out Come Quality Initiative (KDOQI) is distal Radio Cephalic fistula is considered as gold standard followed by elbow Brachio Cephalic Fistula, transposed Brachio-Basilic Fistula, forearm arterio-venous graft. This is a cross sectional-prospective interventional study, 05/2017 to 04/2019, JNMC, Wardha, MH, with sample size of 66 cases. Out of 66 cases 25 % patients had diabetes mellitus, 48% cases were suffering from chronic glomerulonephritis, 15 % cases were suffering hypertension, 6 % cases had COPD and another 6 % cases had some cardiovascular disease. About 54 % cases had previous access failure. In our study the mean flow volume for AV fistula in proximal forearm was 485 ± 291 ml/min on postoperative day1, 695 ± 298 on postoperative day 7 and 755 ± 347 ml/min. Overall postoperative complications in 12% cases was reported in our case study though Yilmiz et al reported postoperative complications in 15% cases.

Keywords: Arterio-venous Fistula, chronic glomerulonephritis, Brachio-Basilic Fistula, cardiovascular disease

Introduction

The native arteriovenous fistula (AVF) is recommended as the best vascular access for hemodialysis by the National Kidney Foundation / Kidney Dialysis Outcomes Quality

Initiative (NFK/DOQI) clinical practice guideline. Patients with AVFs have the lowest risks of death, infections, and cardiovascular events and also the lowest costs to the health care system. Arterio-venous Fistula is life line for long-term hemodialysis for end stage renal disease patients. The order of preference as per National Kidney Foundation/ Kidney Disease Out Come Quality Initiative (KDOQI) is distal Radio Cephalic fistula is considered as **gold standard** followed by elbow Brachio Cephalic Fistula, transposed Brachio-Basilic Fistula, forearm arterio-venous graft. There appears increase in failure rates of distal arterio venous fistulas created over wrists in recent years as a result of poor vein quality in distal veins and arteries combined with history of intravenous catheterizations, diabetes mellitus (PVD) and old age.^[1]

Comment [UdW1]: I suggest to include before PVD, "peripheral vascular disease"

Comment [UdW2]:

Because of 20-50% chance for non-maturation (distal Radio Cephalic » Brachio cephalic fistula) & 10-20% chance for Dialysis Access Steal Syndrome of Brachio Cephalic Fistula, Bonforte et.al (2004) demonstrated Arterio-Venous Fistula of proximal Radial artery to nearest vein (cephalic / median anti-brachial / perforating vein).^[1]

Arterio-Venous Fistulas using proximal Radio Cephalic have excellent patency rate, allowing multiple access sites in proximal forearm.^[1] The latest KDOQI guidelines recommend newly created fistulas to be physically examined within 4–6 weeks and additional imaging studies if the fistula is not adequately matured by the sixth week.^[2] Arterio-Venous Fistula maturation becomes apparent as early as 2–4-wks after creation. Arterio-Venous Fistula is considered sono-graphically matured if its Vein internal diameter \geq 4 mm and flow volume (FV) \geq 500 ml/min. (UAB)^[3] Brachial artery flow volume predicts arterio-venous fistula maturation for both forearm & arm.^[4]

Low FV and small Vein Internal Diameter at 2-wks are predictive of arterio-venous fistulas that may not develop optimally at 6-wks. Michelle L. Robbin et al³ in his study with 602 patients in 2016 suggested that Early (USD) identification of causal lesions and problem-

directed interventions shortens the time of fistula maturation & minimizes complications of central venous catheter use. [2]

Assessing outcomes (maturation & Post-operative complications) of proximal Radial-Median Cubital/ radio cephalic(PR-MC/RC) arterio-venous fistula up to 2-wks and collecting data from dialysis unit about fistula functioning status for 1 year. To assess anatomical maturation at post-operative day one, day seven and 14 th post-operative day (D₁, D₇, & D₁₄). To assess postoperative complications ≤ 2-weeks of surgery. To collect information of fistula functioning status for 1 year.

Material and Method

This is a cross sectional-prospective interventional study, 05/2017 to 04/2019, JNMC, Wardha, MH.

With sample size of 66 cases

- Study design -Cross sectional prospective interventional
- Institute- AVBRH Sawangi Meghe, Wardha
- Sample size – 66 cases
- Statistical analysis done by SPSS (Statistical Package for the Social Sciences) software , version 27.0

Inclusion criteria

1. Artery internal diameter ≥ 2 mm
2. Internal diameter ≥ 2.5 mm
3. Non-dominant hand
4. Failed distal radio cubital arterio-venous fistula
5. Absence of stenosis or thrombosis in the draining vein

Exclusion criteria

1. Blood flow less than 50ml/min

Comment [UdW3]: Include ethical items as Sign the informed consent

Comment [UdW4]: Try to unify the type of letter

2. Vascular access less than 15 days on the target arm

3. Hypotensive \leq 110/70 mmhg

4. Advanced heart failure

Comment [UdW5]: Suggest: exclude the patient do not sign the informed consent(ethical of confidentiality)

Details of procedure-

Patient profile including age, sex, BMI & comorbidities like DM, HTN, hyperlipidemia, CD, PVD & COPD. "Fast, 5-min Dialysis Duplex Scan,"^[4] - (Pre-op. & post-op.)

A lazy 'S' shaped skin incision centering in line with the brachial artery pulsation till 4cm below the elbow crease ↓LA. Identification, dissection, & mobilization of A & V. After applying clamps (6 – 8 mm) arteriotomy and venotomy were made. Vein was distended & flushed with heparinized saline by 6 Fr feeding tube in side vein. The M-CV / CV was anastomosed to the pRA in end to side fashion 2 cm below bifurcation of BA by proline 6-0/7-0 suture in a simple continuous manner. Wound was closed in layers with drain, if oozing present.

PSV, EDV, ID, calcification, compressibility, depth & continuity of vessels. FV (ml/min) =

$TAV \times A^2$

$[TAV \approx EDV + 1/3 (PSV-EDV), A = \pi \times r^2]$

[TAV- Timed Average Velocity, EDV- End diastolic Velocity, PSV- Peak Systolic velocity, A- Area, r- radius]

Statistical analysis done by SPSS (Statistical Package for the Social Sciences) software

M-C V / CV- Median Cubital vein/ Cephalic Vein



Fig. 1. Morphological view of the cases

Comment [UdW6]: Suggest: include a brief description of the procedure in the Figure. Figure 1.a Figure 1. B Figure 3 y 4...

Study definitions as per north American vascular access consortium (NAVAC) given below

- D_1 → Immediately after surgery up to 2 days.
- Technical success → presence of a thrill or a bruit immediately and / 24 h postoperatively.
- Primary failure → Thrombosis or inability to cannulate AV fistula within 3 months of creation
- Immediate failure → failure \leq 72 hrs.
- **Early dialysis suitable failure** → Arterio-venous fistula for which, despite the

interventions (radiological/surgical), it is not possible to use successfully for HD by 3rd month following creation.

- Primary patency → time from arterio-venous fistula maturation to the first intervention.

Results:

Total 66 cases were studied between 05/2017 to 04/2019. Out of 66 cases 6 cases failed at 3 months (primary failure). Remaining 60 cases were mature at 1 year.

Table -1 Patient Characteristics

Characteristics of patients	Present study	Gruidah A. ⁵ et al 2019	Yilmiz S. ⁶ et al 2015	Kumar A. ⁷ et al 2007	Ehsan O. ³ et al 2005
M/F	32/34	9/21	31/35	256/64	13/19
Age (Years)	43 (25-72)	52.7 (28-76)	49.6 (22-77)	32 (14-64)	61 (34-80)
DM	25%	80%	40.9%	25%	96.9%
CGN	48%	-	-	-	-
HTN	15%	30%	57.5%	-	-
CVS disease	6%	22.4%	-	-	-
COPD	6%	-	0.9%	-	-
Previous Access failure	54%	20%	62.1%	85%	0%

Comment [UdW7]: Suggest: To check the Vancouver format for Tables and the title (Comparison of patients characteristics with other authors) is better to include chronic obstructive pulmonary disease (COPD), central venous stenosis (CVS), chronic glomerulonephritis (CGN) to locate readers who are from other health professions

In present study we had male to female ratio of 0.94 with mean age as 43 years. Out of 66 cases 25 % patients had diabetes mellitus, 48% cases were suffering from chronic glomerulonephritis, 15 % cases were suffering hypertension, 6 % cases had COPD and another 6 % cases had some cardiovascular disease. About 54 % cases had previous access failure.

Table -2 Operative and post-operative variables

Comment [UdW8]: Idem as Table 2

Operative & Post-operative variables	Present study	Gruidah A. et al ⁵ 2019	Yilmiz S et al ⁶ 2015	Wu C. et al ¹ 2015	Kumar A. et al ⁷ 2007	Ehsan O et al ⁸ 2005
Operative time (min)	85±10	-	-	-	55±7.15	45-60
Left / Right	10 / 1	-	41 / 25	-	7/1	32 / 0
Technical success	100 %	100 %	-	-	100 %	-
Palpable dR pulse	100 %	-	100 %	-	100 %	-
Primary failure	1.5%	0 %	4 %	1.2-33 %	4 %	9%
Preferred access for cannulation	Cephalic 88%	Cephalic 100%	Cephalic 100%	-	Cephalic 70%	-

In present study mean operative time was observed as 85 min ±10 min, out of all the cases operated 10 cases were operated on non-dominant upper limb i.e., left upper limb. Technical success was 100 percent. Distal radial pulse was palpable in 100 % cases. There were 1.5 % patients who had primary failure. The preferred access for cannulation was cephalic in about 88% cases.

Table -3 Vein internal diameter (VID)

Draining Vein ID (mm) (Cephalic)	Present Study	Toregeani et al ¹⁰ 2008	Han A. et al ² 2016	Toregeani et al ¹⁰ 2008	
	Prox. forearm	Dis. forearm	Mixed	Up. Arm	
Pre Operative	3.47 ± 1.41	3.24 ± 1.43	-	3.71 ± 1.37	
Post. Operative	D ₇	5.28 ± 1.39	4.73 ± 0.87	-	5.84 ± 1.23
	D ₁₄	5.54 ± 1.68	4.86 ± 1.05	4.41 ± 1.07	6.02 ± 1.38

In our study mean venous internal diameter preoperatively was 3.47 ± 1.41.

postoperatively on day 7 and day 14 it was about 5.28 ± 1.39 and 5.54 ± 1.68 respectively.

UNDER PEER REVIEW

Table -4 Flow volume for AVF

Post Oper. Period	Flow volume (mL / min) for AVF						
	Proximal Forearm	Distal Forearm		Distal Forearm + Up. Arm			Up. Arm
	Present Study	Zhu et al⁹ 2016	Toregeani et al¹⁰ 2008	Asif A. et al¹¹ 2006	Yerdel M A. et al¹² 1997	Han A. et al² 2016	Toregeani et al¹⁰ 2008
D₁	485 ± 291	364 ± 151	-	-	539 ± 276	-	-
D₇	695 ± 298	-	493.6 ± 257.5	-	848 ± 565	-	976.3 ± 332.9
D₁₄	755 ± 347	634 ± 239	521.3 ± 264.4	692.3 ± 351.2	-	507.4 ± 194.6	1137.7 ± 430.0

Comment [UdW9]: Suggest: title: Comparative findings with other authors of the Flow volume for AVF

In our study the mean flow volume for AV fistula in proximal forearm was 485± 291 ml/min on postoperative day 1, 695 ± 298 on postoperative day 7 and 755± 347 ml/min.

Table -5 Patency of arterio venous fistula

Patency of Arterio-Venous Fistula		Present study	Gruidah ⁵ et al 2019	Yilmiz S ⁶ et al 2015	Wu C et ¹ al 2015	Kumar ⁷ et al 2007	Ehsan O ³ et al 2005
Primary	D ₁₄	98.5%	-	97.0%	-	-	-
	3 mo	94.0%	-	-	-	95%	-
	6 mo	91.0%	-	84.8%	-	-	-
	1 yr	91.0%	63.3%	-	85.7%	92.5%	90.6%

In present study the primary patency at post-operative day 14 was 98.5 %, at 3 months 94 %, at 6 months it was 91 % and at 1 year it was 91%.

Table -6 Post-operative complications

Comment [UdW10]: Unifi

Post operative Complications	Present study	Gruidah A ⁵ et al 2019	Yilmiz S. ⁶ et al 2015	Kumar A. ⁷ et al 2007	Ehsan O. ⁸ et al 2005
Acute thrombosis	0%	13.3%	6 %	-	6.2%
Bleeding	0%	-	-	0%	-
Hematoma	4.5%	-	1.5%	-	-
Arm oedema	7.5%	6.7%	3 %	-	-
Wound infection	0%	-	0%	6.3%	3.1%
Wound dehiscence	0%	-	0%	2.2%	-
Pseudoaneurysm	3%	6.7%	0%	0.3%	-
Thrombophlebitis	0%	-	1.5%	5.6%	-
DIALYSIS ACCESS STEAL SYNDROME	0%	0%	0.5%	0%	0%
Overall	12%	-	15%	-	-

In this study none of the patients developed acute thrombosis, bleeding, wound infection, wound dehiscence, thrombophlebitis or Dialysis Associated Steal Syndrome though there were 4.5% cases who developed hematoma. 7.5 % cases developed arm edema and 3 % cases developed pseudo-aneurysm.

Discussion

In present study we had male to female ratio of 0.94 with mean age as 43 years. Out of 66 cases 25 % patients had diabetes mellitus, 48% cases were suffering from chronic glomerulonephritis, 15 % cases were suffering hypertension, 6 % cases had COPD and another 6 % cases had some cardiovascular disease. About 54 % cases had previous access failure. Gruidah et al did a case study on 30 patients in 2019 in which 9 patients were males and 21 were females, the mean age was 52.7 years. about 80 percent cases had history of DM, 30 percent cases had hypertension, 22.4 percent cases had cardio-vascular disease and 20 percent cases had history of previous access failure. Yilmiz et al studied 66 cases in 2015 out of which there were 31 males and 35 females with mean age of 49.6 years. In his study about 57.5 % cases had history of hypertension, 0.9% cases had history of COPD and 62.1 % had

history of previous access failure. Kumar et al did case series on 320 patients out of which 256 were males and 64 were females with mean age of 32 years, about 25 % patients had history of diabetes mellitus and 85 % cases had history of primary access failure. Ehsan et al studied 32 cases in 2005 of which 13 were males and 19 were males with mean age of 61 years, 96.9% cases had history of diabetes mellitus (see Table 1).

In present study mean operative time was observed as 85 min \pm 10 min, out of all the cases operated 10 cases were operated on non-dominant upper limb i.e., left upper limb. Technical success was 100 percent. Distal radial pulse was palpable in 100 % cases. There were 1.5 % patients who had primary failure. The preferred access for cannulation was cephalic in about 88% cases. Gruidah et al reported 100 % technical success with cephalic route as 100 % preferred access for cannulation. Yilmiz et al performed 66 AV fistulas of which 41 were performed on non-dominant left upper limb and 25 on dominant limb i.e., right upper limb, distal radial pulse (dR) was palpable in 100 % cases with primary failure rate of 4 % and cephalic route as preferred access for cannulation in 100 % cases. The mean operative time for Kumar et al was 55 \pm 7.15 min, with technical success rate and palpable distal radial pulse in 100% cases, primary failure was noted among 4 % cases. Ehsan et al reported minimum operative time average 45 to 60 min, he performed all the surgeries on left upper limb, also he noted primary failure rate in 9 % cases (see Table 2).

In our study mean venous internal diameter preoperatively was 3.47 \pm 1.41. postoperatively on day 7 and day 14 it was about 5.28 \pm 1.39 and 5.54 \pm 1.68 respectively. Toregeani et al reported 3.24 \pm 1.43 mm as mean draining vein internal diameter (VID) in distal forearm pre operatively whereas at post-operative day 7 there was increase in VID mean VID was 4.73 \pm 0.87 mm and on post-operative day 14 it was 4.86 \pm 1.05 mm. Han et al in 2016 reported mean VID on post-operative day 14 as 4.41 \pm 1.07 mm. Toregeani et al reported 3.71 \pm 1.37 mm as mean VID pre operatively in upper arm which was increased to

Comment [UdW11]: Include Reference here.

Comment [UdW12]:

5.84±1.23mm on post-operative day 7 and 6.02±1.38mm on post-operative day 14.the above findings were consistent with our study(see Table 3).

UNDER PEER REVIEW

In our study the mean flow volume for AV fistula in proximal forearm was 485 ± 291 ml/min on postoperative day 1, 695 ± 298 on postoperative day 7 and 755 ± 347 ml/min. Zhu et al in 2016 evaluated mean flow volume of 364 ± 151 ml/min on post op day 1 where as it was found to be increased to 634 ± 239 ml/min on post op day 14. Toregeani et al reported mean flow volume of 493.6 ± 257.5 ml/min on post op day 7 and 521.3 ± 264.4 ml/min on post op day 14. Both above values were calculated on distal forearm, whereas Asif A et al in 2006 reported 692.3 ± 351 ml/min on post op day 14, Yerdel et al in 1997 reported flow volume of 539 ± 276 ml/min on post op day 1 and 848 ± 565 on post op day 7, Han et al reported mean flow volume of 507.4 ± 194.6 on post op day 14. Both above studies were done on distal forearm with upper arm. Toregeani et al reported flow volume of 976.3 ± 332.9 ml/min whereas on post op day 14 it was 1137.7 ± 430 ml/min (see Table 4).

In present study the primary patency at post-operative day 14 was 98.5 %, at 3 months 94 %, at 6 months it was 91 % and at 1 year it was 91%. Gruidah et al reported primary patency of arterio venous fistula was 63.3 % after 1 year. Yilmiz et al reported primary patency of 97.7 % on post op day 14 and 84.8 % at 6 months. Wu C et al reported primary patency of 85.7% at 1 year, whereas Kumar et al reported 95 % patency at 3 months and 92.5% at 1 year. Ehsan et al reported 90.6 % patency at 1 years (see Table 5).

In this study none of the patients developed acute thrombosis, bleeding, wound infection, wound dehiscence, thrombophlebitis or Dialysis Associated Steal Syndrome though there were 4.5% cases who developed hematoma. 7.5 % cases developed arm edema and 3 % cases developed pseudo-aneurysm. Gruidah et. al. reported 13.3% cases had developed acute thrombosis of cephalic vein. Ehsan et.al. reported 6.2% cases developing cephalic vein thrombosis. In our study none of cases developed the same. Yilmiz reported 1.5% cases developed hematoma but, in our study, higher rate of 4.5% was noted. Arm

edema was developed in 6.7 % cases reported by Gruidah et al whereas 3% was noted in study done by Yilmiz et al. Kumar et al and Ehsan et al noted wound infection in 6.3% and 3.1 % cases respectively fortunately none of the cases developed wound infection in our case study. Kumar et al noted wound dehiscence in 2.2% cases, in our study none developed wound dehiscence. Pseudo-aneurysm was seen in 6.7 % cases in study done by Gruidah et al and 0.3 % in study done by Kumar et al, in our case study 3% cases developed pseudo-aneurysm. Thrombophlebitis was seen in 1.5% and 5.6 % in case studies done by Yilmiz et al and Kumar et al respectively. Dialysis access steal syndrome was reported in 0.5 % cases in study done by Yilmiz et al but none were reported in our case series. Overall postoperative complications in 12% cases was reported in our case study though Yilmiz et al reported postoperative complications in 15% cases (see Table 6).

- Intra-operative options for another artery & vein.
- Number of surgical failures was low comparable to elbow fistula.
- Timely increment of FV & Vein Internal Diameter of proximal forearm fistula were better than distal forearm.

Conclusions

- Proximal forearm arterio-venous fistula **should** be recommended as 2nd option after failed /when not fit for distal forearm arterio-venous fistula.

Comment [UdW13]: It is suggested to expand the conclusions according to the findings and comparison with other studies, surgical techniques, post-surgical care, among others.

References

1. Cong Cong Wu, Hua Jiang, Jun Cheng et al. The outcome of the proximal radial artery arteriovenous fistula. *J Vasc Surg.* **2015**;61:802-8.
2. Ahram Han,* Seung-Kee Min,* Mi-Sook Kim,† Kwon Wook Joo,‡ Jungsun Kim,* Jongwon Ha,* Joongyub Lee,† and Sang-il Min* A Prospective, Randomized Trial of Routine Duplex Ultrasound Surveillance on Arteriovenous Fistula Maturation. *Clin J Am*

Soc Nephrol. **2016**;11:1817–1824.

3. Michelle L. Robbin, Tom Greene, Alfred K. Cheung et al. Arteriovenous Fistula Development in the First 6 Weeks after Creation. *Radiology*. **2016**;279(2):620–629.
4. Sae Hee Ko, Dennis F. Bandyk, Kelley D. Hodgkiss-Harlow et al. Estimation of brachial artery volume flow by duplex ultrasound imaging predicts dialysis access maturation. *J Vasc Surg*. **2015**;61:1521-8.
5. Hesham S. Abu-Gruidah, Mahmoud S. Abd El-Haleim, Mohamed A. Haridy. Evaluation of the proximal radio cephalic arteriovenous fistula for hemodialysis. *Menoufia Medical Journal*. **2019**;32:534–538.
6. Sehyan Yilmaz, Eray Aksoy, Serdar Gunaydin. Radio-medyan cubital arteriovenous creation for hemodialysis access. *Turkish Journal of Thoracic and Cardiovascular Surgery*. **2015**.
7. Anant Kumar, Mrignak S. Jha, Manish Singhla. Radio-median cubital / radiocephalic arteriovenous fistula at elbow to prevent vascular steal syndrome associated with brachiocephalic fistula: Review of 320 cases. *Indian J Urol*. **2007**; 23(3): 261–264.
8. O. Ehsan, D. Bhattacharya, A. Darwish¹ and H. Al-khaffaf. 'Extension Technique': A Modified Technique for Brachio-Cephalic Fistula to Prevent Dialysis Access-Associated Steal Syndrome. *Eur J Vasc Endovasc Surg*. **2005**;Vol 29:.
9. Yu-li Zhu, MD, Hong Ding, MD, Pei-li Fan, MD, Qi-lan Gu, MD, Jie Teng, MD, Wen-ping Wang, MD. Is Brachial Artery Blood Flow Measured by Sonography During Early Postoperative Periods Predictive of Arteriovenous Fistula Failure in Hemodialysis Patients? *J Ultrasound Med*. **2016**;35:1985–1992.

10. Jeferson F, Toregeani, Claudio J, Kimura, Antonio S. et al. Evaluation of hemodialysis arteriovenous fistula maturation by color-flow Doppler ultrasound. *J. vasc.bras.* **2008**;vol.7(3):
11. Asif A, Roy-Chaudhury P, Beathard GA: Early arteriovenous fistula failure: A logical proposal for when and how to intervene. *Clin J Am Soc Nephrol.* **2006**;1:332–339.
12. Yerdel MA, Kesenci M, Yazicioglu KM, Doseyen Z, Turkcapar AG, Anadol E: Effect of haemodynamic variables on surgically created arteriovenous fistula flow. *Nephrol Dial Transplant.* **1997**;12:1684–1688.

UNDER PEER REVIEW