

MICROBIAL ANALYSIS OF CONTACT LENSES

ABSTRACT

Introduction: Contact lenses are small, thin lenses which are worn directly on the surface of the eyes. They can be worn aesthetically or to correct vision. Contact lens related eye infections can lead to serious complications such as blindness, and are associated with several risk factors such as sleeping with lenses, exposure of lenses to water, not adhering to replacement schedules, and reusing disinfecting solutions, among other sete. The severity of the infection may vary with the degree of pathogenicity of the microorganism. Hygiene and handling of contact lenses play a very important role. The main aim of this study is to assess the microbial analysis of contact lenses.

Materials and methods: A total of 15 lenses were collected in 2 ml sterile saline solution and agitated for five minute manually. The lens was then removed using a sterile toothpick. The sterile container was stored at 4°C till it was processed. 50 microliter were pipetted and inoculated on nutrient agar, blood agar and sabouraud dextrose agar. The microorganism (fungus or bacteria) were identified by standard protocol.

Comment [LQ1]: Not clear for me

Results and discussion: Bacillus, Staphylococcus aureus and CONS [Coagulase Negative Staphylococci] were found in blood agar. No fungal growth was found among the samples. More importance could be given to contact lenses handling and hygiene to avoid eye related bacterial and fungal infections.

Conclusion: In the present study, the total CFU was found to be confluent in all the participants who wore contact lenses for 4 months compared to those who have worn contact lenses from 15 and 28 days. Only bacterial growth was seen in the culture plate. There was no fungal growth seen from the samples collected.

Comment [LQ2]: Stands for?

Keywords: contact lenses; microbial keratitis; hygiene; eye infections; innovative technology.

Running title: To assess microbial analysis of contact lenses

INTRODUCTION

The contact lenses market is ever-growing. Contact lenses are small lenses which are worn directly on the surface of the eyes. They can be worn aesthetically or to correct vision. Few of the advantages of contact lenses are that they have wider field of view, better with asymmetrical prescriptions ~~and better~~ for very high prescriptions, cosmetically superior, more practical for sports, avoid weather problems, etc. And few of the disadvantages may be the time required for fitting and adaptation, handling of the patients, wearing time may be limited, lenses can be lost or broken, hygiene maintenance, lens disinfection, problems with foreign bodies, etc (1). There are two main types of contact lenses; hard and soft contact lenses and there are two types of disposable lenses; daily wear disposable lenses and most extended wear disposable lenses (2). Each company manufactures lenses that are made from different materials such as hypergel, hydrogel and silicone hydrogel. Hydrogel lenses with silicone hydrogel were found to be used the most up to date, though many other materials were present in it (3). Contact lens related eye infections can lead to serious complications such as blindness, are associated with several risk factors such as sleeping in lenses, exposure of lenses to water, not adhering to replacement schedules, and reusing disinfecting solutions, etc. ~~Specifically, m~~Microbial keratitis is an ocular infectious disease affecting the cornea and ~~pathogenetically~~ pathogenically resulting from microorganisms which may potentially cause ocular disability (4).

The severity of the infection may vary with the degree of pathogenicity of the microorganism. Microbial keratitis associated with wearing contact lenses still remains a serious concern for patients and the contact lens industry (5). Acanthamoeba is a ubiquitous pathogen which is found worldwide. Acanthamoeba keratitis could potentially cause blinding corneal infection ~~and that can~~ aggressively infect both the eyes. Contact lens related problems depend on lens material, lens hygiene, wearing procedure, degree of compliance of the lenses wearer with contact lenses and related procedures, type of lens caring solution, etc. Basic hygiene if followed by contact lenses wearers can reduce the rate of eye infections (6).

In a study done by Jennifer. R. Cope et al, it was seen that ~~the~~ most frequent behaviour of contact lens wearers was sleeping ~~with~~ contact lenses. And also few of the contact lens wearers do not follow the recommended ~~contact lens~~ replacement schedules leading to eye discomfort and ~~other more~~ complications (7). Strong association was found between the bacterial contamination on

Comment [LQ3]: Is this the right way to cite in this journal?

lens surfaces and the bacterial contamination found on eyelids and conjunctiva. This can be due to the disruption of the normal microbiota due to the use of contact lenses leading to unbalance between staphylococcus and corynebacterium. The underlying mechanism of how contact lenses wear affects the normal microbiota of the eye is still unknown (8). Hygiene and handling of contact lenses play a very important role. More knowledge and awareness on various eye infections resulting from poor hygiene and handling must be spread. The main aim of this study is to assess the microbial analysis of contact lenses.

Comment [LQ4]: Improve writing

MATERIALS AND METHODS

A total of 15 lenses (collected from participants belonging to age below 20 years) were collected in 2 ml sterile saline solution (varying in power and material type) and agitated for five minutes manually. The lens was then removed using a sterile toothpick. The sterile container was stored at 4°C till it was processed. 50 microliter was pipetted and inoculated on nutrient agar, blood agar and sabouraud dextrose agar. Nutrient and blood agar was divided into two halves and 2 samples were inoculated and kept at 37° Celsius for 24 hours. Also sabouraud dextrose agar was divided into two halves and 2 samples were inoculated and kept at 37° Celsius for 24 hours and then at room temperature for 24 hours. The microorganism (fungus or bacteria) were identified by standard protocol.

RESULTS

[Explain and mention the figures and tables in the text](#)



FIGURE 01: Sample inoculated on blood agar.




FIGURE 02: Sample inoculated on nutrient agar.




FIGURE 03: Sample inoculated on sabouraud dextrose agar.

SAMPLES	NO. OF DAYS CONTACT LENS WORN	BACTERIA COUNT
1	15 days	313
2	15 days	412
3	15 days	561
4	15 days	411
5	15 days	456
		Mean= 430.6

TABLE 01: Table depicts total CFU obtained from the contact lenses in relation to 15 days of usage and the total mean value.

Comment [LQ5]: It stands for?

SAMPLES	NO. OF DAYS CONTACT LENS WORN	BACTERIA COUNT
6	28 days	703
7	28 days	590
8	28 days	442
9	28 days	367
10	28 days	421
		Mean= 504.6

TABLE 02: Table depicts total CFU obtained from the contact lenses in relation to 28 days of usage and the total mean value.

SAMPLES	NO. OF DAYS CONTACT LENS WORN	BACTERIA COUNT
11	4 months	Confluent (>1500)
12	4 months	449
13	4 months	487
14	4 months	441
15	4 months	524
		Mean= 680.2

TABLE 03: Table depicts total CFU obtained from the contact lenses in relation to 4 months of usage and the total mean value.

DISCUSSION

In the present study, it was noticed that those who wear lenses daily were less prone to suffer bacterial infection of the eye than those who change their lenses once in a month or year. Bacillus, Staphylococcus aureus and CONS [Coagulase Negative Staphylococci] was found in blood agar (Figure 01) (Figure 02). No fungal growth was found among the samples (Figure 03). The mean value of the total CFU for the samples used for 15 days were found to be 430.6 (Table 01), 504.6 for 28 days (Table 02) and 680.2 for 4 months (Table 03). Wearing contact lenses is a ~~known~~ well-known risk factor for the development of microbial keratitis and other inflammatory eye conditions. A previous study done by Sarah. A. Collier et al, an online survey was prepared and distributed among 4,548 adults and 1618 adolescents. ~~Adolescents and young adults were~~ Both groups were subjected to lower compliance with contact lens hygiene which can result in a greater risk of corneal inflammation and serious eye infection (9–20),(21–25). Young adults were found more likely to replace their contact lenses at regular schedules compared to adolescents. This might be due to daily lifestyle behaviour. The authors stated that awareness and prevention efforts and knowledge on contact lens related infection can be spread through improving communication strategies for a healthier contact lens behaviour (26). In another research article by Maya M Rao et al, it was found that the most frequent behaviour of contact lenses was that the participants slept with contact lenses ~~worn~~. Though the precise mechanism of eye infection is unknown, prolonged lens replacement schedules, non-compliance to recommended lenses can lead to eye infections.

Daily disposable contact lens wearers were less prone to eye infections related to contact lenses (27) (28). Showering while wearing contact lenses, swimming while wearing contact lenses

Comment [LQ6]: Most of the studies should have been previously mentioned in the introduction

Comment [LQ7]: Start with the objectives of the study

Comment [LQ8]: Replace?

Comment [LQ9]: Who does it nowadays?

without wearing goggles, should be avoided as associated with sight threatening infections were found. Contact lenses hygiene was noticed as a compulsory and a very important factor in public health (29). According to Wu Yvonne T et al, the major factors for reducing contact lens contaminants are washing hands with soap and water before use, air drying lens case and matching disinfection solution with lens cases (30). And in another article written by Vinicius Targa Villas Boas et al, it was found that no fungal growth was found in the lenses but bacterial growth was found in 39 cases out of 70 lens cases. Many variables were present in the study not allowing the exact type of bacteria to be identified. But more than 85% of the bacteria were gram- negative bacilli (31).

Bailey CS had stated in their article that polymethyl- methacrylate (PMMA) lenses were 9 times less likely to be associated with extended wear soft lenses. And also extended wear soft lenses were 5 times more prone to be linked with microbial keratitis than daily wear soft lenses (32). Despite the increased use of daily disposable contact lenses, bacterial ulcer incidence is on the rise. Sleeping with contact lenses was one of the main risk factors in contact lens related infections. Acanthamoeba and fungal infections related to contact lenses are also on the rise which could lead to severe vision and blinding threats (33). More importance could be given to contact lenses handling and hygiene to avoid eye related bacterial and fungal infections. Few of the limitations seen in the present study was that the sample size was small and samples from all age groups could have been collected. These limitations can be considered for future studies.

CONCLUSION

In the present study, the mean CFU was found to be confluent in the participants who wore contact lenses for 4 months compared to those who wore contact lenses from 15 and 28 days. Only bacterial growth was seen in the culture plate. There was no fungal growth seen from the samples collected. In future studies, the limitations seen in the present study could be considered for better results and understanding.

REFERENCES

1. Szczotka-Flynn LB, Pearlman E, Ghannoum M. Microbial contamination of contact lenses, lens care solutions, and their accessories: a literature review. *Eye Contact Lens*. 2010 Mar;36(2):116–29.
2. Foulks GN. Prolonging contact lens wear and making contact lens wear safer. *Am J Ophthalmol*. 2006 Feb;141(2):369–73.
3. Musgrave CSA, Fang F. Contact Lens Materials: A Materials Science Perspective. *Materials* [Internet]. 2019 Jan 14;12(2). Available from: <http://dx.doi.org/10.3390/ma12020261>
4. Ezisi C, Ogbonnaya C, Okoye O, Ezeanosike E, Ginger-Eke H, Arinze O. Microbial Keratitis—A Review of Epidemiology, Pathogenesis, Ocular Manifestations, and Management [Internet]. Vol. 26, *Nigerian Journal of Ophthalmology*. 2018. p. 13. Available from: http://dx.doi.org/10.4103/njo.njo_2_18
5. Evans DJ, Fleiszig SMJ. Microbial keratitis: could contact lens material affect disease pathogenesis? *Eye Contact Lens*. 2013 Jan;39(1):73–8.
6. Cope JR, Collier SA, Schein OD, Brown AC, Verani JR, Gallen R, et al. Acanthamoeba Keratitis among Rigid Gas Permeable Contact Lens Wearers in the United States, 2005 through 2011. *Ophthalmology*. 2016 Jul;123(7):1435–41.
7. Cope JR, Konne NM, Jacobs DS, Dhaliwal DK, Rhee MK, Yin J, et al. Corneal Infections Associated with Sleeping in Contact Lenses - Six Cases, United States, 2016-2018. *MMWR Morb Mortal Wkly Rep*. 2018 Aug 17;67(32):877–81.
8. Szczotka-Flynn LB, Bajaksouzian S, Jacobs MR, Rimm A. Risk factors for contact lens bacterial contamination during continuous wear. *Optom Vis Sci*. 2009 Nov;86(11):1216–26.
9. Priyadharsini JV, Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species [Internet]. Vol. 94, *Archives of Oral Biology*. 2018. p. 93–8. Available from: <http://dx.doi.org/10.1016/j.archoralbio.2018.07.001>
10. Vijayashree Priyadharsini J. In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens. *J Periodontol*. 2019 Dec;90(12):1441–8.
11. Paramasivam A, Vijayashree Priyadharsini J, Raghunandhakumar S. N6-adenosine methylation (m6A): a promising new molecular target in hypertension and cardiovascular diseases. *Hypertens Res*. 2020 Feb;43(2):153–4.
12. Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. An insight into the emergence of *Acinetobacter baumannii* as an oro-dental pathogen and its drug resistance gene profile - An in silico approach. *Heliyon*. 2018 Dec;4(12):e01051.

13. Paramasivam A, Vijayashree Priyadharsini J. Novel insights into m6A modification in circular RNA and implications for immunity. *Cell Mol Immunol*. 2020 Jun;17(6):668–9.
14. Paramasivam A, Priyadharsini JV, Raghunandhakumar S. Implications of m6A modification in autoimmune disorders. *Cell Mol Immunol*. 2020 May;17(5):550–1.
15. Girija ASS, Shankar EM, Larsson M. Could SARS-CoV-2-Induced Hyperinflammation Magnify the Severity of Coronavirus Disease (CoViD-19) Leading to Acute Respiratory Distress Syndrome? *Front Immunol*. 2020 May 27;11:1206.
16. Jayaseelan VP, Arumugam P. Exosomal microRNAs as a promising theragnostic tool for essential hypertension. *Hypertens Res*. 2020 Jan;43(1):74–5.
17. Ushanthika T, Smiline Girija AS, Paramasivam A, Priyadharsini JV. An in silico approach towards identification of virulence factors in red complex pathogens targeted by reserpine. *Nat Prod Res*. 2021 Jun;35(11):1893–8.
18. Ramalingam AK, Selvi SGA, Jayaseelan VP. Targeting prolyl tripeptidyl peptidase from *Porphyromonas gingivalis* with the bioactive compounds from *Rosmarinus officinalis*. *Asian Biomed*. 2019 Oct 1;13(5):197–203.
19. Kumar SP, Girija ASS, Priyadharsini JV. Targeting NM23-H1-mediated inhibition of tumour metastasis in viral hepatitis with bioactive compounds from *Ganoderma lucidum*: A computational study. *pharmaceutical-sciences* [Internet]. 2020;82(2). Available from: <https://www.ijpsonline.com/articles/targeting-nm23h1-mediated-inhibition-of-tumour-metastasis-in-viral-hepatitis-with-bioactive-compounds-from-ganoderma-lucidum-a-comp-3883.html>
20. Mathivadani V, Smiline AS, Priyadharsini JV. Targeting Epstein-Barr virus nuclear antigen 1 (EBNA-1) with *Murraya koengii* bio-compounds: An in-silico approach. *Acta Virol*. 2020;64(1):93–9.
21. Samuel SR, Kuduruthullah S, Khair AMB, Shayeb MA, Elkaseh A, Varma SR. Dental pain, parental SARS-CoV-2 fear and distress on quality of life of 2 to 6 year-old children during COVID-19. *Int J Paediatr Dent*. 2021 May;31(3):436–41.
22. Samuel SR. Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life? *Int J Paediatr Dent*. 2021 Mar;31(2):285–6.
23. Barma MD, Muthupandiyam I, Samuel SR, Amaechi BT. Inhibition of *Streptococcus mutans*, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish. *Arch Oral Biol*. 2021 Jun;126:105132.
24. Teja KV, Ramesh S. Is a filled lateral canal - A sign of superiority? *J Dent Sci*. 2020 Dec;15(4):562–3.
25. Reddy P, Krithikadatta J, Srinivasan V, Raghu S, Velumurugan N. Dental Caries Profile and Associated Risk Factors Among Adolescent School Children in an Urban South-Indian

City. *Oral Health Prev Dent*. 2020 Apr 1;18(1):379–86.

26. Cope JR, Collier SA, Nethercut H, Jones JM, Yates K, Yoder JS. Risk Behaviors for Contact Lens–Related Eye Infections Among Adults and Adolescents — United States, 2016 [Internet]. Vol. 66, *MMWR. Morbidity and Mortality Weekly Report*. 2017. p. 841–5. Available from: <http://dx.doi.org/10.15585/mmwr.mm6632a2>
27. Mihailidis M. Legal Issues and Contact Lenses [Internet]. *Contact Lenses*. 2019. p. 529–34. Available from: <http://dx.doi.org/10.1016/b978-0-7020-7168-3.00031-3>
28. Cope JR, Collier SA, Rao MM, Chalmers R, Lynn Mitchell G, Richdale K, et al. Contact Lens Wearer Demographics and Risk Behaviors for Contact Lens–Related Eye Infections – United States, 2014 [Internet]. Vol. 64, *MMWR. Morbidity and Mortality Weekly Report*. 2015. p. 865–70. Available from: <http://dx.doi.org/10.15585/mmwr.mm6432a2>
29. Arshad M, Carnt N, Tan J, Ekkeshis I, Stapleton F. Water Exposure and the Risk of Contact Lens–Related Disease [Internet]. Vol. 38, *Cornea*. 2019. p. 791–7. Available from: <http://dx.doi.org/10.1097/ico.0000000000001898>
30. Wu YT, Willcox MDP, Stapleton F. The effect of contact lens hygiene behavior on lens case contamination. *Optom Vis Sci*. 2015 Feb;92(2):167–74.
31. Bôas VTV, Almeida Júnior GC de, Almeida MTG de, Gonçalves MS, Coelho LF. Microbiological analysis of contact lens cases: impact of the hospital environment. *Arq Bras Oftalmol*. 2018 Sep;81(5):371–5.
32. Bailey CS. A review of relative risks associated with four types of contact lenses. *Cornea*. 1990;9 Suppl 1:S59–61; discussion S62–3.
33. Cheung N, Nagra P, Hammersmith K. Emerging trends in contact lens-related infections. *Curr Opin Ophthalmol*. 2016 Jul;27(4):327–32.