

# **DISTRIBUTION OF FOUR BIOFILM ASSOCIATED GENE AMONG *A.BAUMANNII* BY IN SILICO-PCR**

## **ABSTRACT:**

### **Background:**

*A.baumannii* is an opportunistic pathogen known for its efficient biofilm formation that is attributed for its virulence. *Acinetobacter baumannii* is an inhabitant of oral biofilms as well. Many gene operons are involved in the biofilm formation and the present study is designed to assess the frequency of four vital biofilm forming genes among 19 different strains of *A.baumannii*.

### **Aim:**

The aim of the present study was to detect the distribution of four biofilm associated genes among *A.baumannii*.

### **Materials and methods:**

Four biofilm forming genes viz., *bfms*, *ptk*, *pgaB*, *fimH* of *A.baumannii* were selected. Forward and reverse primers of those four genes as reported from earlier studies were used for in-silico PCR amplification. 19 strains of *A.baumannii* set as default on the server were chosen and the amplicon bands were observed

### **Results:**

The present investigation documents the distribution of four vital biofilm associated gene among 19 different strains of *A.Baumannii* among which *bfms* was distributed at a higher frequency followed by *pgaB* and *ptk*

### **Conclusion:**

The finding of the study suggests the presence of *pgaB*, *bfms*, *ptk* among the 19 different strains of *A.baumannii*. However further experimental validation must be done to frequently monitor the presence of the genes among the clinical strains of *A.baumannii*.

**Keywords:** Biofilm; *A.baumannii*; novel *pgaB* ; *bfms*; *ptk*; innovative in-silico: environmental strains

**Running title:** Detection of biofilm associated genes among *A.baumannii*

## **INTRODUCTION:**

*Acinetobacter baumannii* is a typically short, almost round, rod-shaped gram-negative bacterium. An opportunistic pathogen in humans, affecting people with compromised immune systems, and is becoming increasingly important as a nosocomial pathogen. While other species of the genus *Acinetobacter* are often found in soil samples, it's almost adapted to survive in harsh hospital niches. Although occasionally it's been found in environmental soil and water samples. *A. baumannii*, a Gram-negative, the most successful nosocomial pathogen within a short time period with 2–10% of death rate recorded among patients with chronic tract infections, bacteremia, pneumonia,, and critically ill-patients in ICU. The World Health Organization [WHO] has provided an alert about the carbapenem-resistant *A. baumannii* which secures its place under the critical category (1, 2), (3).

The biofilms produced by the bacteria also are potential reservoirs of pathogens related to pneumonia and chronic obstructive pulmonary disease. Biofilms are collectively one or more sorts of microorganisms that will grow on many various surfaces (4). Microorganisms that form biofilms include bacteria, fungi, and protists. Biofilms are found growing on both biotic and abiotic surfaces and on implanted medical devices like catheters and pacemakers. Biofilm formation is a process whereby microorganisms irreversibly attach to and grow on a surface and produce extracellular polymers that facilitate attachment and matrix formation, resulting in an alteration in the phenotype of the organisms with respect to growth rate and gene transcription (5). The BfmRS two-component system plays a vital role in pathogenesis and antimicrobial resistance of *A. baumannii* via the regulation of bacterial envelope structures. This study investigated the role of the sensor kinase, BfmS, in localization of outer membrane protein (OmpA) within the outer membrane, and the production of outer membrane vesicles (OMVs) (6) (7).

Several studies have indicated that the few other vital genes like *fimA* gene encoding for the massive secondary unit while the *fimF* and *fimG* genes encode the tiny subunits, and therefore, the *fimH* gene encodes the highest of the cilia that are sensitive to the manus and therefore the

fimC gene encode the attached protein that helps fimD encodes the outer membrane proteins and fimI encodes The fimH gene is a crucial virulence agent for bacteria that encodes the sort 1 fimbriae, that helps bacteria to bind to the surface of host cells that cause injury (8). In addition, *A.baumannii* contains a pgaABCD locus that encodes proteins that blend cell-related poly- $\beta$ -(1-6)- N-acetylglucosamine (PNAG) (9). Different expansive consecutive investigations report that pgaABCD operon during *A.baumannii* and various other gram-positive and negative microscopic organisms also encode PNAG

The polymerase chain reaction (PCR) is prime to biology and is the most vital practical molecular technique for the lab. The principle of this system has been further used and applied in many other simple or complex macromolecule amplification technologies (10). In parallel to laboratory experiments for macromolecule amplification technologies in silico or virtual (bioinformatics) approaches are developed, among which in silico PCR analysis is promising. Our team has extensive knowledge and research experience that has translated into high quality publications (11, 12) (13) (14). The present investigation is thus undertaken to evaluate the frequency of four biofilm associated genes among the 19 different strains of *A.baumannii* by computational approach.

## **MATERIALS AND METHODS:**

**Study setting:** This study was carried out as an observational study using a computational approach. Institutional approval for carrying out the study was obtained (SRB NUMBER:IHEC/SDC/UG-1895/21/163). The present study was done with the help of in-silico amplification tools (computerised in silico.ehu.es). Use of pre-identified forward and reverse primers of the four genes (bfms, ptk, pgaB, fimH) we were able to determine their length of the base pairs and the number of bands observed among 19 strains of *A.baumannii* using the in silico amplifier web page. And with the help of these primers obtained we can compare and determine the virulence factors of each genome.

### ***Evolutionary analysis by Maximum Likelihood method:***

The developmental history was derived by utilizing the Maximum Likelihood strategy and Tamura-Nei model (16). The tree with the most elevated log probability (- 1116.82) has appeared. Introductory tree(s) for the heuristic inquiry were obtained consequently by applying Neighbor-Join and BioNJ calculations to a network of pairwise distances assessed utilizing the

Tamura-Nei model, and afterward choosing the geography with prevalent log probability esteem. investigation included 14 nucleotide groupings (16). Codon positions included were 1st+2nd+3rd+Noncoding. There were a sum of 365 situations in the last dataset investigations were directed in MEGA X (17,18).

## RESULTS:

The investigation on the prevalence of the drug resistant genes from 19 different strains of *A.baumannii* using an in-silico amplification server was promising. The results showed the starting position of the amplification in the chromosome or plasmid and the length of each amplicon. Amplicons obtained in each chromosome or plasmid have been tabulated [Table 1] with target genes, primers used, sequenced of primer (5' to 3'), annealing temperature, estimated size of base pair and the frequency of the target genes among the study strains. In the present study, BfmS showed an amplicon size of 1428 bp. PtK Showed an amplicon size of 597 bp [Table 1]. PgaB showed an amplicon size of 490bp. fimH showed no bands (Figure 1-3). Nearly 52.63% of distribution for bfms, 21.05% of distribution for pgaB and about 10.52% of distribution for ptk was observed in the present study among the selected 19 different strains of *A.baumannii*. We further assessed the evolutionary pattern of the distributed genes among the strains as given in figure 4.

**Table 1 showing the target genes, PCR conditions of the biofilm genes selected for the study**

Target Genes	Primers sequences (5–3)	Annealing Temperature (°c)	DNA amplicon Size (bp)	Bands
<i>bfmS</i>	TTGCTCGAACTTCCAATTTATTATAC TTATGCAGGTGCTTTTTTATTGGTC	60	1428	10
<i>ptk</i>	GGCTGAGCATCCTGCAATGCGT ACTTCTGGAGAAGGGCCTGCAA	57	597	2
<i>pgaB</i>	AAGAAAATGCCTGTGCCGACCA GCGAGACCTGCAAAGGGCTGAT	57	490	4
<i>fimH</i>	TGCAGAACGGATAAGCCGTGG GCAGTCACCTGCCCTCCGGTA	60	870	No bands

**Table 2 showing the distribution of the bfmS among the selected strains of *A.baumannii***

<b>Strain name</b>	<b>Sequence id</b>	<b>Start position</b>	<b>End position</b>	<b>Base pair length</b>
Acinetobacter baumannii 1656-2 chromosome	NC_017162	823749	825176	1428
Acinetobacter baumannii ACICU	NC_010611	793047	794474	1428
Acinetobacter baumannii ATCC 17978	NC_009085	889190	890617	1428
Acinetobacter baumannii BJAB07104	NC_021726	834877	836304	1428
Acinetobacter baumannii BJAB0868	NC_021729	830598	832025	1428
Acinetobacter baumannii D1279779	NC_020547	789219	790646	1428
Acinetobacter baumannii MDR-TJ	NC_017847	3143572	3144999	1428
Acinetobacter baumannii MDR-ZJ06	NC_017171	821690	823117	1428
Acinetobacter baumannii TCDC-AB0715	NC_017387	826988	828415	1428
Acinetobacter baumannii TYTH-1	NC_018706	1043958	1045385	1428






Figure 1 showing BfmS with an amplicon size of 1428 bp among the selected strains of *A.baumannii*

UNDER PEER




**Figure 2 showing PtK with an amplicon size of 597bp among the selected strains of *A.baumannii***

UNDER PEER REVIEW



**Figure 3 showing PgaB with an amplicon size of 490bp among the selected strains of *A.baumannii***

UNDER PEER REVIEW



**Figure 4 showing the phylogenetic evolutionary pattern of BfmS among the positive strains of *A.baumannii***

#### **DISCUSSION:**

*A.baumannii* and its associated virulence of biofilm formation attributes for the initiation and progression of the disease in the hospital environment. Many genetic determinants of biofilms

are documented in the pathogenesis of *A.baumannii*. The present study is thus undertaken to determine the frequency of four vital biofilm forming genes among 19 different strains of *A.baumannii* using an in-silico amplification server. Polymerase chain reaction is used for primer designing and is used for selective amplification of the target genetic determinants. Recent improvements in technologies have made it easy to regulate a high specific theoretical possibility of a successful PCR by modifying a high specific and sensitive primer instead of laboratory assay which is expensive (19) (20). There are several free web servers available which we can use to identify the possible outcomes of a target gene using their preexisting forward and reverse primers. Using the specific primers as reported from earlier studies for each different gene, we evaluated frequency of the distribution of the genes among the selected strains (21).

From the results obtained, the present investigation documents the distribution Bfms at a higher frequency followed by *pgaB* and *ptk* these findings correlates with many earlier studies. However *fimH* which was not reported in the study was highly responsible for virulence in earlier reports. (22). In the previous study on *fimH* gene, states that *fimH* can directly stimulate host cell signaling cascades that lead to bacterial internalization and exhibit greater frequencies when compared to our present study (23). Type 1 fimbriae has also been described as a major factor in the formation of biofilm on the abiotic surface. More specifically, *fimH* was shown to be necessary to adhere to ephemeral surfaces in stable growth conditions (24) (25). Thus in the present study, we have chosen *fimH* for detection among the *A.baumannii* strains (26).

Earlier studies on bfms in *A.baumannii* showed decreased biofilm formation, adherence to eukaryotic cells, and greater sensitivity to serum killing as compared to parent strains. Our study had documented the highest frequency of bfms which correlates with many earlier reports. (27–29). In previous studies it has been revealed that the high frequency of biofilm forming genes among the XDR *A. baumannii* from ICU patients. In the same line, presence of biofilm forming genes like *pgAb* and *ptk* has shown greater frequencies which is more significant in our study. In-silico based computational approaches on the amplification of the target genes seem to be highly promising in the detection at a preliminary level. The results were, however, correlated with earlier studies with few contrasting reports as well. The limitation of the study was that the strains were selected from the default set-up in the tool and not with the clinical strains. Future prospects are set to experimentally validate the distribution

of the genes among the clinical strains of *A.baumannii* and using the conventional PCR for amplification (30)(31)(32)

## CONCLUSION:

The present study was undertaken to detect the frequency of the four important biofilm genes among 19 virulent strains of *A.baumannii*. The finding of the study suggests the presence of 3 genes viz., *pgaB*, *bfms*, *ptk* with no distribution of *fimH* gene. However further experimental validation must be done to frequently monitor the presence of the genes among the clinical strains of *A.baumannii* and to curtail the infections in healthcare settings.

## REFERENCES:

1. Mishra DR, Shah DS, Shah N, Prasad JN, Gupta PP, Agrawal KK. Study of microbiological and antibiotic sensitivity pattern of ventilator associated pneumonia (VAP) in ICU of a tertiary care hospital in Nepal. *J Family Med Prim Care*. 2020 Dec;9(12):6171–6.
2. Priyadharsini JV, Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. An insight into the emergence of *Acinetobacter baumannii* as an oro-dental pathogen and its drug resistance gene profile – An in silico approach. Vol. 4, *Heliyon*. 2018. p. e01051.
3. Priyadharsini JV. In silico validation of the non- antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens. Vol. 90, *Journal of Periodontology*. 2019. p. 1441–8.
4. Website [Internet]. Available from: Priyadharsini JV, Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species. Vol. 94, *Archives of Oral Biology*. 2018. p. 93–8.
5. Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species. *Arch Oral Biol*. 2018 Oct;94:93–8.
6. Padmaja S, Smiline Girija AS, Vijayashree Priyadharsini J. Frequency of Adhesive Virulence Factor *fimH* among the Clinical Isolates of *Acinetobacter baumannii* in India [Internet]. *Journal of Pharmaceutical Research International*. 2020. p. 12–7.
7. Paramasivam A, Vijayashree Priyadharsini J, Raghunandhakumar S. N6-adenosine methylation (m6A): a promising new molecular target in hypertension and cardiovascular diseases. *Hypertens Res*. 2020 Feb;43(2):153–4.
8. Paramasivam A, Vijayashree Priyadharsini J. Novel insights into m6A modification in

- circular RNA and implications for immunity. *Cell Mol Immunol.* 2020 Jun;17(6):668–9.
9. Samuel SR, Kuduruthullah S, Khair AMB, Shayeb MA, Elkaseh A, Varma SR. Dental pain, parental SARS-CoV-2 fear and distress on quality of life of 2 to 6 year-old children during COVID-19. *Int J Paediatr Dent.* 2021 May;31(3):436–41.
  10. Forman A, Pfoh R, Eddenden A, Howell PL, Nitz M. Synthesis of defined mono-de-N-acetylated  $\beta$ -(1→6)-N-acetyl-d-glucosamine oligosaccharides to characterize PgaB hydrolase activity. *Org Biomol Chem.* 2019 Nov 6;17(43):9456–66.
  11. Little DJ, Bamford NC, Howell PL. Structure of the PgaB (BpsB) glycoside hydrolase domain from *Bordetella bronchiseptica*. 2018;9:1-9.
  12. Paramasivam A, Priyadharsini JV, Raghunandhakumar S. Implications of m6A modification in autoimmune disorders. *Cell Mol Immunol.* 2020 May;17(5):550–1.
  13. Kumar A, Chordia N. In Silico PCR Primer Designing and Validation. *Methods in Molecular Biology.* 2015. p. 143–51.
  14. Girija ASS, Shankar EM, Larsson M. Could SARS-CoV-2-Induced Hyperinflammation Magnify the Severity of Coronavirus Disease (CoViD-19) Leading to Acute Respiratory Distress Syndrome? *Front Immunol.* 2020 May 27;11:1206.
  15. Kalendar R, Lee D, Schulman AH. FastPCR Software for PCR, In Silico PCR, and Oligonucleotide Assembly and Analysis. *DNA Cloning and Assembly Methods.* 2014. p. 271–302.
  16. Tamura K. On the estimation of the rate of nucleotide substitution for the control region of human mitochondrial DNA [Internet]. Vol. 259, *Gene.* 2000. p. 189–97.
  17. Jayaseelan VP, Arumugam P. Exosomal microRNAs as a promising theragnostic tool for essential hypertension. *Hypertens Res.* 2020 Jan;43(1):74–5.
  18. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol.* 2018 Jun 1;35(6):1547–9.
  19. Yu B, Zhang C. In Silico PCR Analysis [Internet]. *Methods in Molecular Biology.* 2011. p. 91–107.
  20. Ushanthika T, Smiline Girija AS, Paramasivam A, Priyadharsini JV. An in silico approach towards identification of virulence factors in red complex pathogens targeted by reserpine. *Nat Prod Res.* 2021 Jun;35(11):1893–8.
  21. Barma MD, Muthupandiyan I, Samuel SR, Amaechi BT. Inhibition of *Streptococcus mutans*, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish. *Arch Oral Biol.* 2021 Jun;126:105132.
  22. Martinez JJ. Type 1 pilus-mediated bacterial invasion of bladder epithelial cells [Internet]. Vol. 19, *The EMBO Journal.* 2000. p. 2803–12.
  23. Ramalingam AK, Selvi SGA, Jayaseelan VP. Targeting prolyl tripeptidyl peptidase from *Porphyromonas gingivalis* with the bioactive compounds from *Rosmarinus officinalis*.

Asian Biomed . 2019 Oct 1;13(5):197–203.

24. Morris NS, Stickler DJ, McLean RJC. The development of bacterial biofilms on indwelling urethral catheters [Internet]. Vol. 17, World Journal of Urology. 1999. p. 345–50.
25. Clark RB. Imipenem resistance among *Acinetobacter baumannii*: association with reduced expression of a 33-36 kDa outer membrane protein. *J Antimicrob Chemother.* 1996 Aug;38(2):245–51.
26. Zeighami H, Valadkhani F, Shapouri R, Samadi E, Haghi F. Virulence characteristics of multidrug resistant biofilm forming *Acinetobacter baumannii* isolated from intensive care unit patients. *BMC Infect Dis.* 2019 Jul 17;19(1):1–9.
27. Samuel SR. Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life? *Int J Paediatr Dent.* 2021 Mar;31(2):285–6.
28. Teja KV, Ramesh S. Is a filled lateral canal - A sign of superiority? *J Dent Sci.* 2020 Dec;15(4):562–3.
29. Reddy P, Krithikadatta J, Srinivasan V, Raghu S, Velumurugan N. Dental Caries Profile and Associated Risk Factors Among Adolescent School Children in an Urban South-Indian City. *Oral Health Prev Dent.* 2020 Apr 1;18(1):379–86.
30. Ghasemi E, Ghalavand Z, Goudarzi H, Yeganeh F, Hashemi A, Dabiri H, et al. Phenotypic and Genotypic Investigation of Biofilm Formation in Clinical and Environmental Isolates of *Acinetobacter baumannii* [Internet]. Vol. 13, Archives of Clinical Infectious Diseases. 2018.