

## **Original Research Article**

### **RESPIRATORY RESPONSES TO SHORT TERM SUSTAINED ISOMETRIC MUSCLE CONTRACTION AMONG UNDERGRADUATES**

#### **ABSTRACT**

#### **AIM**

This study sets out to investigate whether a short-term isometric abdominal exercise can benefit adults with improvement in ventilatory functions.

#### **MATERIALS AND METHODS**

An experimental design with a convenient sampling technique was used. The study enrolled 20 adults from the student population of I-BDS students with no history of smoking and respiratory illness. The participants did a daily 20-minute static abdominal exercise over a period of three days. The lung function test was assessed using spirometry and the values of forced vital capacity (FVC), forced expiratory volume in the 1st sec (FEV1), FEV1/FVC ratio, peak expiratory flow rate (PEFR) and forced expiratory flow (FEF25-75%) were obtained. Statistical tests?????

#### **RESULTS**

The study demonstrated a significant increase in the mean values of forced vital capacity and forced expiratory volume in one second. The FEV1/FVC ratio, PEFR, FEF 25-75% did not reveal significant changes.

#### **CONCLUSION**

The study concluded an innovative finding that ventilatory functions improved after short term isometric training and higher sample size need to be evaluated to demonstrate its efficacy in adults at high risk of chronic obstructive lung disease??? (This statement cannot be written as a conclusion it would rather be a future direction)

**Key words: sustained (How is this a keyword?), isometric exercise, lung function tests, innovation???(Why is this one of the keywords?)**

## **INTRODUCTION**

An isometric muscle contraction is a form of exercise involving the static contraction of a muscle without any visible movement in the angle of the joint.(1)Isometric exercises help maintain strength(2). **They can also build strength, but not effectively. The cardiovascular responses to sustained, isometric muscle contractions have been well documented, and include FVC - Forced Vital Capacity, FEV1 - Forced Expiratory Volume, PEFR - Peak Expiratory Flow Rate and FEF - Forced Expiratory Flow. (These are not cardiovascular parameters)** For a particular muscle, or group of muscles, the respiratory changes are related to the force of contraction of the muscle after short term sustained isometric muscle contraction.(3)(4) The mass of muscle involved has less effect on the blood pressure and heart rate responses. The respiratory responses to isometric muscle contraction have been less well investigated. The studies have mainly been on short term isometric abdominal muscle contraction. (4)It represents the responses to short term isometric muscle contractions of a small mass of muscle. Some subjects hyperventilate inappropriately which leads to marked falls in alveolar partial pressure of  $\text{CO}_2$  ( $\text{PCO}_2$ ). The present studies aim to compare directly the respiratory responses to contraction of a small mass of muscle and of a larger mass, the short term isometric abdominal muscle contraction. (5)We will test the hypothesis that the increase of ventilation during sustained isometric contractions is independent of the mass of the muscle involved, although oxygen consumption may rise with increasing active muscle mass. The reduction of alveolar  $\text{PCO}_2$ , resulting from hyperventilation, should be less with a larger muscle mass as carbon dioxide production will be increased.(6) Preliminary accounts of this work have already been communicated to the Physiological Society.(7)This study sets out to investigate whether a short-term isometric abdominal exercise can benefit adults with improvement in ventilatory functions.(8)**Why is reference given for this**

statement?

## MATERIALS AND METHODS

A pre-experimental design with a convenience sample was employed. All the study participants signed consent forms before being enrolled. The study group consisted of 20 adult males, ranging from 18 to 25 years old with similar anthropometric measurements like age, gender, height and body mass index (BMI) (Without statistical analysis we cannot say these variables are similar)

Inclusion criteria : Healthy adults from the student population of I-BDS students.

Exclusion criteria – Subjects with any respiratory illness, COPD or spinal deformities were excluded from the study.

The study was introduced and explained to the participants, who were then encouraged to handle the spirometer and arm??. They were taught short term static isometric exercise for 20 mins over a period of three days. Both the pre and post exercise routine spirometry tests were evaluated using HELIOS SPIROMETER and the values of Forced vital capacity (FVC), forced expiratory volume at the end of the first second (FEV1), FEV1/FVC ratio and peak expiratory flow rate (PEFR) and FEF 25-75 % using a spirometer both before and after 3 days of isometric exercise for 20 mins.

Statistical Analysis was done using SPSS Software and the Paired dependant t test was done to evaluate the changes in lung functions pre and post evaluation??.

For anthropometric variables, what statistical test is used???

## RESULTS

The present study showed that there was a significant increase in FVC, FEV1 after short term isometric exercise and it was found to be statistically significant ( $p < 0.05$ ). (Fig 1 & 2). There was a non-significant decrease in FEV1/FVC (equal values are given in the

table), PEFr and FEF25-75 (Fig 3,4 & 5). (Table -1)


Table for anthropometric variables ???

Table-1 represents the lung functions in pre and post sustained isometric exercise training

Lung function parameters	Pre isometric training	Post isometric training
FVC	2.18 $\pm$ 0.444	3.12 $\pm$ 0.571
FEV1	2.28 $\pm$ 0.385	3.32 $\pm$ 0.284
FEV1/FVC	100.01 $\pm$ 0.000	100.01 $\pm$ 0.000
PEFR	4.72 $\pm$ 2.695	2.64 $\pm$ 1.342
FEF 25-75%	3.39 $\pm$ 2.007	3.39 $\pm$ 1.393


Unit of variables???

Values are expressed as mean  $\pm$  S.D




**FIGURE 1.** Represents the changes in FVC among the adults.

FVC shows an increase after short term sustained isometric muscle contraction and the value was found to be statistically significant as in Paired dependant t test  $p= 0.05$  ( $p<0.05$ )??




**FIGURE 2.** Represents the changes in forced expiratory volume in first second (FEV1) among the adults.

FEV1 shows an increase after short term sustained isometric muscle contraction and the value was found to be statistically significant as in Paired dependant t test  $p= 0.00$  ( $p<0.05$ )




**FIGURE 3.** Represents the changes in FEV1/FVC ratio among the adults.

The FEV1/FVC ratio is equal before and after short term sustained isometric muscle contraction and the value was found was not statistically significant as in Paired dependant t test  $p= 0.07$  ( $p>0.05$ )



**FIGURE 4.** Represents the changes in FEF 25-75% among the adults.

FEF 25-75% shows a decrease after short term sustained isometric muscle contraction and the value was not statistically significant as in Paired dependant t test  $p= 0.78$  ( $p>0.05$ )



**FIGURE 5.** Changes in PEFR among the adults.

PEFR shows a decrease after short term sustained isometric muscle contraction and the value was not statistically significant as in Paired dependant t test  $p= 0.66$  ( $p>0.05$ )

## DISCUSSION

A number of research studies have been carried out to investigate the influence of physical activity and sport exercises on pulmonary function in patients who suffer from respiratory problems. The present study sets out to discover whether pulmonary function tests have different results in response to static abdominal exercise for 3 days. The

present study revealed that there were significant improvements in FVC, FEV1. Previous reports suggested that short term isometric exercises can have a facilitatory effect on the lungs particularly FEV1 values. It may be a three days exercise program, but this regimen enhanced the vital capacity and expiratory volumes. Also that the subjects' physical exertion during exercise could have helped to create reduced resistance to respiration and it has given the respiratory muscles an endurance training. This is likely to be an explanation for the significant difference in FVC values noted in our study.

Also that this effect is mediated to increase the ventilation caused by increased alveolar  $PCO_2$ . However, cardiovascular and respiratory changes following short term isometric exercise are partly stimulated by drive from higher centres to the brain stem and role of chemoreceptors. (6)

Previous reports also suggested that aerobic exercise improves respiratory functions and increases the body's ability to use oxygen. It tones up all the muscles, improving the circulation in the process. It lowers blood pressure and reduces workload to the heart. It strengthens the respiratory muscles and, even reduces airflow resistance and facilitates the inflow and outflow of air in the lungs effectively (3)

### **LIMITATIONS OF THE STUDY**

The study population was confined only to a small group. If more sample size is added the results would have been statistically significant.

### **CONCLUSION**

The study concluded that short term isometric exercise significantly improves the lung functions and can reduce the predisposition of developing obstructive lung disease.

### **REFERENCES**

1. Institute NC, National Cancer Institute. Isometric Exercise [Internet]. Definitions. 2020. Available from: <http://dx.doi.org/10.32388/7bxexn>
2. R. R. Review Article on Effectiveness of Isometric Exercise on Osteoarthritis [Internet]. Vol. 24, International Journal of Psychosocial Rehabilitation. 2020. p. 843–6. Available from: <http://dx.doi.org/10.37200/ijpr/v24i5/pr201755>
3. Barnes M. CHRONIC EXERTIONAL COMPARTMENT SYNDROME: MUSCLE CHANGES WITH ISOMETRIC EXERCISE [Internet]. Vol. 35, Medicine & Science in Sports & Exercise. 2003. p. 1794. Available from: <http://dx.doi.org/10.1249/01.mss.0000091501.73963.1d>
4. Imms FJ, Mehta D. Respiratory responses to sustained isometric muscle contractions in man: the effect of muscle mass. *J Physiol*. 1989 Dec;419:1–14.
5. Edwards RHT, Harris RC, Hultman E, Kaijser L, Koh D, Nordesjö L-O. Effect of temperature on muscle energy metabolism and endurance during successive isometric contractions, sustained to fatigue, of the quadriceps muscle in man [Internet]. Vol. 220, The Journal of Physiology. 1972. p. 335–52. Available from: <http://dx.doi.org/10.1113/jphysiol.1972.sp009710>
6. Williams CA. Effect of muscle mass on the pressor response in man during isometric contractions [Internet]. Vol. 435, The Journal of Physiology. 1991. p. 573–84. Available from: <http://dx.doi.org/10.1113/jphysiol.1991.sp018526>
7. Sato H. Electromyographic Study of Muscle Fatigue in Sustained Isometric Contractions [Internet]. Vol. 85, The Journal of Anthropological Society of Nippon. 1977. p. 83–94. Available from: <http://dx.doi.org/10.1537/ase1911.85.83>
8. Davies CTM, Young K. Muscle weakness following sustained and rhythmic isometric contractions in man [Internet]. Vol. 54, European Journal of Applied Physiology and Occupational Physiology. 1985. p. 301–5. Available from: <http://dx.doi.org/10.1007/bf00426149>
9. Barabadi, H. *et al.* (2021) ‘Green synthesis, characterization, antibacterial and biofilm inhibitory activity of silver nanoparticles compared to commercial silver nanoparticles’, *Inorganic Chemistry Communications*, p. 108647. doi: 10.1016/j.inoche.2021.108647.
10. Bharath, B. *et al.* (2021) ‘Evaluation of the anticancer potential of Hexadecanoic acid from brown algae *Turbinaria ornata* on HT–29 colon cancer cells’, *Journal of Molecular Structure*, p. 130229. doi: 10.1016/j.molstruc.2021.130229.
11. Clarizia, G. and Bernardo, P. (2019) *Diverse Applications of Organic-Inorganic Nanocomposites: Emerging Research and Opportunities: Emerging Research and Opportunities*. IGI Global.
12. Egbuna, C., Mishra, A. P. and Goyal, M. R. (2020) *Preparation of Phytopharmaceuticals for the Management of Disorders: The Development of Nutraceuticals and Traditional Medicine*. Academic Press.
13. Ezhilarasan, D. (2020) ‘Critical role of estrogen in the progression of chronic liver diseases’, *Hepatobiliary & pancreatic diseases international: HBPD INT*, 19(5), pp. 429–434.
14. Gowhari Shabgah, A. *et al.* (2021) ‘Shedding more light on the role of Midkine in hepatocellular carcinoma: New perspectives on diagnosis and therapy’, *IUBMB life*, 73(4), pp. 659–669.

15. J, P. C. *et al.* (2018) 'Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study', *Clinical Implant Dentistry and Related Research*, pp. 531–534. doi: 10.1111/cid.12609.
16. Kamath, S. M. *et al.* (2020) 'In vitro augmentation of chondrogenesis by Epigallocatechin gallate in primary Human chondrocytes - Sustained release model for cartilage regeneration', *Journal of Drug Delivery Science and Technology*, p. 101992. doi: 10.1016/j.jddst.2020.101992.
17. Mudigonda, S. K. *et al.* (2020) 'Non-suturing microvascular anastomosis in maxillofacial reconstruction- a comparative study', *Journal of Cranio-Maxillofacial Surgery*, 48(6), pp. 599–606.
18. Nambi, G. *et al.* (2018) 'Spinal manipulation plus laser therapy versus laser therapy alone in the treatment of chronic non-specific low back pain: a randomized controlled study', *European journal of physical and rehabilitation medicine*, 54(6), pp. 880–889.
19. Prakash, A. K. S. and Devaraj, E. (2019) 'Cytotoxic potentials of *S. cumini* methanolic seed kernel extract in human hepatoma HepG2 cells', *Environmental Toxicology*, pp. 1313–1319. doi: 10.1002/tox.22832.
20. Rajakumari, R. *et al.* (2020) 'Grape seed extract-soluplus dispersion and its antioxidant activity', *Drug development and industrial pharmacy*, 46(8), pp. 1219–1229.
21. R, H. *et al.* (2020) 'CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene', *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*, pp. 306–312. doi: 10.1016/j.oooo.2020.06.021.
22. Santhakumar, P. *et al.* (2021) 'Ethanollic Extract of Capparis decidua Fruit Ameliorates Methotrexate-Induced Hepatototoxicity by Activating Nrf2/HO-1 and PPAR $\gamma$  Mediated Pathways', *Indian Journal of Pharmaceutical Education*, 55(1s), pp. s265–s274.
23. Saraswathi, I. *et al.* (2020) 'Impact of COVID-19 outbreak on the mental health status of undergraduate medical students in a COVID-19 treating medical college: a prospective longitudinal study', *PeerJ*, 8, p. e10164.
24. Sridharan, G. *et al.* (2019) 'Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(4), pp. 299–306.
25. Tahmasebi, S. *et al.* (2021) 'The effects of oxygen-ozone therapy on regulatory T-cell responses in multiple sclerosis patients', *Cell biology international*, 45(7), pp. 1498–1509.
26. Vivekanandhan, K. *et al.* (2021) 'Emerging Therapeutic Approaches to Combat COVID-19: Present Status and Future Perspectives', *Frontiers in Molecular Biosciences*. doi: 10.3389/fmolb.2021.604447.
27. Wadhwa, R. *et al.* (2021) 'Anti-inflammatory and anticancer activities of Naringenin-loaded liquid crystalline nanoparticles in vitro', *Journal of food biochemistry*, 45(1), p. e13572.
28. Wahab, P. U. A. *et al.* (2018) 'Scalpel Versus Diathermy in Wound Healing After Mucosal Incisions: A Split-Mouth Study', *Journal of Oral and Maxillofacial Surgery*, pp. 1160–1164. doi: 10.1016/j.joms.2017.12.020.

UNDER PEER REVIEW