

Study Protocol

PROTOCOL ON: TO EVALUATE THE EFFECTIVENESS OF NURSING PROTOCOL ON KNOWLEDGE AND PRACTICE REGARDING HEMODIALYSIS CARE AMONG PROFESSIONAL STAFF

ABSTRACT

Background: The prevalence rate of chronic kidney disease (CKD) ranges from 1 < percent to 13 percent and recently international society of nephrology reported In India, the prevalence rate of chronic kidney disease was 17 percent in 2015. The risk factors and etiological factors are different in overall India. **Objective :** 1.To assess the existing level of knowledge regarding hemodialysis care among professional staffs. 2. To assess the existing Practice regarding hemodialysis care among professional staffs. 3.To assess the effectiveness of nursing protocol on knowledge regarding hemodialysis care among professional staffs.4.To assess the effectiveness of nursing protocol on practice regarding hemodialysis care among professional staffs. 5.To find out the correlation between knowledge and practice regarding hemodialysis care among professional staffs. 6.To associate the level of knowledge and practice regarding hemodialysis with selected demographic variable. **Methodology:** It is Quasi experimental one group pre-test post-test research design will be used in this research study conducted on the professional staff in the selected hospital of Wardha. Non-probability purposive sampling technique will be used to select samples for analysis. This research study included 60 professional staff of selected hospital of Wardha district. . Professional staff must select according to requirements for inclusion and exclusion. For this study, Professional staffs who are giving care to the hemodialysis patients. Professional staffs who are willing to participate. **Expected Results:** Outcome includes To evaluate the nursing care for hemodialysis patients. Through, improve and evaluate the knowledge and performance of hemodialysis care among professional staff before and after the education and training interventions. To assess the existing practice of hemodialysis care and the generation of new evidence. **Where is correlations**

Comment [D1]: Objective 3 and 4 summed in one point ... it should be To assess the effectiveness of nursing protocol on knowledge and practice regarding hemodialysis care among professional staffs

Comment [D2]: No need for this objective

Comment [D3]: Pre-posttests

Comment [D4]: Is their more than 60 nurses already working in hemodialysis unite????

Ethics approval was obtained from (DMIMS(DU)/IEC/DEC-2019/8689). The conclusion will be drawn from the results.

Conclusion: Conclusion will be drawn from the statistical analysis.

Keyword: nursing protocol, knowledge, practice, hemodialysis care ,professional staff

INTRODUCTION

"Nurses Are The Heart Of Hospital"

Comment [D5]: Introduction usually without titles

Dialysis nurses should have good knowledge and have skillful practices in hemodialysis (HD) treatment. In the dialysis unit have an important role in monitoring, promoting the treatment, and educating of patients. In the developing and developed countries, with changes in lifestyle and advances in life expectancy, chronic diseases such as end-stage renal diseases, cardiac diseases, and diabetes mellitus are increasing steadily.¹⁻³

A major health problem in the world is chronic renal disease. End-stage renal disease (ESRD) is the number of patients that are irreversible and progressive kidney failure disease in developed and developing countries to increase Chronic Renal Failure (CRF). Chronic renal failure (CRF) results from total loss of kidney function. Total loss of kidney function results in chronic renal failure (CRF).^{3,4}

The prevalence rate of chronic kidney disease (CKD) ranges from 1 < percent to 13 percent and recently international society of nephrology reported In India, the prevalence rate of chronic kidney disease was 17 percent in 2015. The risk factors and etiological factors are different in overall India.⁵⁻⁷

End-stage renal disease or chronic kidney disease are common health problems in India. It includes diabetes mellitus, elderly, obese people. Many people around the world suffer from End-stage renal disease (ESRD) and require long-term dialysis.⁸⁻¹⁰

Chronic renal failure is medically managed with fluid adjustments, correction of electrolyte imbalances, proper low protein diet, and drug .dialysis and renal replacement are the ultimate renal replacement therapy.^{11,12}

Standards of nursing care that the nurse shows and learns through nursing activities include assessment, diagnosis, recognition of outcomes, planning, implementation and evaluation. The nursing method considers clinical decision-making concepts and fundamentals, clinical guidance and includes all essential nurses' practices in delivering treatment to all patients.^{13,14}

Professional nursing practice standards are authoritative descriptions of the duties that are required to be performed competently by all registered nurses, regardless of their position, level of care or efficiency, beliefs and preferences, population or specialization, by which the quality of nursing practice can be measured.¹⁵⁻¹⁷

AIM OF THE STUDY

To evaluate the nursing care for hemodialysis patients. Through, improve and evaluate the knowledge and performance of hemodialysis care among professional staff before and after the education and training interventions. To assess the existing practice of hemodialysis care and the generation of new evidence.

OBJECTIVES

1. To assess the existing level of knowledge regarding hemodialysis care among professional staff.
2. To assess the existing Practice regarding hemodialysis care among professional staff.
3. To assess the effectiveness of nursing protocol on knowledge regarding hemodialysis care among professional staff.
4. To assess the effectiveness of nursing protocol on practice regarding hemodialysis care among professional staff.
5. To find out the correlation between knowledge and practice regarding hemodialysis care among professional staff.
6. To associate the level of knowledge and practice regarding hemodialysis with selected demographic variable.

Comment [D6]: Objective 3 and 4 summed in one point ... it should be To assess the effectiveness of nursing protocol on knowledge and practice regarding hemodialysis care among professional staffs

CRITERIA OF THE STUDY

Inclusion Criteria

1. Professional staff who are giving care to the hemodialysis patients.
2. Professional staff who are willing to participate.

Exclusion Criteria

1. Those who are not directly involved in hemodialysis patient care.
2. Those who are participants in similar type of study.

Comment [D7]: Remove

RESEARCH HYPOTHESIS

H₀

Comment [D8]: Where is null hypothesis

H₁. There is a significant difference between pre-test and post-test knowledge and practices on hemodialysis care among professional staff.

RESEARCH METHODOLOGY

It is an academic based study. In this study will be used interventional research approach. Quasi experimental one group pre-test post-test research design will be used. The study will be conducted in selected hospital of Wardha district. Sampling technique will be used non probability purposive sampling technique. In this research study the accessible population will be the professional staff in selected hospital of Wardha district.




Figure 2: Schematic presentation of one group pre test and post-test design

SAMPLE SIZE –

To calculated to according working staffs.

In consideration of 10% dropout,120 participants will be assigned

The sample consists of the unite which comprise the population. Sixty (60) professional staffs were selected using calculated formula to suit the study. Formula for calculating sample size.¹⁶

$$N = \frac{X^2 \cdot N \cdot P(1-P)}{C^2 (N-1) + X^2 \cdot P(1-P)}$$

Where;

X^2 is the chisquare value for 1. df at 95% confidence interval = 1.96

N= Total professional staff = 70

P= 50% Proportion = 0.50

C^2 = Desired error of level = 0.05

$$N = \frac{3.84 \times 70 \times 0.50 \times 0.50}{(0.05^2 \times 69) + (3.84 \times 0.50 \times 0.50)}$$
$$= 59.64$$

N= 60 professional staff needed in the study

INTERVENTIONS –

On the first day of the study, first the pre-test knowledge regarding hemodialysis care among professional staff will be assessed through a structured knowledge questionnaire. Observe the per practice regarding hemodialysis care among professional staff will be assessed through the observational checklist.

After the education and training interventions will be given on the same day to the professional staff.

On the 7th day post-test the effectiveness of nursing protocol on knowledge and practice regarding hemodialysis care among professional staff.

OUTCOME MEASURES

Primary outcomes assessment of existing knowledge and practice regarding hemodialysis care among professional staff.

Secondary outcomes assessment of post- test knowledge and practice regarding hemodialysis care among professional staff.

The effectiveness of nursing protocol on knowledge and practice regarding hemodialysis care among professional staff.

Correlation between knowledge and practice regarding hemodialysis care among professional staff.

Association of knowledge and practice regarding hemodialysis with selected demographic variable.

DATA MANAGEMENT AND MONITORING - The demographic data (like age, education, place of work, total year experience.....), in this pre-test to assess the exiting level of awareness. Than given the intervention of nursing protocol on knowledge and practice regarding hemodialysis care after 7 the day to taken the post test.

Comment [D9]: Add..
Attendance of workshops or training programs
Gender

STRUCTURED QUESTIONNAIRE

It consists of three sections

Section A. Consist of demographic such as age, qualification, place of work, total year experience

Section B. Consists of 20 questions on knowledge questionnaire on hemodialysis care

Section C. Consists of 24 steps on the Observation checklist

Scoring

- The correct answer was given 1 score.
- The wrong answer was given a 0 score
- Knowledge was categorized from poor knowledge to excellent knowledge based on the scores
- Practice was categorized from poor practice to excellent knowledge based on the scores

Table No-1 : Scoring method to assess the knowledge regarding the care of hemodialysis

Sr.no.	Level of knowledge score	Score range
1.	Poor	0-5
2.	Average	6-10
3.	Good	11-15
4.	Excellent	16-20

Table No-2: Scoring method to assess the practice regarding the care of hemodialysis

Sr.no.	Level of knowledge score	Score range
1.	Poor	0-6
2.	Average	7-12
3.	Good	13-18
4.	Excellent	19-24

STATISTICAL ANALYSIS - Statistical analyses will be performed using SPSS software version

23.

ETHICS AND DISSEMINATION- This study is approved by the Institutional Ethics Committee of DMIMS (DMIMS(DU)/IEC/DEC-2019/8689). All participants will be asked to read and sign the informed consent. The study results will be disseminated to study participants and published in peer-reviewed publications.

EXPECTED OUTCOMES/RESULTS:

This study is planned to improve effectiveness of nursing protocol on knowledge and practice regarding hemodialysis care among professional staff.

DISCUSSION :

A quasi-experimental study was conducted in Egypt. The convenience samples of forty one nurses were all women and had prior experience caring for patients on hemodialysis. 68.3 percent of nurses were unaware of hemodialysis nursing care quality, according to the report. More than half of nurses (61.0 percent - 56.1 percent) had not attended a workshop in the previous five years and had not attended educational lectures. In comparison to the pre-test time, an equal proportion of nurses achieved very good or excellent overall total knowledge and efficiency in the post-test and follow-up times, as well as an improvement in total mean scores for findings related to overall infection control results.¹⁷

A quasi-experimental study was conducted in Egypt. 90.6 percent of the nurses had adequate levels of expertise. In addition to the overall pre score, the nurses' practice scores improved significantly in the areas of general precautions, hand washing procedure, procedure of wearing gloves, wearing masks, procedure of catheter insertion, skin preparation procedure, taking care of patients' instruments, dealing with sharp items, dealing with blood, body secretions, and fluids, and ensuring a clean environment and healthy injection practices. Several aspects of the nurses' work, such as laboratory-style covering coats, complete isolation, and catheter treatment, saw major changes. The practice score level was satisfactory in 68.7 percent of the nurses, while it was insufficient in 31.3 percent. Older nurses and those with a bachelor's degree in nursing had slightly higher awareness scores and levels of progress after completing the education program.¹⁸

CONCLUSION: Conclusion will be drawn from the statistical analysis.

BIBLIOGRAPHY

1. Manandhar DN, Chhetri PK, Poudel P, Baidya SK, Agrawaal KK. Knowledge and practice of hemodialysis amongst dialysis nurses. *JNMA J Nepal Med Assoc.* 2017 Jul 1;56(207):346-51
2. Nasri H, Rafieian-Kopaei M. Diabetes mellitus and renal failure: Prevention and management. *Journal of research in medical sciences: the official journal of Isfahan University of Medical Sciences.* 2015 Nov;20(11):1112.
3. Schieppati A, Remuzzi G. Chronic renal diseases as a public health problem: Epidemiology, social, and economic implications. *Kidney International.* 2005 Sep 1;68:S7–10
4. Anand S, Shivashankar R, Ali MK, Kondal D, Binukumar B, Montez-Rath ME, Ajay VS, Pradeepa R, Deepa M, Gupta R, Mohan V. Prevalence of chronic kidney disease in two major Indian cities and projections for associated cardiovascular disease. *Kidney international.* 2015 Jul 1;88(1):178-85.
5. Abdelsalam KE, muslem em. Influence of end stage renal disease in alteration of some trace elements in sudanese patients. *Int J Cur Res Rev| Vol.* 2016 May 22;8(10):16.
6. lewis , bucher, heitkemper , harding , kwong , roberts. *medical surgical nursing.* 3rd ed. Vol. volume II. elsevier; 2018. 995–1009 p.
7. John T. Daugirdas , peter G.B lake. *handbook of dialysis.* In: 5th ed. Wolters Kluwer Health; p. 66–215
8. Potter PA, Perry AG, Hall AE, Stockert PA. *Fundamentals of nursing.* Elsevier mosby; 2009.
9. King CR, NP R, Sally Gerard DN, Rapp CG, editors. *Essential knowledge for CNL and APRN nurse leaders.* Springer Publishing Company; 2018 Dec 28.

10. Saleh M, Ali J, Afifi W. Nurses Compliance to Standards of Nursing Care for Hemodialysis Patients: Educational and Training Intervention. IOSR Journal of Nursing and Health Science (IOSR-JNHS) e-ISSN. 2018;2320-1959.
11. Siegel JD, Rhinehart E, Jackson M, Chiarello L. 2007 guideline for isolation precautions: preventing transmission of infectious agents in health care settings. American journal of infection control. 2007 Dec 1;35(10):S65-164.
12. Thomas A, Silver SA, Rathe A, Robinson P, Wald R, Bell CM, Harel Z. Feasibility of a hemodialysis safety checklist for nurses and patients: a quality improvement study. Clinical kidney journal. 2016 Jun 1;9(3):335-42.
13. Kim S, Lee HZ, Hwang E, Song J, Kwon HJ, Choe K. Lived experience of Korean nurses caring for patients on maintenance haemodialysis. Journal of clinical nursing. 2016 May;25(9-10):1455-63.
14. Baidoo IK, Asare-Kumi AA, Nortey EN, Kodom EB. Profiling patients healthcare quality perceptions: A baseline study of the University of Ghana hospital. Health Science Journal. 2016;10(3):1.
15. Castner D. Management of patients on hemodialysis before, during, and after hospitalization: challenges and suggestions for improvements. Nephrology nursing journal. 2011 Jul 1;38(4):319.
16. Dr. suresh k sharma . nursing research and statistics. New delhi: Elsevier publications ; p.p. no 93,99,114
17. Yousef YE, Elashir UM, Mahmoud SR, Maghraby N. The effect of nursing educational program on knowledge and practice of nurses regarding infection control measures for children under hemodialysis. Egyptian Nursing Journal. 2019 Jan 1;16(1):1.

18. Gomez NJ, editor. Nephrology nursing scope and standards of practice. American Nephrology Nurses Association; 2017.

UNDER PEER REVIEW