

## Original Research Article

# A CROSS SECTION ASSESSMENT STUDY ON THE AWARENESS OF COVID-19 DURING THE ONSET OF THE PANDEMIC AMONG THE URBAN POPULATION OF THE SAUDI ARABIA

### Abstract:

This study was aimed to seek out the awareness and knowledge about the covid-19 during the onset of the pandemic among the urban population of the Saudi arabia. This study was carried out for a period of three months from 01.04.2020 to 31.07.2020 and a sample of 1000 participants. The significant difference of P- value obtained by performing the Pearson Chi-Square test for all the questionnaires to analyse the study of assessment. This study was a small attempt for conducting such valuable studies in the preventive medicine. Thus, this study suggests that the overall satisfactory report that the community is well aware of the preventive measure procedures and well prepared to tackle the covid-19 pandemic.

**Keywords:** Covid-19, Urban population, Cross section assessment, Saudi Arabia, Pandemic

### Introduction

The corona virus which caused Covid-19 resulting in the pandemic outbreak which caused severe damage to the mankind by impacting on the both monetary and human lives throughout the world resulting as the one of the biggest disasters of the century<sup>1,2,3</sup>. This study was aimed to seek out the awareness and knowledge about the covid-19 during the onset of the pandemic among the urban population of the Saudi Arabia<sup>4,5,6</sup>. This study was carried out for a period of three months from 01.04.2020 to 31.07.2020 and a sample of 1000 participants were taken for the study by means of well structured online survey questionnaires. The data was collected in the form of well designed specific questionnaires related to the awareness of covid-19 along with other related informations related to the bio-informatics of the individual, etc both in English and Arabic language. This cross-sectional survey study was conducted by blending both qualitative and quantitative methods of data collection protocol. The collected data were analysed by the Microsoft Excel and SPSS. The values were presented as numbers (%). IBM SPSS Statistics for

Windows, version 23 was utilized for analysis (IBM SPSS, IBM Corp., Armonk, N.Y., USA). Statistical comparisons were made by Pearson Chi-Square test. The data were significant if P-values were  $< 0.05$ <sup>7,8</sup>. The Coronavirus which causes covid-19 impacts various individuals in different ways. The most individuals who were infected may have experienced gentle to mild ailment and recuperate without hospitalization. The most sought symptoms in the affected individuals include fever, cough, tiredness, loss of taste or smell, sore throat, headache, aches and pains, diarrhea, a rash on skin, or discolouration of fingers or toes, red or irritated eyes, etc<sup>9,10,11,12</sup>. However, the severity of the disease leads to the high mortality among the individuals and requires an immediate action to get the prescribed laboratory investigation to be done in the suspected individuals who were to be isolated in the hospitals in a separate wards to prevent the spread of the disease to the other patients<sup>1, 13,14</sup>. The lack of breathlessness leading to the severe pneumonia due to the virus possessed severe threat and the hospitalization becomes mandatory for such individuals to get the appropriate prescribed treatment. The World Health Organization has warned against self prescribed remedies for the treatment of covid-19 as it may cause more severe consequences. The WHO prompts individuals to ensure themselves as well as other people around you by knowing current realities to avoid potential risk and follow guidance given by your neighborhood wellbeing authority<sup>9,11</sup>.

### **Methodology:**

This study was carried out for a period of three months from 01.04.2020 to 31.07.2020 and a sample of 1000 participants were taken for the study by means of well structured online survey questionnaires. The data was collected in the form of well designed specific questionnaires related to the awareness of covid-19 along with other informations related to the bio-informatics of the individual, etc both in English and Arabic language. This cross-sectional survey study was conducted by blending both qualitative and quantitative methods of data collection protocol. The collected data were analysed by the Microsoft Excel and SPSS. The values were presented as numbers (%). IBM SPSS Statistics for Windows, version 23 was utilized for analysis (IBM SPSS, IBM Corp., Armonk, N.Y., USA). Statistical comparisons were made by Pearson Chi-Square test. The data were significant if P-values were  $< 0.05$ <sup>7,8,15</sup>

### **Results and Discussion**

The target population for the sample collection for this study was 1000 samples but received

overwhelming responses but we collected about 1000 samples participates in this survey as per the strategy. The collected data were well tabulated in the Table-1 and the appropriate responses were categorily separated and depicted in the form of well brewed chart figures 1 to 7 for better understanding about the explanation of the study.


**Table-1.** Assessment study data report of the awareness of covid-19 among the urban population of the saudi arabia

<b>Questionnaires</b>	<b>Response percentage</b>	<b>p-value significance</b>
<b>Distribution of Region</b>		<0.07
Riyadh	29	
Makkah	27	
Madina	13	
Eastern	15	
Others	16	
<b>Age groups</b>		<0.003
12 to 18	10	
19-24 years	24	
25 to 40 years	34	
41-60 years	27	
More than 60	5	
<b>Gender</b>		<0.087
Male	19	
Female	81	
<b>Nationality</b>		<0.06
Saudi	93	
Non-Saudi	7	
<b>Language</b>		<0.913
Arabic	95	
English	5	
<b>Education levels</b>		<0.002
Undergraduate	21	
Graduate	52	
Post Graduate	19	
Others	8	
<b>Do you Suffer from chronic disease</b>		<0.003
yes	27	
No	73	
<b>Do you take any medications</b>		<0.001
yes	53	
No	47	
<b>How worried are you to be infected with corona virus</b>		<0.02

worried	39.5	
Not worried	16	
Little bit worried	44.5	
<b>Do you think you are infected with corona virus</b>		<0.426
Yes	12.2	
No	24.5	
May be	63.3	
<b>Do you prepared to tackle pandemic</b>		<0.213
Yes	44	
No	34	
May be	22	
<b>Are you regular in wearing face mask</b>		<0.133
Yes	77	
No	09	
May be	14	
<b>Do you apply social distance</b>		<0.346
Yes	53	
No	34	
May be	13	
<b>Propability of you or any one of you know him is infected with corona</b>		<0.001
High propability	11.2	
Medium probability	66.8	
No propability	16.5	
Never any one will be infected	5.5	
<b>What do you think the fatality rate of corona patients</b>		<0.202
Less than 10 %	56	
Between 10 to 25%	28.8	
25 to 50%	9.7	
More than 50 %	5.5	
<b>What is the rate of corona patients suffer from mild symptoms</b>		<0.001
Less than 10 %	15	
Between 10 to 25%	24	
25 to 50%	26	
More than 50 %	35	
<b>What are the symptoms of corona virus do you think</b>		<0.04
High fever	27	
Dyspnea and chest pain	9	
Sore throat	11	
Anosomia	8	
Dry cough	22	
Diarrhea	5	
Fatigue, Body ache&Headache	12	


Rash or change color of finger nails	6	
<b>How can you protect yourself from corona</b>		<0.325
Wearing a face mask	35	
Stay at home	22	
Disinfecting Hand	11	
Social Distancing	27	
others	5	
<b>What will be the impact rate of the virus corona in public health in the scale of 1 to 10</b>		<0.28
1	2.5	
2	1.5	
3	2.6	
4	5.5	
5	14.6	
6	9.2	
7	12.8	
8	16.4	
9	6.8	
10	27.9	

This study was conducted among the urban population of the different regions of the Kingdom of Saudi Arabia which includes Riyadh, Makkah, Madina, Eastern province cities and other urban areas of the region. The study shows Makkah and Riyadh participants were in large numbers with the participants of Makkah were 27%, Riyadh were 29%, Madina were 13%, Eastern province were 15% and other regional urban areas were 16%. The distribution of the participants was described in the form of chart figure-1. Symbolically with the P-value obtained by performing the Person Chi-Square test was < 0.07 for the Makkah and Riyadh when compared to the other regions which shown the significant differences in the participants in the region studied.



**Fig. 1. Distribution of the region**


The general bio-information survey questionnaires were conducted based upon the nationality, age group, gender, language and education and was depicted in the form of chart figure-2 based on the participants. The most of the participants were Saudi nationals with 93% to the 7% non – Saudi participant with the clear significant difference of  $< 0.06$  P-value was obtained by performing the Pearson Chi-Square test. The most that participated in the survey group of age 25-40 years which were 34%, with 27% in the age group of 41-60 and 24.% were between 19-24 where as 10% were between 12-18 years old, while 5% participants were more than 60 years old respectively. The achieved significant difference in terms of the P- value obtained by performing the Pearson Chi-Square test was  $< 0.003$ . Regarding the gender distribution about 81% were female participants and 19%. were male participants with the P- value obtained by performing the Pearson Chi-Square test was  $< 0.087$ . There was a clear significant difference in the P- value obtained by performing the Pearson Chi-Square test which was  $< 0.913$  regarding the language distribution of the participants with 95% were Arabic and only 5% were English participants. The distribution of education level showed that the large group of participants were were graduate with 52%. followed by under graduate with 21%, post graduate 19% and others with 8% with the P- value obtained by performing the Pearson Chi-Square test was  $< 0.002$ .



**Fig. 2. General bio-information**

The set of public awarenessbased questionnaires about the onset of covid-19 pandemic were described in the form of the chart figure-3 based on the percentage of response received. Regarding the questionnaire “Do you Suffer from chronic disease”, the response was 27% participants answered Yes while 73% answered No while none answered May be with the significant difference of  $< 0.003$  P- value obtained by performing the Pearson Chi-Square test. The P-value of  $< 0.001$  was obtained by performing the Pearson Chi-Square test for the continuation to the previous questionnaire “Do you take any medications”which received the response rate of 53% answered Yes while 47% answered No. For answering to the questionnaire “How worried are you to be infected with corona virus”, the response answers of 39.5% felt worried with 16 % not worried and little worried were 44.5% was achieved with the significant difference of  $< 0.02$  P- value obtained by performing the Pearson Chi-Square test for the questionnaire. The next questionnaire in this category was about “Do you think you are infected with corona virus” for which the response answer for Yes was 12.2% while No was 24.5% and May be was 63.3% with the significant difference of  $< 0.426$  P- value obtained by performing the Pearson Chi-Square test. The important questionnaire of this category was “Do you prepared to tackle pandemic” for which the significant difference of P- value obtained by performing the Pearson Chi-Square test for the questionnaire was  $< 0.213$  with majority of the participants


answered Yes with 44% while No was 34% and May be with 22%. The second last questionnaire of this category was “Are you regular in wearing face mask” for which the significant difference of P- value obtained by performing the Pearson Chi-Square test for the questionnaire was < 0.133 with majority of the participants answered Yes with 77% while No was 09% and May be with 14% answers respectively. The final questionnaire in this category of questionnaire was regarding “Do you apply social distance ” for which a significant 53% answered Yes while 34% answered No while may be was 13% with the significant difference of P- value obtained by performing the Pearson Chi-Square test for the questionnaire was < 0.346.



**Fig. 3. Public awareness based questionnaire percentage**

A category of questionnaires were designed specifically based upon the presumption and probability based scenario towards the target population which were presented in the chart figure-4 based upon the percentage of responses achieved for each. The reponse for the first questionnaire of this category was based on the questionnaire “Probability of you or any one of you know him is infected with corona” for which answer for the option high probability was 11.2% while for medium probability was 66.8% and less probability was 16.5 with never any one will be infected was 5.5% with the significant difference of P- value obtained by performing the Person Chi-Square test for the questionnaire was < 0.001. The second questionnaire of this


category was “What do you think the fatality rate of corona patients” for which the significant difference of P- value obtained by performing the Pearson Chi-Square test for the questionnaire was  $< 0.202$  with majority of the participants answered for less than 10 % with 56% while for 10 to 25% was 28.8% and for 25 to 50 % with 9.7% while for more than 50% it was 5.5% answers respectively. The next target questionnaire of this category was “What is the rate of corona patients suffer from mild symptoms“ for which the response answer of less than 10% was 15% while 10 to 25% was 24% with 25 to 50% was 26% and more than 50% was with 35% with the significant difference of  $< 0.001$  P- value obtained by performing the Pearson Chi-Square test.



**Fig. 4. Probability scenario based questionnaire percentage**


A questionnaire was framed to study the category about the awareness of the general covid-19 symptoms to assess the knowledge of the participants and the percentage of the results were described in the chart Figure-5. The framed questionnaire towards the target population was “What are the symptoms of corona virus do you think” for which the response for the High fever symptom was 27% which out performed other symptoms due to its regularity associated with majority of the infections. The other responses were 9% for Dyspnea or chest pain while sore throat was 11% with the anosomnia received 8% and dry cough symptom response received 22%. The response for diarrhea symptom received 5% and Fatigue, body ache or Head ache

received 12% while 6% response was received for the symptom of rash or change of color of the finger. The P- value obtained by performing the Pearson Chi-Square test was  $< 0.04$  which was clear significant difference




**Fig. 5. Public awareness about general Covid-19 symptoms questionnaire percentage**

A questionnaire was framed to study the category about the awareness of the general covid-19 precautionary measures to assess the knowledge of the participants and the percentage of the results were described in the chart Figure-6. The framed questionnaire towards the target population was “How can you protect yourself from corona” for which the response for the wearing the mask was 35% which out performed other responses due to its popularity associated with majority of the infections and ease of availability. The other responses were 22% for Stay at home while Disinfecting hand received 11% response with the social distancing received received 27% and other responses received mere just 5%. The P- value obtained by performing the Pearson Chi-Square test was  $< 0.325$  which was clear significant difference



**Fig. 6. Awareness about self protection questionnaire percentage**

The final questionnaire category of this cross-sectional study was specifically designed to study the impact of the covid-19 on the public health which was analysed by the scale rating 1 to 10 and the response percentage was depicted in the form of chart Figure-6. This questionnaire was strategically placed at the end of the cross-sectional survey to create the awareness among the target population regarding the awareness of the target population to understand the psychological condition of the participants towards the onset of the pandemic condition. The questionnaire designed was “What will be the impact rate of the virus corona in public health in the scale of 1 to 10”. The procured responses for this questionnaire were majority of the participants felt the impact rate of 10 with 27.9% while the impact rate of 5,7 & 8 received almost the same percentage responses of moderate 14.6%, 12.8% and 16.4% respectively. The mere percentage of responses were received for the other impact rate scales of 1,2,3,4,6 and 9 with the response percentages of 2.5%, 1.5%, 2.6%, 5.5%, 9.2% and 6.8% respectively. The significant difference of P-value obtained by performing the Pearson Chi-Square test for the questionnaire category was  $< 0.028$ .



**Fig. 7. Impact rate scale of Covid-19 on public health**

## **Conclusion**

In this cross section assessment study received responses regarding the well formulated questionnaires regarding prevalence of awareness of covid-19 during the onset of the pandemic situation among the urban population of the Kingdom of Saudi Arabia it was observed that the community was well aware of the knowledge regarding the covid-19 situation. An overwhelming satisfactory responses were received from the target population and the data were limited to the planned 1000 participants for the study was procured. The collected data were well tabulated in the Table-1 and depicted in the form of chart figures 1 to 7 respectively. The significant difference of P-value obtained by performing the Pearson Chi-Square test for all the questionnaires to analyse the study of assessment. This conducted study was well satisfactory and more such studies need to be conducted to assess the psychological factors of the community to support the valuable efforts taken by the authorities which clearly suggests the most encouraging facts. This study was a small attempt for conducting such valuable studies in the preventive medicine. Thus, this study suggests that the overall satisfactory report that the community is well aware of the preventive measure procedures and well prepared to tackle the covid-19 pandemic.

**COMPETING INTERESTS DISCLAIMER:**

**Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.**

## References

1. CDC. When to quarantine; March 12, 2021. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/quarantine.html>. Accessed June 28, 2021.
2. World Health Organization. Considerations for quarantine of contacts of COVID-19 cases; June 25, 2021. Available from: <https://www.who.int/publications/i/item/WHO-2019-nCoV-IHR-Quarantine-2021.1>. Accessed June 20, 2021.
3. World Health Organization. WHO virtual press conference on COVID-19; March 11, 2020. Available from: [https://www.who.int/docs/default-source/coronaviruse/transcripts/who-audio-emergencies-coronavirus-press-conference-full-and-final-11mar2020.pdf?sfvrsn=cb432bb3\\_2](https://www.who.int/docs/default-source/coronaviruse/transcripts/who-audio-emergencies-coronavirus-press-conference-full-and-final-11mar2020.pdf?sfvrsn=cb432bb3_2). Accessed March 16, 2020.
4. Bedford J, Enria D, Giesecke J, et al. COVID-19: towards controlling of a pandemic. *Lancet*. 2020;395(10229):1015–1018. doi:10.1016/S0140-6736(20)30673-5
5. Ministry of Health. What do you need to know if you get infected with COVID-19?; July 28, 2020. Available from: <https://covid19awareness.sa/wp-content/uploads/2020/07/WhatToDoWhenGettingInfected.pdf>. Accessed June 28, 2021.
6. Alahdal, H.; Basingab, F.; Alotaibi, R. An analytical study on the awareness, attitude and practice during the COVID-19 pandemic in Riyadh, Saudi Arabia. *J. Infect. Public Health* 2020, 13, 1446–1452.

7. Bäuerle, A.; Teufel, M.; Musche, V.; Weismüller, B.; Kohler, H.; Hetkamp, M.; Dörrie, N.; Schweda, A.; Skoda, E.-M. Increased generalized anxiety, depression and distress during the COVID-19 pandemic: A cross-sectional study in Germany. *J. Public Health* 2020, *42*, 672–678
8. Algahtani, F.D.; Zrieq, R. Avoiding medical visits even when needed during the COVID-19 pandemic: A cross-sectional survey from Saudi Arabia. *Syst. Rev. Pharm.* 2020, *11*, 649–654.
9. World Health Organization. Clinical management of severe acute respiratory infection (SARI) when COVID-19 disease is suspected; 2020. Accessed 16 April 2020.
10. WHO. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19). Available online: <https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf> (accessed on 15 January 2021).
11. World Health Organization, Coronavirus Disease (COVID-19) Pandemic. 2020. Available online: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> (accessed on 15 January 2021).
12. World Health Organization. Clinical Management of Severe Acute Respiratory Infection When Novel Coronavirus (2019-nCoV) Infection Is Suspected: Interim Guidance, 28 January 2020. Art. no. WHO/nCoV/Clinical/2020.3. 2020.
13. World Health Organization. Critical preparedness, readiness and response actions for COVID-19; March 7, 2020. Available from: <https://apps.who.int/iris/handle/10665/331422>. Accessed March 13, 2020.
14. Chudasama, Y.V.; Gillies, C.L.; Zaccardi, F.; Coles, B.; Davies, M.J.; Seidu, S.; Khunti, K. Impact of COVID-19 on routine care for chronic diseases: A global survey of views from healthcare professionals. *Diabetes Metab. Syndr. Clin. Res. Rev.* 2020, *14*, 965–967.
15. Alyami, H.S.; Orabi, M.A.; Aldhabbah, F.M.; Alturki, H.N.; Aburas, W.I.; Alfayez, A.I.; Alharbi, A.S.; Almasuood, R.A.; Alsuhaibani, N.A. Knowledge about COVID-19 and Patients Beliefs about and Use of Herbal Products during the COVID-19 Pandemic: A Cross-Sectional Study in Saudi Arabia. *medRxiv* 2020

UNDR PEER REVIEW