

COMPARISON OF RANDOM BLOOD SUGAR AND pH OF BLOOD- A CLINICOPATHOLOGICAL STUDY.

Running Title: Comparison of random blood sugar and pH of blood -A clinicopathological study.

ABSTRACT:

Aim:

This study aims to compare the random blood sugar pH of the blood.

Introduction:

Hyperglycemia is a risk factor for the impairment of periodontal health while periodontal disease, in turn, is responsible for the adverse effects on glycemic control and the pathophysiology of diabetes-related complications. An increase in blood sugar levels has a direct effect on pH with respect to age. Hence this study deals with the association of random blood sugar levels and pH with age, gender, and dental complaints.

Materials and Method:

20 random blood samples were collected from the patients in Saveetha Dental College. The random blood sugar level was analyzed by the semi-quantitative method and the pH was analyzed by using a multi reagent strip. And the results were statistically analyzed using SPSS software and represented as pie charts and bar graphs.

Comment [A1]: Write 20 after blood samples

Results and Discussion:

The overall mean value of RBS and blood pH was found to be 138.4 mg/dl and 7.06 respectively. The mean RBS value is higher in the 41-50 years age group of people than in other age categories and the mean pH value is same for all age categories. In the present study,

samples with periodontic complaints have higher mean value of RBS values(193) than other dental complaints and the mean pH value is same for all dental complaints

Conclusion: Within the limits of the study, it can be observed that there is a significant association between the age group, gender, and dental complaints with RBS and pH of the patients. However, further studies ~~have to~~ must be conducted to have more accurate results and elicit the association between pH and RBS with age and gender.

Keywords: Random blood sugar, Blood pH, Dental complaints, Hyperglycemia, ~~innovative~~ innovative technique.

INTRODUCTION:

pH is a key criterion for evaluating body homeostasis. Hemoglobin and albumin are high pH buffering molecules in the blood, and the pH of blood remains around 7.35 and 7.45 on the pH scale.(1,2). Disturbance of the mechanisms which maintain normal blood pH is a defining feature of diabetic ketoacidosis (DKA), an acute and life-threatening complication of the chronic metabolic disease, diabetes(3,4), and Hyperglycemia. A complex synergy of action involving the lungs, kidneys, and the blood's bicarbonate buffering system maintains the pH of blood (ECF) within very narrow limits (7.35-7.45).(5,6).

Normal metabolism is associated with continuous production of hydrogen ions (H⁺) which tend to reduce pH, but bicarbonate in the blood (ECF) combines with these hydrogen ions forming carbonic acid. CO₂ is eliminated from the body by the lungs and bicarbonate is regenerated in the kidneys. By this means, the concentration of hydrogen ions in the blood, and therefore the pH, remains constant despite the continuous production of metabolic acids(7,8). The abnormal accumulation of ketoacids in the blood that occurs in DKA overwhelms this buffering system. In the process of DKA, pH is less because the bicarbonate buffer is exhausted, i.e., bicarbonate concentration is reduced. So, the compensatory mechanism invoked by DKA is increased respiration(9–11). Kussmaul's respiration, in patients suffering from DKA(12,13), provides clinical evidence of the compensatory mechanism for blood pH. The decreased *p*CO₂ that results from this increased respiration returns the pH towards normal but may not be

sufficient to achieve a normal pH(14–16). The patient ~~is considered to be~~ fully compensated if the blood pH is within the normal range or partially compensated if the blood pH remains below the lower limit of the normal values(12,17). The three main components of DKA treatment are insulin administration, replacement of lost fluid with an intravenous injection of isotonic saline, and replacement of potassium losses(18). Bicarbonate infusion is often used to deal with acidosis correction(19,20).

Careful and intensive monitoring of blood biochemistry (particularly blood glucose, potassium, and pH) and fluid balance is essential for a favorable outcome_(21–23). Our team has extensive knowledge and research experience –that has translated into high quality publications (13)_(2,4,6,10,12,16,19,24–30) ,(31–35)_.

Hence, the study aims to compare the Random Blood Sugar levels and pH of blood among the population.

MATERIALS AND METHOD:

20 rRandom blood samples (20) were collected from the patients in Saveetha Dental College and the low sample size is one of the limitations in this study. The Random blood sugar for the 20 samples was determined by a semi-quantitative method. The blood pH was analyzed from the samples using a multi-reagent strip and the pH was determined. The pH and RBS of the patients were tabulated and were statistically analyzed using SPSS software version (23.0) and represented as pie charts and bar graphs_(34). A correlation test was done to check the association between RBS and pH with Gender, age group, and dental complaints.

RESULTS AND DISCUSSION:

The following graphs indicate the correlation between RBS and pH with Gender, age group, and dental complaints.




Figure 1: Bar graph represents the correlation between RBS and pH with the Gender. The X-axis represents the gender, and the Y-axis represents the mean value of RBS and pH. The green color represents pH, and the blue color represents RBS. It is analyzed that the mean RBS value is higher in males than females and the mean pH value is the same for males and females. A correlation test was done, and the association was found to be statistically significant with Pearson's correlation value ($r = 1$).



Figure 2: Bar graph represents the correlation between RBS and pH with age. The X-axis represents the age, and the Y-axis represents the mean value of RBS and pH. The green color represents pH, and the blue color represents RBS. It is analyzed that the mean RBS value is higher in the 41-50 years age group of people than in other age categories and the mean pH value is same for all age categories. Pearson's correlation test was done, and the association was found to be statistically significant with Pearson's correlation value of (r=1).



Figure 3: Bar graph represents the correlation between RBS and pH with the dental complaints. The X-axis represents the dental complaints, and the Y-axis represents the mean value of RBS and pH. The green color represents pH, and the blue color represents RBS. Samples with periodontic complaints have higher mean value of RBS values(193) than other dental complaints and the mean pH value is same for all dental complaints. Pearson's correlation test was done, and the association was found to be statistically significant with Pearson's correlation value of (r=1).

In our study, it was found that 2(13%) of the study population were aged between 21-30 years, 5(33%) of them between 31-40 years, 5(33%) of them between 41-50 years and the remaining 3(20%) of the study population were above 50 years. Gender distribution shown in this study was 6(40%) males and 9(60%) females. The study population showed 6(40%) out of 15 with endodontic complaints, 5(33%) with periodontal complaints, 2(13%) with surgical complaints, 2(13%) with neuralgic pain. The Random Blood Sugar value and pH of the blood were calculated for all the participants and that showed an overall mean value of RBS and blood pH was found to be 138.4 mg/dl and 7.06 respectively.

Comment [A2]: Check spaces

The mean pH for the age group between 21-30 years was 7.0, between 31-40 years was 7.0, the age group between 41-50 years was 7.13, and the mean pH of the age group more than 50 years was 7.10. Results of this study show that blood pH values are positively correlated with the age group ($r=1$). The mean value of pH was almost same in all the age groups. It showed that when the age group increases the pH of the blood also slightly moves towards a higher end. In the present study, the mean value of pH was almost equal both in males and females. In a previous study with similar findings, RBSRBS, and the prevalence of glycosuria among men were much higher than females with respect to glucose tolerance tests(36–38).

In the case of dental complaintscomplaints, the mean value of RBS was higher in the 41-50-year-old age group(190 mg/dl) when compared to the other age groups. The mean value of RBS was higher(168 mg/dl) in males when compared to females (119 mg/dl). With respect to dental complaints in the present study, the mean value of RBS for periodontic complaints, neuralgic pain, endodontic complaints, and surgical complaints were 193mg/dl, 126 mg/dl, 112 mg/dl, and 95 mg/dl respectively. In a study performed by K almas et al.2001, the subjects who had high random blood sugar levels were compared against their Community Periodontal Index of Treatment Needs score(CPITN) score and the results showed that periodontal disease severity was high among subjects who had high RBS levels than the rest of the subjects. The mean CPITN score was then compared with the mean RBS level and the severity of periodontal disease present. There was a steady increase in the RBS level with an increase in CPITN scores, i.e., CPITN score was found to be 13.5 to 19.12 with the RBS values of 184.2 and 199.12 mg/dl among patients who had comparatively high RBS (31,35,39).

Comment [A3]: Check spaces. Do for all

It can be observed that the mean value of RBS was comparatively higher in case of periodontal problems in males between the age group of 41-50 years which is in accordance with studies that suggests hyperglycemia as a risk factor for the impairment of periodontal health while periodontal disease(40–42), in turn, is responsible for the adverse effects on glycemic control and the pathophysiology of diabetes-related complications(18,33,43).

Limitations of this study include the less sample size and the other parameters which affect the pH were not considered. Future studies may be done to overcome these limitations.

CONCLUSION:

Within the limits of the study, it can be observed that there is a significant association between the age group, gender, and dental complaints with RBS and pH of the patients. The average mean value for RBS is 138.4mg/dl and for pH is 7.06. However, further studies have to be conducted to have more accurate results and elicit the association between pH and RBS with age and gender.

REFERENCES:

1. Mieno H, Marunaka Y, Inaba T, Kojima K, Yoneda K, Nagata K, et al. pH balance and lactic acid increase in the vitreous body of diabetes mellitus patients. *Exp Eye Res.* 2019 Nov;188:107789.
2. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. *Arch Oral Biol.* 2021 Feb;122:105030.
3. Jeng CY. Glucagon does not increase plasma free fatty acid and glycerol concentrations in patients with noninsulin-dependent diabetes mellitus [Internet]. Vol. 77, *Journal of Clinical Endocrinology & Metabolism.* 1993. p. 6–10. Available from: <http://dx.doi.org/10.1210/jc.77.1.6>
4. Girija SA, Priyadharsini JV, Paramasivam A. Prevalence of carbapenem-hydrolyzing OXA-

Comment [A4]: Check all references for similar pattern

type β -lactamases among *Acinetobacter baumannii* in patients with severe urinary tract infection. *Acta Microbiol Immunol Hung*. 2019 Dec 9;67(1):49–55.

5. Marunaka Y. The Proposal of Molecular Mechanisms of Weak Organic Acids Intake-Induced Improvement of Insulin Resistance in Diabetes Mellitus via Elevation of Interstitial Fluid pH. *Int J Mol Sci* [Internet]. 2018 Oct 19;19(10). Available from: <http://dx.doi.org/10.3390/ijms19103244>
6. Priyadharsini JV, Paramasivam A. RNA editors: key regulators of viral response in cancer patients. *Epigenomics*. 2021 Feb;13(3):165–7.
7. Rodríguez-Gutiérrez R, Cámara-Lemarroy CR, Quintanilla-Flores DL, González-Moreno EI, González-Chávez JM, Lavallo-González FJ, et al. Severe Ketoacidosis ($\text{pH} \leq 6.9$) in Type 2 Diabetes: More Frequent and Less Ominous Than Previously Thought [Internet]. Vol. 2015, *BioMed Research International*. 2015. p. 1–5. Available from: <http://dx.doi.org/10.1155/2015/134780>
8. Kumar SP, Praveen Kumar S, Smiline Girija AS, Vijayashree Priyadharsini J. Targeting NM23-H1-mediated Inhibition of Tumour Metastasis in Viral Hepatitis with Bioactive Compounds from *Ganoderma lucidum*: A Computational Study [Internet]. Vol. 82, *Indian Journal of Pharmaceutical Sciences*. 2020. Available from: <http://dx.doi.org/10.36468/pharmaceutical-sciences.650>
9. Newton CA, Raskin P. Diabetic ketoacidosis in type 1 and type 2 diabetes mellitus: clinical and biochemical differences. *Arch Intern Med*. 2004 Sep 27;164(17):1925–31.
10. Mathivadani V, Smiline AS, Priyadharsini JV. Targeting Epstein-Barr virus nuclear antigen 1 (EBNA-1) with *Murraya koengii* bio-compounds: An in-silico approach. *Acta Virol*. 2020;64(1):93–9.
11. Jaisankar AI, Smiline Girija AS, Gunasekaran S, Vijayashree Priyadharsini J. Molecular characterisation of *csgA* gene among ESBL strains of *A. baumannii* and targeting with essential oil compounds from *Azadirachta indica* [Internet]. Vol. 32, *Journal of King Saud University - Science*. 2020. p. 3380–7. Available from:

<http://dx.doi.org/10.1016/j.jksus.2020.09.025>

12. Jayaseelan VP, Paramasivam A. Emerging role of NET inhibitors in cardiovascular diseases. *Hypertens Res.* 2020 Dec;43(12):1459–61.
13. Anita R, Paramasivam A, Priyadharsini JV, Chitra S. The m6A readers YTHDF1 and YTHDF3 aberrations associated with metastasis and predict poor prognosis in breast cancer patients. *Am J Cancer Res.* 2020 Aug 1;10(8):2546–54.
14. Muhammad R. Diabetic ketoacidosis: clinical and biochemical parameters in patients with type 2 diabetes [Internet]. Available from: <http://dx.doi.org/10.26226/morressier.59d51843d462b80296ca371e>
15. Timothy CN, Samyuktha PS, Brundha MP. Dental pulp Stem Cells in Regenerative Medicine--A Literature Review. *Research Journal of Pharmacy and Technology.* 2019;12(8):4052–6.
16. Jayaseelan VP, Ramesh A, Arumugam P. Breast cancer and DDT: putative interactions, associated gene alterations, and molecular pathways. *Environ Sci Pollut Res Int.* 2021 Jun;28(21):27162–73.
17. Sivakumar S, Smiline Girija AS, Vijayashree Priyadharsini J. Evaluation of the inhibitory effect of caffeic acid and gallic acid on tetR and tetM efflux pumps mediating tetracycline resistance in *Streptococcus* sp., using computational approach [Internet]. Vol. 32, *Journal of King Saud University - Science.* 2020. p. 904–9. Available from: <http://dx.doi.org/10.1016/j.jksus.2019.05.003>
18. Girija ASS, Smiline Girija AS. Delineating the Immuno-Dominant Antigenic Vaccine Peptides Against gacS-Sensor Kinase in *Acinetobacter baumannii*: An in silico Investigational Approach [Internet]. Vol. 11, *Frontiers in Microbiology.* 2020. Available from: <http://dx.doi.org/10.3389/fmicb.2020.02078>
19. Anchana SR, Girija SAS, Gunasekaran S, Priyadharsini VJ. Detection of csgA gene in carbapenem-resistant *Acinetobacter baumannii* strains and targeting with *Ocimum sanctum* biocompounds. *Iran J Basic Med Sci.* 2021 May;24(5):690–8.

20. As SG, Vijayashree PJ, Paramasivam A. Prevalence of Acb and non-Acb complex in elderly population with urinary tract infection (UTI) [Internet]. Vol. 76, Acta Clinica Belgica. 2021. p. 106–12. Available from: <http://dx.doi.org/10.1080/17843286.2019.1669274>
21. Chua HR, Schneider A, Bellomo R. Bicarbonate in diabetic ketoacidosis - a systematic review. *Ann Intensive Care*. 2011 Jul 6;1(1):23.
22. Fujikawa K, Numasaki H, Kobayashi M, Sugano N, Tomura S, Murai S. pH determination in human crevicular fluids. Examination of the pH meter and evaluation of the correlation between pH level and clinical findings or the microflora in each periodontal pocket [Internet]. Vol. 31, *Nihon Shishubyo Gakkai Kaishi (Journal of the Japanese Society of Periodontology)*. 1989. p. 241–8. Available from: <http://dx.doi.org/10.2329/periodo.31.241>
23. Girija ASS, Smiline Girija AS, Shoba G, Vijayashree Priyadharsini J. Accessing the T-Cell and B-Cell Immuno-Dominant Peptides from *A.baumannii* Biofilm Associated Protein (bap) as Vaccine Candidates: A Computational Approach [Internet]. Vol. 27, *International Journal of Peptide Research and Therapeutics*. 2021. p. 37–45. Available from: <http://dx.doi.org/10.1007/s10989-020-10064-0>
24. Sivakumar S, Smiline Girija AS, Vijayashree Priyadharsini J. Evaluation of the inhibitory effect of caffeic acid and gallic acid on tetR and tetM efflux pumps mediating tetracycline resistance in *Streptococcus* sp., using computational approach. *Journal of King Saud University - Science*. 2020 Jan 1;32(1):904–9.
25. Smiline Girija AS. Delineating the Immuno-Dominant Antigenic Vaccine Peptides Against gacS-Sensor Kinase in *Acinetobacter baumannii*: An in silico Investigational Approach. *Front Microbiol*. 2020 Sep 8;11:2078.
26. Iswarya Jaisankar A, Smiline Girija AS, Gunasekaran S, Vijayashree Priyadharsini J. Molecular characterisation of csgA gene among ESBL strains of *A. baumannii* and targeting with essential oil compounds from *Azadirachta indica*. *Journal of King Saud University - Science*. 2020 Dec 1;32(8):3380–7.

27. Girija ASS. Fox3+ CD25+ CD4+ T-regulatory cells may transform the nCoV's final destiny to CNS! *J Med Virol* [Internet]. 2020 Sep 3; Available from: <http://dx.doi.org/10.1002/jmv.26482>
28. Kumar SP, Girija ASS, Priyadharsini JV. Targeting NM23-H1-mediated inhibition of tumour metastasis in viral hepatitis with bioactive compounds from *Ganoderma lucidum*: A computational study. *pharmaceutical-sciences* [Internet]. 2020;82(2). Available from: <https://www.ijpsonline.com/articles/targeting-nm23h1mediated-inhibition-of-tumour-metastasis-in-viral-hepatitis-with-bioactive-compounds-from-ganoderma-lucidum-a-comp-3883.html>
29. Girija As S, Priyadharsini J V, A P. Prevalence of Acb and non-Acb complex in elderly population with urinary tract infection (UTI). *Acta Clin Belg*. 2021 Apr;76(2):106–12.
30. Girija ASS, Shoba G, Priyadharsini JV. Accessing the T-Cell and B-Cell Immuno-Dominant Peptides from *A.baumannii* Biofilm Associated Protein (bap) as Vaccine Candidates: A Computational Approach. *Int J Pept Res Ther*. 2021 Mar 1;27(1):37–45.
31. Arvind P TR, Jain RK. Skeletally anchored forsus fatigue resistant device for correction of Class II malocclusions-A systematic review and meta-analysis. *Orthod Craniofac Res*. 2021 Feb;24(1):52–61.
32. Venugopal A, Vaid N, Bowman SJ. Outstanding, yet redundant? After all, you may be another Choluteca Bridge! *Semin Orthod*. 2021 Mar 1;27(1):53–6.
33. Ramadurai N, Gurunathan D, Samuel AV, Subramanian E, Rodrigues SJL. Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial. *Clin Oral Investig*. 2019 Sep;23(9):3543–50.
34. Varghese SS, Ramesh A, Veeraiyan DN. Blended Module-Based Teaching in Biostatistics and Research Methodology: A Retrospective Study with Postgraduate Dental Students. *J Dent Educ*. 2019 Apr;83(4):445–50.
35. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding

gingival inflammation in primary molars: randomized controlled trial [Internet]. Vol. 24, Clinical Oral Investigations. 2020. p. 3275–80. Available from: <http://dx.doi.org/10.1007/s00784-020-03204-9>

36. Apoorva SM, Sridhar N, Suchetha A. Prevalence and severity of periodontal disease in type 2 diabetes mellitus (non-insulin-dependent diabetes mellitus) patients in Bangalore city: An epidemiological study. J Indian Soc Periodontol [Internet]. 2013 Jan [cited 2021 Mar 11];17(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/23633768/>
37. Harsha L, Brundha MP. Prevalence of Dental Developmental Anomalies among Men and Women and its Psychological Effect in a Given Population. Journal of Pharmaceutical Sciences and Research; Cuddalore. 2017 Jun 20;9(6):869–73.
38. Brundha MP, Pathmashri VP, Sundari S. Quantitative Changes of Red Blood cells in Cancer Patients under Palliative Radiotherapy-A Retrospective Study. Research Journal of Pharmacy and Technology. 2019;12(2):687–92.
39. Almas K, Al-Qahtani M, Al-Yami M, Khan N. The relationship between periodontal disease and blood glucose level among type II diabetic patients. J Contemp Dent Pract. 2001 Nov 15;2(4):18–25.
40. Su L, Liu W, Xie B, Dou L, Sun J, Wan W, et al. Toothbrushing, Blood Glucose and HbA1c: Findings from a Random Survey in Chinese Population. Sci Rep [Internet]. 2016 Jul 7 [cited 2021 Mar 11];6. Available from: <https://pubmed.ncbi.nlm.nih.gov/27385509/>
41. Venugopal A, Vaid N, Jay Bowman S. Outstanding, yet redundant? After all, you may be another Choluteca Bridge! [Internet]. Vol. 27, Seminars in Orthodontics. 2021. p. 53–6. Available from: <http://dx.doi.org/10.1053/j.sodo.2021.03.007>
42. Girija ASS, Smiline Girija AS. Fox3 CD25 CD4 T- regulatory cells may transform the nCoV's final destiny to CNS! [Internet]. Vol. 93, Journal of Medical Virology. 2021. p. 5673–5. Available from: <http://dx.doi.org/10.1002/jmv.26482>
43. Kim EK, Lee SG, Choi YH, Won KC, Moon JS, Merchant AT, et al. Association between diabetes-related factors and clinical periodontal parameters in type-2 diabetes mellitus.

BMC Oral Health [Internet]. 2013 Nov 7 [cited 2021 Mar 11];13. Available from:
<https://pubmed.ncbi.nlm.nih.gov/24195646/>

UNDER PEER REVIEW

