

AROMATIC PROFILING AND BIOACTIVE POTENTIAL OF *LAVENDULA ANGUSTIFOLIA*: A REVIEW

ABSTRACT

Plants are the vital source around which human life revolves. Starting from our food to medicine we obtain, all such valuable resources come from resilience creatures called plants. Plants are gifted with certain properties that make them precious in all aspects of life. They produce bioactive substances that make them medicinally important for pharmaceutical use. The aromatic plants which are important as industrial raw material are used as essence material in various baking and brewing industries because of their pleasant aroma. These plants undergo distillation processes to extract the essential oils from oil secreting glands. The present review deals with aromatic profiling and bioactive potential of *Lavendula spp* and their commercial importance as a plant material.

Keywords: Aromatic, Bioactivity, Lavender, essential oil.

INTRODUCTION:

Aromatic plants are significant and great importance because of their wide commercial uses. Growing use of aromatic flowers as cut flowers and their use in perfumery products lead to their flourishing as a raw material. Angiosperms are distributed all around the globe and there are about 3,00,00 spp and 427 families around globe while in India about 251 angiosperm families exist as per present statistics of BSI, Kolkata. Angiosperms are plants that are rich in aroma while it contain certain families rich in essential oils such as Lamiaceae, Lauraceae, Myricaceae etc. to name a few. The largest genera include *Salvia*, *Stachys*, *Plectranthus*, *Scutellaria*, *Thymus*, *Vitex*, *Nepeta* [Kuethe *et.al*,2017]. The family is important as it has ornamental and condimental herbs which are commercially vital. The present plant belongs to one such family Labiatae which is also called sage or **mint** family. The important species of this genus that are *Lavendula latifolia*, *Lavendula angustifolia* Mill which are native to mediterranean region from Africa to Asia [Verma *et.al*,2010]. In India it is well marked from Utrakhand to Kashmir valley [Singh

et.al,2015]. The plant holds great importance since ancient period as Iranian folk medicine.[Sharif *et.al*,2003]

Classification

Kingdom:Plantae

Clade:Tracheophyta

Order:Lamiales

Family:Lamiaceae

Genus:*Lavendula*

Species:*angustifolia*

PLANT DESCRIPTION

The plant is an semi-woody, evergreen, perennial plant with lanceolate leaves covered by tomentum, square stem curled at edges. The inflorescence is spike like and has bright purple flowers that are important ornamental blooms. The plant is native to Spain, and some neighboring countries but now naturalized in India. This plant requires fertile loamy soil, and it is propagated vegetatively as well as by seed. The flowers have excellent flavour properties and sell as sachets. The plant possesses majestic flowers that are known for their aroma and used as

cut flowers and have various bioactive compounds of economic importance.



Fig. 1. Plant morphology

https://commons.wikimedia.org/wiki/File:Lavandula_angustifolia_002.JPG

BIOACTIVE POTENTIAL

Lavender augustifolia Mill. is a member of Lamiaceae which is a family rich in *polyphenols* and other bioactive substances. The flowers are the economical part of the plant that yield phenolic compounds and other active constituents. Cardia *et.al*,2018 used GC-MS chromatography and NMR spectroscopy to examine the bioactive compounds like 1,8 cineole, linalool, camphor which help to control acute inflammatory response and Lavender oil induced the leukocyte chemotaxis in vitro by just varying the concentration of bioactive compounds. Linalool, p-cymene, limonene and camphor were known impart the anti-inflammatory activity which was observed in lavender by lipoxygenase inhibitory activity as reported by Carrasco *et.al*,2015. Terpenoids found in Lavender oil are known to play predominant role in plant-insect communication which was examined by cytological investigation of TPS-TPS, TPS-CYP450 linked to substances that function as attractants and repellants [Li *et.al*,2021]. Characterization of the genetic diversity were examined in *Lavender angustifolia* Mill. using sequence related amplified polymorphism which showed higher genetical diversity as reported by Rusanov *et.al*,2020. Cristina *et.al*,2014 examined the volatile oils from the tea samples of *Lavendula angustifolia* Mill. using GC-MS spectroscopy. The dominant bioactives were Linalyl acetate and linalool which corresponds to 1.05% essential oil in the given species and it shows subsequent amounts of chlorogenic acid and Rosmarinic acid using its alcoholic extracts was reported [Ghelardini *et.al*,1999].




Fig. 2. Lavender oil properties

AROMATIC PROFILING IN *LAVENDULA ANGUSTIFOLIA*:

Lavandula angustifolia Mill is rich in various aromatic compound that makes it a majestically useful and mainstay herb in the gardens worldwide. It's also called as English lavender and it finds its place in all the restaurants and kitchens where it is used as an flavouring agent and for seasoning purposes and cosmetic industry. Wang *et.al*,2020 evaluated the volatile or odour characteristics [LE] with lavender essential oil [LEO] using Gas chromatography-Mass chromatography [GC-MS] and Principal chemical analysis [PCA].He found out the LE contains high amount of bioactive substances like lavendulyl acetate, linalool oxides I and II than LEO,and has herbal and clove like odour, whereas more earthy green and watery odour of LEO was due to higher amount of Linalool and camphor and finally he concluded that LE is better LEO in terms of odour characteristics. Methanolic extracts of Lavender were used to estimate flavone content by using UV-vis spectroscopy as reported by Margaoan *et.al*, 2017. Lazari *et.al*,2018 elucidated the antiviral and scavenging activity while Gezizi *et.al*,2018 reported the Anti-proliferative property of Essential oils. Roj *et.al*,2019 used Box-Behnken method to extract the phenolic compounds under controlled conditions and found highest phenolic content was found in *Lavender angustifolia*Mill.at 54.5°C.

Comment [U1]: #1 antiviral
#2 scavenging

Comment [U2]: #3 anti-proliferative

Table 1. Aromatic profiling

S.NO	Volatile compounds	total % in Lavender oil
------	--------------------	-------------------------

1	<i>α – terpineol</i>	1.35
2	Camphor	9.67
3	Sebinene	0.5
4	Myrcene	0.68
5	Limonene	0.24
6	1,8 cineole	8.6
7.	Linalool acetate	3.75
8.	Cryptone	0.98
9.	Terepiene-4-ol	3.8
10.	Linalool	47.55

Predoi *et.al* ,2018 examined the aromatic potential of plants based on the compounds present in it and using Gas chromatography they found that dominant compounds in the essential oil of basil and lavender are oxygenated monoterenes responsible for the characterstic essence. The other compounds were in trace amounts and the plant is researched in its infancy stage and other properties are yet to be unveiled. The antimicrobial activity of the *Lavendula aungustifolia* Mill was seen by Aires *et.al*, 2017 who used HPLC-DAD and minimum inhibitory concentration [MIC] to characterize phenolic substances and flavanoids in lavender and they found out that hydroxycinnamic acid [Chlorogenic acid, rosmarinic acid], flavonoids were responsible for high antibacterial activity against *Staphylococcus aureus* isolated from diabetic foot ulcers. Lavender due to its characterstic aroma and beauty was indiscriminately used which lead to its overexploitation and therefore certain strategies and tools under biotechnology were used to revive it such as micropropagation vis meristem proliferation and organogenesis as reported by Romanno *et.al* 2013. Ayaz *et.al*, 2017 examined the essential oils of lavender to provide protection against neural disorder like Alzhemier disease and dementia, it shows improved cognitive performance against epilepsy and convulsions in animals and human subjects and they are cheap, easy to obtain and safe to use with good efficiency. Oran *et.al*,2011 inferred the methanolic extracts of *Lavendula angustifolia* Mill. used to control the diabetic dyslipidemia

Comment [U3]: Check for typo

Comment [U4]: #4 anti-microbial

Comment [U5]: #5 anti-bacterial/microbial

Comment [U6]: # 6 protect neural disorder

Comment [U7]: # 7 anti diabetic

using the hormone sensitive lipase and pancreatic lipase potential which was imparted by Rosmarinic acid and gallic acid in Lavender oil which protects a person from cardiovascular diseases. Souri *et.al*, 2019 reported the Apoptotic activity using ELISA and TTC staining by targeting oxidative stress.

Comment [U8]: #8 cardiovascular

CONCLUSION

Lavender oil is a known element which has found place in cosmetic, perfumery and culinary industry due to bioactive potential and characteristic aroma of the flower and leaves. It is an important plant from time immemorial when it was used in bath by romans in ancient time, so we can trace back its importance as a plant. Lavender is being researched at a pretty infant stage because of localised occurrence and it is only reviewed for bioactive and pharmacological basis that too on a small pace and much more research is required in this part so as to flourish its literature about other such potentials and secrets that are to be unravelled in the coming time. The present study reviewed the aromatic profiling and bioactive substances that impart the plant certain useful properties that make it a mainstay herb in the gardens and adds to its economic production as a plant. So, there is a lot of scope of this plant to be researched and reviewed as with time and better understanding, we can explore the plant on a large scale based on new properties and can make it a multifarious drug.

REFERENCES

1. Tamokou, J. D. D., Mbaveng, A. T., & Kuete, V. (2017). Antimicrobial activities of African medicinal spices and vegetables. In *Medicinal spices and vegetables from Africa* (pp. 207-237). Academic Press
2. Verma, R. S., Rahman, L. U., Chanotiya, C. S., Verma, R. K., Chauhan, A., Yadav, A., ... & Yadav, A. K. (2010). Essential oil composition of *Lavandula angustifolia* Mill. cultivated in the mid hills of Uttarakhand, India. *Journal of the serbian chemical society*, 75(3), 343-348.
3. Singh, P., Andola, H., Rawat, M. S. M., & Jangwan, J. S. (2015). GC-MS Analysis of Essential Oil from *Lavandula angustifolia* Cultivated in Garhwal Himalaya. *The Natural Products Journal*, 5(4), 268-272.

4. Hajhashemi, V., Ghannadi, A., & Sharif, B. (2003). Anti-inflammatory and analgesic properties of the leaf extracts and essential oil of *Lavandula angustifolia* Mill. *Journal of ethnopharmacology*, 89(1), 67-71.
5. Cardia, G. F. E., Silva-Filho, S. E., Silva, E. L., Uchida, N. S., Cavalcante, H. A. O., Cassarotti, L. L., ... & Cuman, R. K. N. (2018). Effect of lavender (*Lavandula angustifolia*) essential oil on acute inflammatory response. *Evidence-Based Complementary and Alternative Medicine*, 2018.
6. Li, J., Wang, Y., Dong, Y., Zhang, W., Wang, D., Bai, H., ... & Shi, L. (2021). The chromosome-based lavender genome provides new insights into Lamiaceae evolution and terpenoid biosynthesis. *Horticulture research*, 8(1), 1-14.
7. Zagorcheva, T., Stanev, S., Rusanov, K., & Atanassov, I. (2020). SRAP markers for genetic diversity assessment of lavender (*Lavandula angustifolia* mill.) varieties and breeding lines. *Biotechnology & Biotechnological Equipment*, 34(1), 303-308.
8. Apetrei, Cristina & Corciova, Andreia & Şpac, Adrian & Ciobanu, Constantin & Ivanescu, B.. (2014). evaluation of bioactive compounds from commercial lavender products and comparative histo-anatomical study. *Analele Ştiinţifice ale Universităţii „Al. I. Cuza” Iaşi, s. II a. Biologie vegetală*. 60. 11-19
9. Ghelardini, C., Galeotti, N., Salvatore, G., & Mazzanti, G. (1999). Local anaesthetic activity of the essential oil of *Lavandula angustifolia*. *Planta medica*, 65(8), 700-703.
10. Guo, X., & Wang, P. (2020). Aroma Characteristics of Lavender Extract and Essential Oil from *Lavandula angustifolia* Mill. *Molecules*, 25(23), 5541
11. Duda, S. C., Mărghiţaş, L. A., Dezmirean, D., Duda, M., Mărgăoan, R., & Bobiş, O. (2015). Changes in major bioactive compounds with antioxidant activity of *Agastache foeniculum*, *Lavandula angustifolia*, *Melissa officinalis* and *Nepeta cataria*: Effect of harvest time and plant species. *Industrial Crops and Products*, 77, 499-507.
12. Kontogiorgis, C., Deligiannidou, G. E., Hadjipavlou, L. D., Lazari, D., Constantinides, T., Papadopoulos, A., & Shift, D. (2018). Antioxidant Contribution Of Lavender (*Lavandula angustifolia*). *Journal of Food Science and Technology*, 3(4), 360-377.
13. Tyśkiewicz, K., Konkol, M., & Rój, E. (2019). Supercritical Carbon Dioxide (scCO₂) Extraction of Phenolic Compounds from Lavender (*Lavandula angustifolia*) Flowers: A Box-Behnken Experimental Optimization. *Molecules*, 24(18), 3354.

14. Gezici, S. (2018). Promising anticancer activity of lavender (*Lavandula angustifolia* Mill.) essential oil through induction of both apoptosis and necrosis. *Annals of Phytomedicine*, 7(2), 38-45
15. Predoi, D., Groza, A., Iconaru, S. L., Predoi, G., Barbuceanu, F., Guegan, R., ... & Cimpeanu, C. (2018). Properties of basil and lavender essential oils adsorbed on the surface of hydroxyapatite. *Materials*, 11(5), 652
16. Zenão, S., Aires, A., Dias, C., Saavedra, M. J., & Fernandes, C. (2017). Antibacterial potential of *Urtica dioica* and *Lavandula angustifolia* extracts against methicillin resistant *Staphylococcus aureus* isolated from diabetic foot ulcers. *Journal of herbal medicine*, 10, 53-58.
17. Gonçalves, S., & Romano, A. (2013). In vitro culture of lavenders (*Lavandula* spp.) and the production of secondary metabolites. *Biotechnology advances*, 31(2), 166-174
18. Ayaz, M., Sadiq, A., Junaid, M., Ullah, F., Subhan, F., & Ahmed, J. (2017). Neuroprotective and anti-aging potentials of essential oils from aromatic and medicinal plants. *Frontiers in aging neuroscience*, 9, 168.
19. Issa, A., Mohammad, M., Hudaib, M., Tawah, K., Rjai, T. A., Oran, S., & Bustanji, Y. (2011). A potential role of *Lavandula angustifolia* in the management of diabetic dyslipidemia. *Journal of Medicinal Plants Research*, 5(16), 3876-3882
20. Sourì, F., Rakhshan, K., Erfani, S., Azizi, Y., Maleki, S. N., & Aboutaleb, N. (2019). Natural lavender oil (*Lavandula angustifolia*) exerts cardioprotective effects against myocardial infarction by targeting inflammation and oxidative stress. *Inflammopharmacology*, 27(4), 799-807

UNDER PEER REVIEW

