

Original Research Article

Anti – Androgenic activity of *Cesalpenia bonducella* in
androgen induced Polycystic ovarian syndrome rats

Abstract

Aim: Our goal was to use the androgen and anabolic steroid (AAS) Testosterone Propionate (T.P) to create a novel animal model to study polycystic ovaries.

Materials and Methods: Thirty albino female wistar rats weighing 200 – 250 g were split into V groups with six rats in each group. I Group was treated as normal control, II Group was treated as the PCOS induced group, III Group was treated as a standard group, Clomiphene citrate (20 mg per kg body weight), intraperitoneally (I.P) along with a 4 mg T.P injection intramuscularly (I.M) in an oily solution, IV Group was treatment control group treated with a hydro-alcoholic extract of *Cesalpenia bonducella* (CBHAE) at a dose of 200 mg per kg body weight. V Group was given CBHAE at 400 mg per kg of body weight. Blood collected from animals examined for hormonal parameters and ovaries is subjected to histopathological studies.

Results: There was a fall in testosterone volume and an increase in the efflux of female hormone constituents, which improved the ovarian development and helped to regulate menses.

Conclusion: The extract had a significant impact on reducing Testosterone production

Comment [R1]: The spelling is not correct. Caesalpinia

Comment [R2]: androgen-induced (consider adding the hyphen)

Comment [R3]: Double-check the capitalization here

Formatted: Line spacing: Exactly 14 pt

Comment [R4]: The abstract is usually a "gateway" encouraging the reading of the complete work. In this sense, I suggest modifying the abstract, briefly describing the methodology, including more specific details about its results, and answering the working hypothesis (that is, answering if and how it reached the objective).

Comment [R5]: Your sentence may be unclear or hard to follow. Consider rephrasing. Something similar to this example: "To create a novel animal model to study polycystic ovaries, our goal was to use the androgen and anabolic steroid (AAS) testosterone propionate (TP)."

What is the difference between your study model and the model used by Ayyadurai et al.? "Ayyadurai, T., Thirupathi, S. K., Shanmugam, A., & Diana, R. K. B. (2021). Effect of aqueous seed extract of *Caesalpinia bonduc* (L.) Roxb., on hormonal assay and lipid profile in induced Polycystic Ovary Syndrome albino female rats. *International Journal of Botany Studies*, 6(3), 139–146. Retrieved from <https://www.botanyjournals.com/archives/2021/vol6/issue3/6-2-117>"

Comment [R6]: Hidden formatting symbols are present throughout the document (That's where 'Show All' becomes essential).

Comment [R7]: Consider capitalizing the word "Wistar". Because is a proper noun.

Comment [R8]: The word **normal** is often overused. Consider using a more specific synonym (e.g. standard) to improve the sharpness of your writing.

Comment [R9]: You have written the same word with and without a hyphen in your document. Both ways are acceptable, but it's best to be consistent.

Comment [R10]: Correct your spelling.

and had improved anti-androgenic effects. However, further intervention is suggested to investigate the therapeutic effect of *Cesalpenia bonducella* seed.

Comment [R12]: Correct your spelling.

Keywords- anti-androgens, *Cesalpenia bonducella*, follicles, PCOS, tubular injury.

Comment [R13]: Correct your spelling.

1. INTRODUCTION

Androgens in normal levels is beneficial, but in abnormal range (women), it leads to serious consequences i.e. forbid ovaries from releasing egg (ovulation), excessive hair growth, acne, Insulin resistance e.t.c eventually leads to PCOS.

Comment [R14]: are

Comment [R15]: The word **serious** is often overused. Consider using a more specific synonym (e.g. severe) to improve the sharpness of your writing.

Comment [R16]: ???

Luteinizing hormone (LH) is a pituitary hormone release in response to Luteinizing hormone-Releasing hormone. It regulates the length and duration of menstrual cycle, ovulation, implantation of

Comment [R17]: released

fertilized egg in uterus, secretion of estrogen and progesteron. Theca cells secrete testosterone in response to LH surge and inturn converts to estrogen by adjoining granulose cells. Ovulation is stimulated by excess LH, whereas a relative FSH deficiency affects follicular growth. LH imbalance:

Comment [R18]: the menstrual

FSH (3:1) induces ovary theca cells to proliferate, which results in increased steroidogenesis and eventually in hyperandrogenism in PCOS women. *Cesalpenia bonducella* is abundant in

Comment [R19]: a fertilized

Comment [R20]: an egg

Comment [R21]: the uterus

Comment [R22]: progesterone

Comment [R23]: secrete

Comment [R24]: in turn convert

Comment [R25]: steroidogenesis

furanoditerpenes beneficial in management of PCOS [1]. Seeds comprises of phytochemicals includes β - caesalpin, γ - caesalpin, caesalpin, citrullin and bonducellin, as well as fatty acid; steric acid, palmitic acid, and oleic acid; amino acids such as arginine and aspartic acid [2-3].

Comment [R26]: Your sentence may be unclear or hard to follow. Consider rephrasing. Something similar to this example: "LH imbalance: FSH (3:1) induces ovary theca cells to proliferate, resulting in increased **steroidogenesis** and eventually, hyperandrogenism in PCOS women."

Comment [R27]: Correct your spelling.

Comment [R28]: the management

Comment [R29]: comprised

Comment [R30]: including

Comment [R31]: citrulline

Comment [R32]: stearic

Comment [R33]: Correct your spelling.

Comment [R34]: a variety

Phytochemicals in *Cesalpenia bonducella* potent against variety of PCOS attributes [4]. -There are numerous documented medicinal properties like [5] anti-oxidant [6], adaptogenic [7], anti-inflammatory [8], anti-estrogen [9], and Immunomodulatory [10]. In order to appease the consequences anti-androgens came to light-~~in~~, in the present study by using the anti-androgenic property of *Cesalpenia bonducella* to appease PCOS in rodents.

2. MATERIAL AND METHODS

2.1 Collection of Plant Material

The marketed *Cesalpenia bonducella* was collected from an herbal production factory named ANNAIARAVINDHHERBALS located in 1, 2&3, Janaki Nagar, Maduravoyal, Chennai, and Tamil Nadu 600095.

Comment [R35]: Correct your spelling.

2.2 Plant Authentication

The roots were recognized by professor P Jayaraman, Ph.D. from the plant anatomy and research center located at Tambaram (west). The authentication ID is **PARC/2019/4111** and the certificate is enclosed below in the Appendix-I

2.3 Extraction Procedure

Cesalpenia bonducella fresh fruits were collected and authenticated. After the seeds were rubbed out, the skin was left to dry in the shade. The shade-dried seeds were mashed into a coarse powder and sieved through mesh no 60, were subjected to extraction.

Comment [R36]: Correct your spelling.

Comment [R37]: were

500mg of coarsely powdered dried Grains of *Cesalpenia bonducella* was placed in a maceration flask with 70% of 2-liter Ethanol at 70°C temperature for 72 hours with occasional shaking. Then it was filtered, and the macerate was evaporated to attain the yield. Finally, the percentage of yield was calculated.

Comment [R38]: Correct your spelling.

2.4 Experimental model

2.4.1 Grouping of animal's

Female albino rats (180-220gm) were purchased from VETERINARY AND ANIMAL SCIENCES UNIVERSITY, TAMIL NADU, Laboratory Animal Medicine, and Centre for Animal Health Studies, Chennai- 51 the study was taken from an experimental facility, they were kept in a poly glycol boxes in a temperature-controlled (25±20°C) environment with a 12 hours dark/light cycle with standard ad

libitum. All animals were acclimatised to the laboratory condition for a period of 15 days before the commencement of the study. The animals were divided into five groups, each of which had five animals.

Comment [R39]: acclimatized

2.4.2 PCOS Induction in the Animals

The treatment of PCOS is evaluated in this study using T.P-caused PCOS. [11] Thirty female virgin Wistar rats, weighing 180-220g of 10-12 weeks aged, having regular estrus cycles as determined by vaginal smear, were employed in the study. [12]. Excluding the normal control group, all the other groups were treated with 4mg TP intramuscular injection in oily solution per rat. [13, 14] each animal had subjected to vaginal smear examination daily. The absence of cyclicity was shown as criteria for induction of PCOS by prolonged cornification of vaginal smears.

Formatted: Font color: Red

Comment [R40]: The word **normal** is often overused. Consider using a more specific synonym (e.g. standard) to improve the sharpness of your writing.

Comment [R41]: You used abbreviations both with and without periods. Both styles are acceptable, but it's best to use one style throughout your document.

Formatted: Font color: Red

2.4.3 Vaginal smears

The stage of cyclicity was determined via microscopic examination of the main cell type in vaginal smears. Estrous cycle cyclicity was assessed with 800 to 1200 hrs, and the relative components of epithelium, leukocytes, and cornified cells collect in daily vaginal lavages was studied using light microscopy. Which are known to alter at different stages of the estrous cycle. In the control group or PCOS, rats (proestrus, estrus, diestrus1, and diestrus2,) lasts usually about 4 days. [15]

Comment [R42]: primary

Comment [R43]: collected

Comment [R44]: were

Comment [R45]: usually lasts

Comment [R46]: Your sentence may be unclear or hard to follow. Consider rephrasing. Something similar to this example: "The control group or PCOS rats (proestrus, estrus, diestrus1, and diestrus2) usually lasts about 4 days."

2.4.4 Vaginal Cytology

Every day vaginal smears were checked, while having lumbar support, the rats were confined at the upper ventral side and thorax. Cotton soaked in a drop of physiological saline solution was used to collect vaginal secretions. The cotton swab was inserted almost one or two inches into the female rat's vagina and rotated for two to three rotations. (Allowing the cotton swab end to collect an acceptable volume of cells), and the cotton tip was rolled along the glass slide and swab has been gently removed. The dried smear was preserved by soaking it in 70% alcohol. After that, the slides

Comment [R47]: along with

Comment [R48]: the swab was

were stained with methylene blue 0.5 % solution, ~~cleansed~~ with tap water and inspected under a light microscope with a ×10 objective lens (without the condenser lens).[16]

Comment [R49]: cleansed

2.4.5 Treatment Protocol

Except for I Group, which receives TP intramuscularly, after induction, all rats in Groups IV and V were given oral gavage for 15 days. Whereas group 3 rats were treated with Clomiphene citrate Standard drug for ~~a period of~~ 15 days. On 16th day, ~~a~~ Retro orbital puncture was performed to collect blood samples and serum was subjected to hormonal analysis (LH, FSH, estradiol, progesterone and testosterone). Animal in each group was selected and euthanized in ~~a~~ carbon dioxide incubator. Finally, ~~a~~ incision was made to collect the ovaries then weighed and subjected to histopathological examination.

Comment [R50]: Except for I Group, which receives TP intramuscularly, after induction, all rats in Groups IV and V were given oral gavage for 15 days after induction.

2.4.6 Serum Hormonal Assay

LH, FSH, Serum testosterone, progesterone and estradiol, were measured using ~~an~~ Enzyme-Linked immunoassay kit for quantitative estimation of respective hormones.^[16]

2.4.7 Histopathological examination

Bouin's solution was used to repair the removed ovaries and the ovaries were drained in a series of alcohol, and then cleansed in xylene before being immersed in heated paraffin wax at 60 degrees Celsius. They were dehydrated in a series of alcohol, and then cleaned in xylene by being embedded in paraffin wax that was melted at 60°C. Consecutive slices were placed on ~~a~~ slide with 3-aminopropyl ~~triethylsilane~~-coated sections were dried at 37°C for 24 hours. Before being mounted for histology, the slides were deparaffinized, hydrated, and stained with Mayer shematoxylin & eosin dyes. Scope picture 3.0 imaging instrument (ScopeTek DCM 200) was used to examine the ovaries at a magnification of 40X (Hangzhou Scope tekOpto-Electric Co Ltd, USB 2.0). The thickness and

Comment [R51]: They were ~~was~~ dehydrated in a series of alcohol, and then cleaned in xylene by being embedded in paraffin wax ~~that was~~ melted at 60°C.

Comment [R52]: triethylsilane

diameter of cystic follicles were measured. Cystic follicles have a thicker, fibrotic cortex with a visible surface theca and interior layer. [17]

2.4.8 Statistical analysis

The results are expressed as Mean \pm SEM. Data were evaluated using ONE WAY ANOVA followed by Newman – Keul's multiple range test. Probability values less than ($p < 0.01$) were considered significant.

3. RESULTS AND DISCUSSION

Table 1: Percentage yield (w/w) of Plant extract

Sr No	Extract	Color	Percentage yield (w/w)
1.	Hydroalcoholic extract of <i>Cesalpenia bonducella</i>	Brownish Black	3.70

Comment [R53]: Correct your spelling.

3.1 Hormonal parameter evaluation:

3.1.1 The effect of CBHAE on LH levels in rats with T.P induced PCOS rats

When compared to the normal control group (G1), T.P causes a considerable increase in LH levels & decrease in toxic control FSH (G2) levels ($P < 0.001$). The levels of LH/FSH differed considerably between control groups. When compared to toxic control group, extracted treated animal's i.e., and both doses of CBHAE 200mg/kg & 400mg/kg exhibited a low levels of LH/FSH ratio and a remarkable drop in LH ($P < 0.01$) & ascent in FSH levels (Tab: 2)

Comment [R54]: and a
Formatted: Highlight

Comment [R55]: the toxic

Comment [R56]: a low level or low levels

3.1.2 The effect of CBHAE in T.P induced PCOS rats

Estradiol levels decline markedly when T.P injection are given. Considerable rise in estradiol levels ($P < 0.01$) result from simultaneous treatment for 15 days of CBHAE extracts. In standard group animals, estradiol levels are also significantly increased (Tab: 2)

Comment [R57]: injections

Comment [R58]: A considerable

3.1.3 The Effect of CBHAE on progesterone in TP induced PCOS rats

The drug-induced group, which was given T.P, has considerably decreased in progesterone levels. CBHAE treatment at both dosages in accordance with T.P (200mg/kg & 400mg/kg) caused a substantial rise in progesterone ($P < 0,001$) levels to nearly normal levels. The control group found similar findings with CBHAE (Tab: 2).

3.1.4 The effect of CBHAE on ovarian morphology

There was more cystic follicle in the ovaries of the positive control group (T.P), but, but in 200mg/kg & and 400mg/kg CBHAE shows near to normal follicles. Atretic follicles were seen and present at 200mg/kg. There were several healthy follicles in the 400 mg/kg group (Tab: 3)

3.1.5 The Effect of CBHAE on follicular diameter & thickness

The diameter and thickness of the cysts in the PCOS treatment group have increased, but the standard group and extract-treated group has fallen. (Tab: 3)

3.1.6 The Effect of CBHAE on ovarian weight

The T.P. control group's ovarian weight fall significantly ($P < 0.01$) as compared to the other groups, but it was restored to normal levels in the treatment groups 200mg/kg & and 400mg/kg. (Tab: 3)

Comment [R59]: fell

Table 2: Effect of CBHAE on serum hormone in T.P induced PCOS

GROUP	FSH	LH	Estradiol	TSH	PRGSN
G1	8.27±0.2 5	6.146±0.33	54.12±2.31	0.28±0.04	14.15±0. 61
G2	2.26±0.2 35**a	11.36±0.75* *a	14.32±0.82**a	0.37±0.02**a	7.055±0. 712**a
G3	7.02±0.4 6	5.25±0.38	46.49±1.74	0.32±0.02	12.12±0. 29
G4	6.63±0.5 3**b	3.70±0.18** b	38.17±1.33**b	0.34±0.01**b	10.8±0.7 2*b
G5	6.13±0.5 6**b	4.57±0.21** b	41.18±0.92**b	0.32±0.01**b	11.60±0. 76*b

All data's were analyzed G1-Normal, G2-Toxic, G3-Standard, G4-Low dose (CBHAE), G5-High dose (CBHAE)

For each group of 6 animals, all values represented as means ± SEM.

**a- values are significantly differ from Normal control (G1) at P<0.001

**b- values are significantly differ from PCOS control (G2) at P<0.001

*b- values are significantly differ from PCOS control (G2) at P<0.01

Table 3: Effect of CBHAE on ovarian morphology of PCOS rats

Dose mg.kg ovarian feature	Normal	Std control	Toxic control	Low dose	High dose
Atretic follicle	0.00±0.0 0	1.12±0.0 5	4.43±0.3 07	3.06±0.17* *b	0.05±0.11* b

Cystic follicle	0.00±0.0 0	3.7±10.5 8	10.55±1. 21	0.00±0.00	0.00±0.00
Cystic follicle diameter	0.00±0.0 0	71.19±2. 35	87.73±2. 367	0.00±0.00	0.00±0.00
Cystic follicle thickness	0.00±0.0 0	34.63±2. 18	42.35±1. 48	0.00±0.00	0.00±0.00

G1-Normal, -G2-Toxic, G3-Standard, G4-Low_dose, G5-High dose

For each group of 6 animals, all values represented as means ± SEM.

**b- values are considerably differ from PCOS control (G2) at P<0.001


Comment [R60]: different

*b- values are considerably differ from PCOS control (G2) at P<0.01

Comment [R61]: different

3.2 Histopathological Examination

Comment [R62]: Histopathological



G1: Normal Control (10ml/kg Normal Saline) Fig.1, G2: Drug-Induced Group (T.P 4mg/kg) Fig.2, G3: Standard Control (Clomiphene Citrate 20mg/ml), G4: Treatment Control (LOW DOSE of CBHAE 200mg/kg), G5: Treatment Control (High Dose of CBHAE 400mg/kg)

G1 Antral follicles, Corpus luteum, Oocyte are surrounded by granulosa cells, and theca cells is visible in this section, G2 Many cystic degenerating follicle and atretic follicles with degenerated granulosa layer is observed in an ovarian section from PCOS rat, G3 –A normal follicle with clear antrum, and Oocyte in granulosa layer is visible in the ovarian section of PCOS rat, treated with clomiphene citrate, G4 The ovary of a PCOS rat treated with a low dose of CBHAE 200mg/kg reveals mild degenerative follicle and absence of cystic, atretic follicle, G5 The ovary of a PCOS rat treated with a low dose of CBHAE 400mg/kg reveals the existence of growing regenerating follicles and corpus luteum, as well as Oocytes within granulosa- and thicker theca.

Comment [R63]: Corpus

Comment [R64]: cells. Theca

Comment [R65]: clomiphene

Comment [R66]: corpus

Comment [R67]: within

4. Discussion

Although many models can be used to study PCOS, induction of PCOS by Testosterone Propionate can also be considered as one of the best models for studying PCOS. Hence, in terms of exhibiting the majority of reproductive and endocrine symptoms associated with PCOS, rodent PCOS models appear to closely parallel the human condition. This study investigated the effect CBHAE on the serum levels of LH, FSH, estradiol, testosterone & progesterone in TP induced PCOS. After 30 days of PCO induction, animals were analysed both harmonically & histologically, on 16th day after treatment with hydroalcoholic extract, animals were also analysed irrespective of their estrous cycle. In PCOS condition, normal gonadotropin-ovarian axis is disturbed results in hormonal imbalance reflected by the higher levels of LH, lower FSH levels and reversal of LH: FSH ratio. An elevated LH/FSH ratio and anovulation are typical findings in women with PCOS. [22,23] The extract treated

Comment [R68]: most

Comment [R69]: to parallel the human condition closely

Comment [R70]: analyzed

Comment [R71]: analyzed

Comment [R72]: estrous

groups shows better reduction in this LH/ FSH ratio indicate that extract could reverse PCOS condition. Oestrogen similar to other steroids becomes altered in PCOS. [24] Repetitive administration of CBHAE led to a significant rise in estradiol. Similarly, the reduction in the level of progesterone level in the PCOS-induced animals could be responsible for the persistent oestrus phase. [104] Elevation in the concentration of serum progesterone by CBHAE may be responsible for the reversal of reversing the luteal phase dysfunction and restoration of restoring normalcy of the estrous cycle.

Comment [R73]: The extract treated groups show shows better reduction in this LH/ FSH ratio, indicating indicate that extract could reverse PCOS condition.

Comment [R74]: oestrus phase

Our study showed that CBHAE induced an increase in serum estradiol implies, implying that plant causes marked improvement in endocrine function and the recovery of ovulatory functions in the rats. Hyperandrogenism (as a result of high testosterone levels) which is evident in human PCOS. [25, 26] was not present in this animal model of TP induced PCOS. [27] Therefore no effect of the extract on androgen levels was observed using this model of PCOS induction. Ovarian weight in PCOS induced rats was more than the normal rats which is in accordance with, according to earlier findings. [28-30] Treatment with HAECBE prevented further increase in ovarian weight & and returned to normal normally. The biochemical results are also supported by the histopathological observation of light microscopy. The histomorphometry of PCOS was a suitable measurement for describing the cystic status because differences were observed in the morphological characteristic and in the presence or absence of follicular cysts. [119] It is reported that the histopathological study of PCOS induced rats shows the formation of poly cysts in the ovary. [31, 32]. Ovaries exhibited increased follicle atresia and multiple cysts with thin granulosa cell layers and thickened theca cell layers. [33] After treatment with extract of *Cesalpenia bonducella*, PCOS condition was reversed, the number of cystic follicles reduces & and found numerous healthy follicles at different stage of development. This indicates that the treatment group shows a marked recovery of ovarian tissue, and the animals may probably be preparing for ovulation.

Comment [R75]: characters

Comment [R76]: granulosa

Comment [R77]: Correct your spelling.

Comment [R78]: reduced

Comment [R79]: a different stages

Comment [R80]: It may unclear who or what This refers to.

After treatment with CBHAE, the incidence of cystic follicles reduced and many healthy follicles at various stages of development were detected. This indicates that the therapy group has made substantial progress, and the ovarian tissue and animals are most likely preparing for ovulation. In addition, CBHAE therapy restores gonadotropin feedback inhibition (LH & FSH), resulting in a rise in estradiol and progesterone. The capability of the extract for reducing the histological, clinical, and biochemical aspects of the PCOS indicates all of a! these variables. The presence of flavanoids may explain the pharmacological effect of CBHAE extracts. Aside from the flavonoids discovered to have an antioxidant function in PCOS rats, the PCOS illness has been linked to a reduction in the quantity of antioxidant enzyme/molecule. Antioxidants aid in the protection of the human body from the negative effects of reactive oxygen species (ROS) (ROS). Plants with a strong antioxidant characteristic, particularly flavonoids, have shown themselves. The well-known usage of plants in the past in the management of gynaecological disorders may be responsible for these plant phytochemicals.

Comment [R81]: was reduced, and

Comment [R82]: It may unclear who or what This refers to.

Comment [R83]: extract's capability

Comment [R84]: flavonoids

Comment [R85]: reducing

Comment [R86]: protecting

Comment [R87]: gynecological

5. CONCLUSION

PCOS is a prevalent endocrine disorder of women, which can results in infertility. The prospective function of Herbal medicine in the treatment of PCOS is consistent with few adverse effects. Herbal medicine improves body immunity without changing hormonal level and also regulariseregularize the menstrual cycle. In conclusion, CBHAE reveals a considerable recovery of PCOS animals from, LH, FSH, estradiol & and progesterone as well as, and the irregular cycle and ovarian physiology. T.P injection enhances the ovary weight and reproductive system of normal rats, which returned to normal, after treating with CBHAE. Polycyst's were seen as a group exception of the normal standard

Comment [R88]: The conclusion is not a repetition of the results and discussion. On the contrary, it must be in the present and bring the main contributions of the work and future perspectives.

control group, which has no polycysts ~~at all~~. CBHAE was found to correct the hormonal imbalance caused by T.P in PCOS rats, indicating that it might used be used to treat infertility.

Comment [R89]: publicists

CONSENT

It is not applicable.

ETHICAL APPROVAL

The research has been reviewed and approved by the Institutional animal ethical committee and was carried out ~~in accordance with~~ according to CPCSEA guidelines (approval number **IAEC/219/2019**).

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. Pascoe, Keith O., Basil A. Burke, and Wilfred R. Chan. "Caesalpin F: a new furanoditerpene from *Caesalpinia bonducella*." *Journal of Natural Products* 49.5 (1986): 913-915. <https://doi.org/10.1021/np50047a025>
2. Prasad, GoliPenchala, et al. "Analytical study of Kuberaksha/KantakikaranjapatraChurna [*Caesalpinia bonduc* (L.) Roxb. leaf powder]." *Ayu* 31.2 (2010): 251.
3. Ramadurai, Sivasankari, and Usha Balasundaram. "Rhizomicrobiomics of *Caesalpinia bonducella*, a wonder plant for PCOS treatment." *Physiology and Molecular Biology of Plants* 26.12 (2020): 2453-2463.
4. Kandasamy, Veerapandiyan, and Usha Balasundaram. "*Caesalpinia bonduc* (L.)

Roxb. As a promising source of pharmacological compounds to treat Poly Cystic Ovary Syndrome (PCOS): A review." *Journal of Ethnopharmacology* (2021): 114375.

5. Shukla, Shruti, et al. "Antioxidant activity and total phenolic content of ethanolic extract of *Caesalpinia bonducella* seeds." *Food and chemical Toxicology* 47.8 (2009): 1848-1851.
6. Kannur, D. M., V. I. Hukkeri, and K. S. Akki. "Adaptogenic activity of *Caesalpinia bonducella* seed extracts in rats." *Journal of Ethnopharmacology* 108.3 (2006): 327-331.
7. Shukla, Shruti, et al. "Studies on anti-inflammatory, antipyretic and analgesic properties of *Caesalpinia bonducella* F. seed oil in experimental animal models." *Food and Chemical Toxicology* 48.1 (2010): 61-64.
8. Biswas, T. K., et al. "Oral hypoglycemic effect of *Caesalpinia bonducella*." *International journal of pharmacognosy* 35.4 (1997): 261-264.
9. Kannur, Dayanand M. "Antidiabetic and antihyperlipidemic activity of *Bonducella* (*Caesalpinia bonducella*) seeds." *Nuts and Seeds in Health and Disease Prevention*. Academic Press, 2011. 237-244.
10. Shukla, Shruti, et al. "In vivo immunomodulatory activities of the aqueous extract of bonduc nut *Caesalpinia bonducella* seeds." *Pharmaceutical biology* 48.2 (2010): 227-230.
11. Alaa El-Din, A. B., Vern, J.C., Michelle, M., Jenny, H.Z., and Richard, S. Antioxidant activities, sensory and anti-influenza activity of grape skin tea infusion. *Food Chemistry*, 2011, 129: 837–845.
12. Dehghani, F., Panjehshahin, M. R., and Vojdani, Z. The Effect of hydroalcoholic extract of caraway on thyroid gland structure and hormones in female rat. *Iranian Journal of Veterinary Research*, Shiraz University, 2010, 11(4): 337-341.
13. Sunanda, P., and Anand, K. studied the anti-thyroid action of leaf extract of *Annona squamosa* in albino rats. *Current Science*, 2003, 84(11):1402-1404.
14. Kar, A., Panda, S., and Bharti, S. The Relative efficacy of three medicinal plant extracts in the alteration of thyroid hormone concentrations in male mice. *Journal of Ethnopharmacology*, 2002, 81:281-285.
15. Panda, S., Jafri, M., Kar, A., and Meheta, B.K. The Thyroid inhibitory, antiperoxidative and hypoglycemic effects of stigmasterol isolated from *Butea monosperma*. *Fitoterapia*, 2009, 80:123-126.
16. Jean-Marc, S., Franck, B., and Michel, M. The Polymeric proanthocyanidins from grape skins. *Phytochemistry*, 1996, 43(2):509-512.
17. Sourgens, H., Winterhoff, H., Gumbinger, H.G., and Kemper, F.H. Antihormonal effects of plant extracts, TSH and prolactin-suppressing properties of *Lithospermum officinale* and other plants. *Planta Med.*, 1982, 45:78-86.
18. Kim, S.M. Effects of ginseng radix on the rat hypothyroidism induced by PTU (6-n-propyl-2-thiouracil). *Korean Journal of Herbology*, 2010, 25, (3):11–18.
19. Kavutcu, M.O., and Canbolat, ztu"rk, S. O", et al. Reduced enzymatic antioxidant defense mechanism in kidney tissues from gentamicin-treated guinea pigs: effects of vitamins E and C. *Nephron*, 1996, 72(2): 269–274.

20. Jamall, I. S., Smith, J. C. Effects of cadmium on glutathione peroxidase, superoxide dismutase, and lipid peroxidation in the rat heart: a possible mechanism of cadmium cardiotoxicity. *Toxicology and Applied Pharmacology*, 1985, 80(1): 33–42.
21. Sedlak, J., Lindsay, R. H. Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Analytical Biochemistry*, 1968;25(1):192–205.
22. Ivana N., Patricia J., Marijan S., and Ana, M. O. B. Ultrasound-extracted flavonoids from four varieties of Portuguese red grape skins determined by reverse-phase high-performance liquid chromatography with electrochemical detection. *Analytica Chimica Acta*, 2008, 630:107–115.
23. Mukesh, Y., Shalini, J., Aarti, B., and Ravinder, N., et al., Biological and Medicinal Properties of Grapes and Their Bioactive Constituents: An Update. *Journal Of Medicinal Food*, 2009, 12 (3): 473–484.
24. Panda, S., and Kar, A. Amelioration of L-thyroxine-induced hyperthyroidism by coumarin (1, 2-benzopyrone) in female rats. *Clinical and Experimental Pharmacology and Physiology*, 2007, 34(11):1217–1219.
25. Seong-Mo, K., Sang-Chan, K., In-Kwon, C., Woo-Hyun, C., and Sae K. Antioxidant and Protective Effects of *Bupleurum falcatum* on the L-Thyroxine-Induced Hyperthyroidism in Rats. *Evidence-Based Complementary and Alternative Medicine*, 2012, pp.1- 12.
26. Lei, Z., Shelly, H., Jianrong, L., Shi, S., Corene, C., Shao, J.Z., and Kequan, Z. Grape skin extract inhibits mammalian intestinal α -glucosidase activity and suppresses postprandial glycemic response in streptozocin-treated mice. *Food Chemistry*, 2011, 126:466–471.
27. Tahiliani, P., and Kar, A. The combined effects of *Trigonella* and *Allium* extracts in the regulation of hyperthyroidism in rats. *Phytomedicine*, 2003, 10(8): 665–668.
28. Subudhi, K.U.D., Paital, B., Bhanja S., and Chainy, G. B. N. Alleviation of enhanced oxidative stress and oxygen consumption of L-thyroxine-induced hyperthyroid rat liver mitochondria by vitamin E and curcumin. *Chemico-Biological Interactions*, 2008,
29. Kim, S.M. Effects of ginseng radix on the rat hypothyroidism induced by PTU (6-n-propyl-2-thiouracil). *Korean Journal of Herbology*, 2010, 25, (3):11–18.
30. Kavutcu, M.O., and Canbolat, ztu'rk, S. O., et al. Reduced enzymatic antioxidant defense mechanism in kidney tissues from gentamicin-treated guinea pigs: effects of vitamins E and C. *Nephron*, 1996, 72(2):269–274.
31. Jamall, I. S., Smith, J. C. Effects of cadmium on glutathione peroxidase, superoxide dismutase, and lipid peroxidation in the rat heart: a

possible mechanism of cadmium cardiotoxicity. *Toxicology and Applied Pharmacology*, 1985, 80(1): 33–42.

33. Sedlak, J., Lindsay, R. H. Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Analytical Biochemistry*, 1968; 25(1): 192–205.

34. Aebi, H. Catalase, in *Methods in Enzymatic Analysis*, H. U. Bergmeyer, Ed., Academic Press, New York, NY USA, 1974: 673–686.

35. Sun, L. Y., Oberley, W., Y. Li. A simple method for clinical assay of superoxide dismutase. *Clinical Chemistry*, 1988; 34(3): 497–500..

36. Balu, M.S.P., Haripriya, D., Panneerselvam, C. Rejuvenation of antioxidant system in central nervous system of aged rats by grape seed extract. *Neurosci. Lett.*, 2005; 383: 295-300.

UNDR PEER REVIEW