

## Original Research Article

### ANTICANCER EFFECT OF DIGERA MURICATA LEAF EXTRACT AGAINST PROSTATE CANCER CELL LINES

**Comment [F1]:** Follow the naming rules.

**Running title:** Anticancer effect of *Digera muricata* leaf extract against prostate cancer cell lines.

**Comment [F2]:** This one is correct.

#### ABSTRACT

**Background:** Prostate cancer has become a major health problem globally during the last few decades. It is the second most frequently diagnosed cancer in men worldwide and the fifth most common cancer overall. Chemotherapy or other cancer treatments including androgen depletion therapy, show high toxicity and cause serious side effects in cancer patients. Thus, there have been many studies conducted to find various natural products as potential anticancer drug candidates with low toxicity and fewer side effects for the treatment and prevention of prostate cancer. The *Digera muricata* of the genus *Digera* Forssk. and family Amaranthaceae Juss. is a wild edible plant. The presence of phenol, flavonoids, alkaloids, terpenes, sterols, tannins, glycosides and lignins is seen in *Digera muricata* leaf extract, which are reported to have cytotoxic activities.

**Formatted:** Font: Italic

**Purpose:** The aim of this study is to assess the cytotoxic effects of *Digera muricata* leaf extract on prostate cancer cell line.

**Materials and methods:** The cytotoxic potency of *Digera muricata* leaf extract was carried out by MTT assay against the prostate cancer cell line. PC-3 cells were treated with different concentrations of (25-150µg/ml) *Digera muricata* leaf extract for 24h. Furthermore, the morphological changes were analysed using phase contrast microscopy and nuclear morphological changes examined using DAPI staining under the fluorescence microscopy

**Results:** The MTT assay showed decreased cell vitality with increased concentration of *Digera muricata* leaf extract. The morphological study showed that the number of cells decreased after

treatment and the cells exhibited cell shrinkage and cytoplasmic membrane blebbing. The treated cells also showed condensed chromatin and nuclear fragmentation

**Conclusion:** Within the limits of this study it can be demonstrated that the leaf extract of *Digera muricata* were cytotoxic and induced apoptosis to the cancer cells at higher concentration and incubation period.

**KEYWORDS:** *Digera muricata*, prostate cancer, cell viability, apoptosis

## INTRODUCTION

Prostate cancer is a major health problem and is the second most frequently diagnosed cancer in men worldwide and the fifth most common cancer overall<sup>1</sup>. Prostate cancer primarily affects elderly males with more than three quarter of the cases affecting men above the age of 65. Prostate cancer has become a major health problem globally during the last few decades. There has been lower incidences of prostate cancer in India and non resident Indians when compared to the Western population. Within India, prostate cancer has a wide disparity in disease characteristics, incidence, and mortality, with the urban population being more increasingly affected<sup>2</sup>. Incidence of prostate cancer is increasing in various parts of India. Changes in the diagnostic criteria and modalities, awareness levels of the public, and changing lifestyles may be responsible for much of the observed change<sup>3</sup>.

The prostate is a gland located between the bladder and the external genital organ of the males. The prostate is present in front of the rectum. The urethra runs through the center of the prostate which aids in the excretion of urine. The function of the prostate is to secrete fluid that nourishes and protects sperm. Androgens are required for growth of the cancerous cells in the early stages but eventually moves on to an androgen-independent stage, and progresses despite androgen extirpation<sup>4</sup>. The pathological mechanism at a molecular level is poorly understood for androgen-independent cancer progression. Among prostate cancer cell lines, the PC-3 cell line is known to be corresponding to androgen-independent cancer cells<sup>5,6</sup>. Chemotherapy or other cancer treatments including androgen depletion therapy, show high toxicity and cause serious

**Comment [F3]:** When cited for the first time, the acronyms must appear with their meaning so that the reader understands what is being said right away.

side effects in cancer patients. Drug resistance and low anticancer efficacy also pose difficulties in the clinical treatment<sup>7,8</sup>. For these reasons, there have been many studies conducted to find various natural products as potential anticancer drug candidates with low toxicity and fewer side effects for the treatment and prevention of prostate cancer.

The *Digera muricata* of the genus *Digera* Forssk. and family Amaranthaceae Juss. is a wild edible plant commonly known as 'Cancali soppu'. It is native to Egypt and eastern Kenya but can be commonly seen distributed throughout India. In Ayurveda the herb is considered to be a cooling astringent to the bowels and is also used as a laxative. The flowers and seeds are used to treat urinary discharges<sup>9</sup>. while the boiled root infusion is given to the mother after parturition for lactation purposes<sup>10</sup>. The presence of phenol, flavonoids, alkaloids, terpenes, sterols, tannins, glycosides and lignins in *Digera muricata* leaf extract was observed on conducting phytochemical tests<sup>11</sup>.

Tetrachloride-induced nephrotoxicity can cause the generation of reactive oxygen species which are in accumulation of amyloids, lipid peroxidation and thus can lead to kidney injuries<sup>12-14</sup>. *Digera muricata* is seen to have antioxidant properties which acts as a defence mechanism against toxicity induced by carbon tetrachloride and thus may have a role in the therapeutic treatment of free radical mediated diseases<sup>15</sup>. Cytotoxicity is the ability of a material to be toxic to a cell and induce apoptosis to that cell<sup>16,17</sup>. The cytotoxicity of extract of the plants may be due to the presence of flavonoids having mono to poly phenolic groups in the structure. The flavonoids have phenolic groups which are reported to have cytotoxic activities. Previous studies have been done to assess the association between flavonoids and cancer risk and it has been proven that there is a decrease in cancer risk with consumption of vegetables and fruits rich in flavonoids. Our team has extensive knowledge and research experience that has translated into high quality publications<sup>18-38</sup>. Thus in this study, we are assessing the cytotoxic effects of *Digera muricata* leaf extract on prostate cancer cells.

**Comment [F4]:** These studies were with *Digera muricata*?

**Comment [F5]:** Needless to write this here.

## MATERIAL AND METHODS

Reagents:

DMEM (Dulbecco's Modified Eagle Medium), Phosphate Buffered Saline (PBS), Trypsin-EDTA, Fetal bovine serum (FBS), were purchased from Gibco, Canada. Dimethyl sulfoxide (DMSO), [3-(4,5-dimethylthiazol-2-yl) 2,5-diphenyl tetrazolium bromide (MTT), DAPI, were purchased from Sigma Chemical Pvt Ltd, USA. All other chemicals used were extra pure of molecular grade and were purchased from SRL, India.

[Put the references that were reproduced in the methodology.](#)

Cell line maintenance:

PC-3 Prostate cancer cell lines were obtained from the National Center For Cell Science (NCCT), Pune. The cells were grown in T25 culture flasks containing DMEM supplemented with 10% FBS and 1% antibiotics. Cells were maintained at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub>. Upon reaching confluency, the cells were trypsinized and passaged.

Preparation of the Herbal Extract:

*Digera muricata* Leaf powder obtained from The Indian Medical Practitioners' Cooperative Pharmacy & Stores (IMPCOPS) (Chennai, India) was used for the present study. About 50g of *Digera muricata* powder was soaked in 500 mL of 95% ethanol and kept at room temperature for 3 days in a static condition. Then the solution was filtered with crude filter paper followed by whatman paper. Fine filtrate was subjected to rota evaporation after that 3g of the material was obtained. The total ethanol extract was concentrated in a vacuum and immediately stored at 4°C.

[Put the references that were reproduced in the methodology.](#)

Cell viability (MTT) assay:

The cell viability of *Digera muricata* extract treated A549 cells was assessed by MTT assay. The assay is based on the reduction of soluble yellow tetrazolium salt to insoluble purple formazan crystals by metabolically active cells. A549 cells were plated in 48 well plates at a concentration of 2x10<sup>4</sup> cells/well 24 hours after plating, cells were washed twice with 500µl of serum-free medium and starved by incubating the cells in serum-free medium for 3 hours at 37°C. After starvation, cells were treated with *Digera muricata* leaf extract in different concentrations (25 - 150µg) for 24 hours. At the end of treatment, the medium from control and *Digera muricata* treated cells were discarded and 200µl of MTT containing DMEM (0.5 mg/ml) was added to each well. The cells were then incubated for 4h at 37°C in the CO<sub>2</sub> incubator.

The MTT containing medium was then discarded and the cells were washed with 1x PBS. The crystals were then dissolved by adding 200µl of solubilization solution and this was mixed properly by pipetting up and down. Then the formazan crystals formed were dissolved in dimethyl sulfoxide (200µl) and incubated in dark for an hour. Then the intensity of the color developed was assayed using a Micro ELISA plate reader at 570 nm. The number of viable cells was expressed as the percentage of control cells cultured in serum-free medium. Cell viability in the control medium without any treatment was represented as 100%. The cell viability is calculated using the formula: % cell viability = [A570 nm of treated cells/A570 nm of control cells]×100.

[Put the references that were reproduced in the methodology.](#)

Morphology study:

Based on MTT assay we selected the optimal doses (IC-50: 50µg/ml) for further studies. Analysis of cell morphology changes by a phase contrast microscope.  $3 \times 10^4$  cells were seeded in 6 well plates and treated with *Digera muricata* (concentration for MCF-7 cells) for 24h. At the end of the incubation period, the medium was removed and cells were washed once with a phosphate buffer saline (PBS pH 7.4). The plates were observed under a phase contrast microscope.

[Put the references that were reproduced in the methodology.](#)

Determination of nuclear morphological changes of cells (DAPI staining):

For the nuclear morphological analysis, the monolayer of cells was washed with PBS and fixed with 3% paraformaldehyde for 10 min at room temperature. The fixed cells were permeabilized with 0.2% Triton X-100 in PBS for 10 min at room temperature and incubated with 0.5µg/ml of DAPI for 5 min. The apoptotic nuclei (intensely stained, fragmented nuclei, and condensed chromatin) were viewed under a fluorescent microscope.

[Put the references that were reproduced in the methodology.](#)

Statistical analysis:

Statistical analyses were performed using one-way ANOVA followed by Student–Newman–Keuls (SNK) tests for comparison between treatment values and control values. Data were

expressed as mean  $\pm$  SEM. The level of statistical significance was set at  $p < 0.05$ . [Put the references that were reproduced in the methodology.](#)

## RESULTS

In the current study, we analysed the effect of *Digera muricata* leaf extract on prostate cancer cell lines. [Figure 1] shows the MTT assay which determines the cytotoxic effect of the leaf extract on the cancer cells. Here, the cancer cell activity was inhibited in a dose dependent manner by the leaf extract of *Digera muricata*. The Cells were treated with increasing concentrations of the leaf extract (0, 25, 50, 75, 100, 125 and 150 $\mu$ g) for 24hrs. The 50% of inhibition observed in concentration of 50 $\mu$ g/ml. Maximum cancer cell viability is seen with the control (100%) while the minimum viability is seen with 150 $\mu$ g/ml (10%). [Figure 2] depicts the morphological changes in the prostate cancer cell lines after treatment with *Digera muricata* at 50  $\mu$ g/mL. Here, the cell number decreased after treatment and the cells exhibited cell shrinkage and cytoplasmic membrane blebbing. [Figure 3] depicts the nuclear changes in the treated cancer cells. The cells treated with the leaf extract showed condensed chromatin and nuclear fragmentation, characteristic features of apoptosis.





Figure 1: The cytotoxic effects of *Digera muricata* leaf extract on PC-3 prostate cancer cell line was determined by MTT assay. The Cells were treated with different concentrations (0, 25, 50, 75, 100, 125 and 150 $\mu$ g) for 24hrs. The 50% of inhibition observed in concentration of 50 $\mu$ g/ml, (*p* value: 0.0033) which has been taken as IC50 value and fixed for further experiments. \* represents statistical significance between control versus treatment groups at *p* < 0.05 using Student's-Newman-Keuls test

20X



Induction of apoptosis in *Digera muricata* leaf extract treated prostate cancer cell line

PC-3


Figure 2: Represents the morphological changes in prostate cancer cell line with and without treatment of *Digera muricata* at 50  $\mu$ g/mL for 24hrs by phase contrast microscope at 20x magnification. Here, the number of cells decreased after treatment and the cells exhibited cell shrinkage and cytoplasmic membrane blebbing.

[Follow the naming rules for the scientific name of the species in this image, in addition to placing the source.](#)

20X

Control

*Digeria muricata* leaf extract (50µg/ml)



*Figure 3: This figure represents the DAPI staining of PC-3 prostate cancer cell line upon, without and with treatment of Lippia nodiflora leaf extract at 50 µg/ml for 24 hrs. The nuclei were stained with DAPI staining and observed under a fluorescence microscope. The treated cells clearly showed condensed chromatin and nuclear fragmentation.*

Follow the naming rules for the scientific name of the species in this image, in addition to placing the source.

**Comment [F6]:** And the images of the other concentrations that also showed statistical difference? It is important for the reader to notice this difference in the images as well.

## DISCUSSION

Treatment for cancer including chemotherapy has been proven to have many side effects. Many new chemotherapeutic agents have been developed and are undergoing clinical trials. However, they have a narrow therapeutic spectrum considering their toxicity. Thus, many researches have been done to develop anticancer compounds derived from natural sources<sup>39</sup>. Certain plants contain natural compounds called phytochemicals which have anticancer effects and can be used to prevent cancer. These phytochemicals include phenol, flavonoids, alkaloids, terpenes, sterols, tannins, glycosides and lignins<sup>40</sup>.

In our study, based on the MTT assay, the cell viability is seen to decrease with the increased viability of the leaf extract. Maximum cancer cell viability is seen with the control (100%) while the minimum viability is seen with 150µg/ml (10%). IC-50 dose was observed at 50 µg/ml [Figure 1]. The morphology tests showed that the number of cells decreased after treatment and

the cells exhibited cell shrinkage and cytoplasmic membrane blebbing. [Figure 2]. This is a feature of cell apoptosis. The reason for cytoplasmic blebbing is not quite clear. A few hypotheses suggest that the cytoplasmic blebs can represent the structural and functional communication of the cell with the neighbouring cells to initiate phagocytosis or it also may be a way to deplete ATP in order to mix compartments to aid in cellular packaging or it may be a prerequisite for the formation of apoptosis bodies<sup>41</sup>. Also, The nuclei were stained with DAPI staining and observed under a fluorescence microscope. The treated cells clearly showed condensed chromatin and nuclear fragmentation which are features of apoptosis[Figure 3].

In our study the cell viability decreased with increased concentration of the leaf extract. This is in concordance with previous researches where the extract of *Digera muricata* showed inhibition of cancer cells at all concentrations in a dose dependent manner with a significant number of decrease in percentage proliferation against lung cancer cell lines<sup>40,42</sup>. A study by Shazia et al., on cervical and lung cancer cell lines provide similar results. It is suggested that the methanolic and aqueous fraction of the *Digera muricata* leaf extract played an important role in its anticancer effect. Here the anticancer activity was at its peak in the extract containing methanol which indicated that most of the active components were extracted using methanol<sup>40,42,43</sup>. The cytotoxic changes that were observed were cell aggregation, cell rounding and cell death<sup>44,45</sup>.

In the above study, the presence of flavonoids in the extract seems to have an important role in the anticancer effect of *Digera muricata*. In a similar study, Ashokkumar et al., stated that the activity of inhibition of the cancerous cells can be because of the interaction of the metabolic nature of the cancer cells with the nature of the compounds found in each crude extract or may be due to the effectiveness of some enzymes which act as antioxidants mainly in cancer cells<sup>46</sup>. While not many studies have been done regarding the anticancer effects of *Digera muricata*, there has been studies highlighting the antioxidant, antiinflammatory and antibacterial of the plant<sup>47-50</sup>. The limitation is that the study does not involve any in vivo study, So its effect is not assessed. This paves way for various future studies such as to view the drug action in in vivo studies and also to know about the side effects of the extract.

## CONCLUSION

Within the limits of this study it can be demonstrated that the leaf extract of *Digera muricata* were cytotoxic and induced apoptosis to the prostate cancer cells at 50 µg/ml concentration and 24h incubation period<sup>51 52 53 54 55 56 57 5859 60 61 62 63 64 65</sup>. However more research is needed to understand the mechanisms of cytotoxicity effect of *Digera muricata* extract.

**Comment [F7]:** Why is only this concentration being highlighted, if the graph shows that the others, with the exception of the smallest, were significant? These are points that confuse the reader. This information must be made explicit in the text.

**Comment [F8]:** ?????????? not necessary

**NOTE:**

The study highlights the efficacy of "ayurveda which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

**COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

**REFERENCES**

1. International Agency for Research on Cancer. *IARC Biennial Report 2010-2011*. 2012.
2. Bussemakers MJ, van Bokhoven A, Verhaegh GW, et al. DD3: a new prostate-specific gene, highly overexpressed in prostate cancer. *Cancer Res* 1999; 59: 5975–5979.

3. Hariharan K, Padmanabha V. Demography and disease characteristics of prostate cancer in India. *Indian J Urol* 2016; 32: 103–108.
4. Chen L. *ErbB3 in the Development of Androgen-independent Prostate Cancer*. 2010.
5. Weissmesser E. Growth hormone (GH) receptors in prostate cancer: gene expression in human tissues and cell lines and characterization, GH signaling and androgen receptor regulation in LNCaP cells. *Molecular and Cellular Endocrinology*. Epub ahead of print 2004. DOI: 10.1016/s0303-7207(04)00101-7.
6. Boo H-J, Hong J-Y, Kim S-C, et al. The anticancer effect of fucoidan in PC-3 prostate cancer cells. *Mar Drugs* 2013; 11: 2982–2999.
7. Blomberg K, Wengström Y, Sundberg K, et al. Symptoms and self-care strategies during and six months after radiotherapy for prostate cancer – Scoping the perspectives of patients, professionals and literature. *European Journal of Oncology Nursing* 2016; 21: 139–145.
8. Mandalà M, Romano E. *Mechanisms of Drug Resistance in Cancer Therapy*. Springer, 2019.
9. Parrotta JA. *Healing Plants of Peninsular India*. C A B International, 2001.
10. Gahukar RT. Potential of Minor Food Crops and Wild Plants for Nutritional Security in the Developing World. *Journal of Agricultural & Food Information* 2014; 15: 342–352.
11. Mathad P, Mety SS. Phytochemical and Antimicrobial Activity of *Digera Muricata*(L.) Mart. *E-Journal of Chemistry* 2010; 7: 275–280.
12. Khan MR, Ahmed D. Protective effects of *Digera muricata* (L.) Mart. on testis against oxidative stress of carbon tetrachloride in rat. *Food and Chemical Toxicology* 2009; 47: 1393–1399.
13. M S, Sohaib M, Ezhilarasan D. Carbamazepine, a Histone Deacetylase Inhibitor Induces Apoptosis in Human Colon Adenocarcinoma Cell Line HT-29. *Journal of Gastrointestinal Cancer* 2020; 51: 564–570.
14. Singh TR, Rohit Singh T, Ezhilarasan D. Ethanolic Extract of *Lagerstroemia Speciosa*(L.) Pers., Induces Apoptosis and Cell Cycle Arrest in HepG2 Cells. *Nutrition and Cancer* 2020; 72: 146–156.
15. Mv HMBH. Aqueous Extract of *Prosopis strombulifera* (LAM) BENTH Induces Cytotoxic Effects against Tumor Cell Lines without Systemic Alterations in BALB/c Mice. *Journal of Clinical Toxicology*; 04. Epub ahead of print 2014. DOI: 10.4172/2161-0495.1000222.
16. Shathviha PC, Ezhilarasan D, Rajeshkumar S, et al.  $\beta$ -sitosterol Mediated Silver Nanoparticles Induce Cytotoxicity in Human Colon Cancer HT-29 Cells. *Avicenna J Med Biotechnol* 2021; 13: 42–46.

17. Solai Prakash AK, Devaraj E. Cytotoxic potentials of *S. cumini* methanolic seed kernel extract in human hepatoma HepG2 cells. *Environ Toxicol* 2019; 34: 1313–1319.
18. Rajeshkumar S, Kumar SV, Ramaiah A, et al. Biosynthesis of zinc oxide nanoparticles using *Mangifera indica* leaves and evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells. *Enzyme Microb Technol* 2018; 117: 91–95.
19. Nandhini NT, Rajeshkumar S, Mythili S. The possible mechanism of eco-friendly synthesized nanoparticles on hazardous dyes degradation. *Biocatalysis and Agricultural Biotechnology* 2019; 19: 101138.
20. Vairavel M, Devaraj E, Shanmugam R. An eco-friendly synthesis of *Enterococcus* sp.-mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells. *Environ Sci Pollut Res Int* 2020; 27: 8166–8175.
21. Gomathi M, Prakasam A, Rajkumar PV, et al. Green synthesis of silver nanoparticles using *Gymnema sylvestre* leaf extract and evaluation of its antibacterial activity. *South African Journal of Chemical Engineering* 2020; 32: 1–4.
22. Rajasekaran S, Damodharan D, Gopal K, et al. Collective influence of 1-decanol addition, injection pressure and EGR on diesel engine characteristics fueled with diesel/LDPE oil blends. *Fuel* 2020; 277: 118166.
23. Santhoshkumar J, Sowmya B, Venkat Kumar S, et al. Toxicology evaluation and antidermatophytic activity of silver nanoparticles synthesized using leaf extract of *Passiflora caerulea*. *South African Journal of Chemical Engineering* 2019; 29: 17–23.
24. Raj R K, D E, S R.  $\beta$ -Sitosterol-assisted silver nanoparticles activates Nrf2 and triggers mitochondrial apoptosis via oxidative stress in human hepatocellular cancer cell line. *J Biomed Mater Res A* 2020; 108: 1899–1908.
25. Saravanan M, Arokiyaraj S, Lakshmi T, et al. Synthesis of silver nanoparticles from *Phenochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria. *Microb Pathog* 2018; 117: 68–72.
26. Gheena S, Ezhilarasan D. Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells. *Human & Experimental Toxicology* 2019; 38: 694–702.
27. Ezhilarasan D, Sokal E, Najimi M. Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets. *Hepatobiliary Pancreat Dis Int* 2018; 17: 192–197.
28. Ezhilarasan D. Oxidative stress is bane in chronic liver diseases: Clinical and experimental perspective. *Arab J Gastroenterol* 2018; 19: 56–64.
29. Gomathi AC, Xavier Rajarathinam SR, Mohammed Sadiq A, et al. Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of *Tamarindus indica* on MCF-7 human breast cancer cell line. *Journal of Drug Delivery Science and Technology* 2020; 55: 101376.

30. Dua K, Wadhwa R, Singhvi G, et al. The potential of siRNA based drug delivery in respiratory disorders: Recent advances and progress. *Drug Dev Res* 2019; 80: 714–730.
31. Ramesh A, Varghese S, Jayakumar ND, et al. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *Journal of Periodontology* 2018; 89: 1241–1248.
32. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. *Arch Oral Biol* 2021; 122: 105030.
33. Joseph B, Prasanth CS. Is photodynamic therapy a viable antiviral weapon against COVID-19 in dentistry? *Oral surgery, oral medicine, oral pathology and oral radiology* 2021; 132: 118–119.
34. Ezhilarasan D, Apoorva VS, Vardhan NA. Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. *Journal of Oral Pathology & Medicine*. Epub ahead of print 2018. DOI: 10.1111/jop.12806.
35. Duraisamy R, Krishnan CS, Ramasubramanian H, et al. Compatibility of Nonoriginal Abutments With Implants. *Implant Dentistry* 2019; 28: 289–295.
36. Gnanavel V, Roopan SM, Rajeshkumar S. Aquaculture: An overview of chemical ecology of seaweeds (food species) in natural products. *Aquaculture* 2019; 507: 1–6.
37. Markov A, Thangavelu L, Aravindhana S, et al. Mesenchymal stem/stromal cells as a valuable source for the treatment of immune-mediated disorders. *Stem Cell Res Ther* 2021; 12: 192.
38. Veerasamy R, Roy A, Karunakaran R, et al. Structure-Activity Relationship Analysis of Benzimidazoles as Emerging Anti-Inflammatory Agents: An Overview. *Pharmaceuticals* ; 14. Epub ahead of print 11 July 2021. DOI: 10.3390/ph14070663.
39. Khan T, Gurav P. PhytoNanotechnology: Enhancing Delivery of Plant Based Anti-cancer Drugs. *Frontiers in Pharmacology*; 8. Epub ahead of print 2018. DOI: 10.3389/fphar.2017.01002.
40. McKeage MJ. Clinical Toxicology of Platinum- Based Cancer Chemotherapeutic Agents. *Platinum-Based Drugs in Cancer Therapy*; 251–275.
41. Mills JC, Stone NL, Pittman RN. Extranuclear Apoptosis. *Journal of Cell Biology* 1999; 146: 703–708.
42. Khare CP. *Digera muricata* (Linn.) Mart. *Indian Medicinal Plants* 2007; 1–1.
43. Usmani S, Hussain A, A. FAH, et al. Anti-proliferative Activity of Crude Extract and Fractions Obtained from *Digera muricata* on HeLa Cell Lines of Human Cervix and A549 Cell Lines of Human Lung. *Pharmacognosy Journal* 2014; 6: 32–38.

44. Thakur RS, Devaraj E. Lagerstroemia speciosa (L.) Pers. triggers oxidative stress mediated apoptosis via intrinsic mitochondrial pathway in HepG2 cells. *Environ Toxicol* 2020; 35: 1225–1233.
45. Iyengar RM, Devaraj E. Silibinin Triggers the Mitochondrial Pathway of Apoptosis in Human Oral Squamous Carcinoma Cells. *Asian Pacific Journal of Cancer Prevention* 2020; 21: 1877–1882.
46. Prabhu PN, Ashokkumar P, Sudhandiran G. Antioxidative and antiproliferative effects of astaxanthin during the initiation stages of 1,2-dimethyl hydrazine-induced experimental colon carcinogenesis. *Fundamental & Clinical Pharmacology* 2009; 23: 225–234.
47. Khan MR. Protective effects of *Digera muricata* (L.) Mart. against carbon tetrachloride induced oxidative stress in thyroid of rat. *AFRICAN JOURNAL OF BIOTECHNOLOGY*; 10. Epub ahead of print 2011. DOI: 10.5897/ajb11.691.
48. Muhammad RK, Muhammad A, Naima S, et al. Protective potential of methanol extract of *Digera muricata* on acrylamide induced hepatotoxicity in rats. *African Journal of Biotechnology* 2011; 10: 8456–8464.
49. Shah R, Shah SA, Shah S, et al. Green Synthesis and Antibacterial Activity of Gold Nanoparticles of *Digera muricata*. *Indian Journal of Pharmaceutical Sciences*; 82. Epub ahead of print 2020. DOI: 10.36468/pharmaceutical-sciences.659.
50. al. AM et AM et, al. AM et, al. AM et, et al. Evaluation of Antibacterial Efficacy of *Digera Muricata* Plant using Microbroth Dilution and Autobiography Assays. *International Journal of Botany and Research* 2018; 8: 9–16.
51. Ezhilarasan D, Apoorva VS, Ashok VN. Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. *J Oral Pathol Med*; 48. Epub ahead of print February 2019. DOI: 10.1111/jop.12806.
52. Danda AK, Krishna TM, Narayanan V, et al. Influence of primary and secondary closure of surgical wound after impacted mandibular third molar removal on postoperative pain and swelling--a comparative and split mouth study. *J Oral Maxillofac Surg*; 68. Epub ahead of print February 2010. DOI: 10.1016/j.joms.2009.04.060.
53. Ramadurai N, Gurunathan D, Samuel AV, et al. Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial. *Clin Oral Investig*; 23. Epub ahead of print September 2019. DOI: 10.1007/s00784-018-2775-5.
54. Sathivel A, Raghavendran HR, Srinivasan P, et al. Anti-peroxidative and anti-hyperlipidemic nature of *Ulva lactuca* crude polysaccharide on D-galactosamine induced hepatitis in rats. *Food Chem Toxicol*; 46. Epub ahead of print October 2008. DOI: 10.1016/j.fct.2008.07.016.
55. Panda S, Doraiswamy J, Malaiappan S, et al. Additive effect of autologous platelet concentrates in treatment of intrabony defects: a systematic review and meta-analysis. *J*

- Investig Clin Dent*; 7. Epub ahead of print February 2016. DOI: 10.1111/jicd.12117.
56. Neelakantan P, Varughese AA, Sharma S, et al. Continuous chelation irrigation improves the adhesion of epoxy resin-based root canal sealer to root dentine. *Int Endod J*; 45. Epub ahead of print December 2012. DOI: 10.1111/j.1365-2591.2012.02073.x.
  57. Govindaraju L, Neelakantan P, Gutmann JL. Effect of root canal irrigating solutions on the compressive strength of tricalcium silicate cements. *Clin Oral Investig*; 21. Epub ahead of print March 2017. DOI: 10.1007/s00784-016-1922-0.
  58. Sekhar CH, Narayanan V, Baig MF. Role of antimicrobials in third molar surgery: prospective, double blind, randomized, placebo-controlled clinical study. *Br J Oral Maxillofac Surg*; 39. Epub ahead of print April 2001. DOI: 10.1054/bjom.2000.0557.
  59. DeSouza SI, Rashmi MR, Vasanthi AP, et al. Mobile phones: the next step towards healthcare delivery in rural India? *PLoS One*; 9. Epub ahead of print 18 August 2014. DOI: 10.1371/journal.pone.0104895.
  60. Nasim I, Neelakantan P, Sujeer R, et al. Color stability of microfilled, microhybrid and nanocomposite resins--an in vitro study. *J Dent*; 38 Suppl 2. Epub ahead of print 2010. DOI: 10.1016/j.jdent.2010.05.020.
  61. Danda AK, Muthusekhar MR, Narayanan V, et al. Open versus closed treatment of unilateral subcondylar and condylar neck fractures: a prospective, randomized clinical study. *J Oral Maxillofac Surg*; 68. Epub ahead of print June 2010. DOI: 10.1016/j.joms.2009.09.042.
  62. Molecular structure and vibrational spectra of 2,6-bis(benzylidene)cyclohexanone: A density functional theoretical study. *Spectrochim Acta A Mol Biomol Spectrosc* 2011; 78: 113–121.
  63. Putchala MC, Ramani P, Herald J. Sherlin, et al. Ascorbic acid and its pro-oxidant activity as a therapy for tumours of oral cavity – A systematic review. *Archives of Oral Biology* 2013; 58: 563–574.
  64. Neelakantan P, Grotra D, Sharma S. Retreatability of 2 mineral trioxide aggregate-based root canal sealers: a cone-beam computed tomography analysis. *J Endod* 2013; 39: 893–896.
  65. Suresh P, Marimuthu K, Ranganathan S, et al. Optimization of machining parameters in turning of Al-SiC-Gr hybrid metal matrix composites using grey-fuzzy algorithm. *Transactions of Nonferrous Metals Society of China* 2014; 24: 2805–2814.