

**EVALUATION OF EFFICACY OF *NIGELLA SATIVA* ON GINGIVAL HEALTH**

Running title: Efficacy of *Nigella sativa* on Gingival Health

**ABSTRACT**

**Background:** Gingivitis is a reversible inflammation of the gingiva. If it is left untreated it will progress to irreversible damage to the tissues like mobility and tooth loss. Previous studies have found that use of an antibacterial agent along with scaling and root planing is ideal for treating periodontal diseases. *Nigella sativa* is a herb with excellent antioxidant, antiinflammatory and antibacterial properties and hence was used in this study.

**Aim:** The aim of the study was to evaluate the efficacy of *Nigella sativa* mouthwash on gingival health in comparison to 0.12% chlorhexidine mouthwash.

**Materials and Methods:** The study was performed as a double-blinded, randomised controlled trial research study. A total of 30 patients were selected randomly, comprising two groups, 15 in each, aged between 18 and 28 years with gingivitis. The first group was advised to oral rinse with 0.12% chlorhexidine mouthwash and the second group with new formulated *Nigella sativa* herbal mouthwash for 1 week. The plaque index and gingival index scores were calculated before administering them the mouthwashes. Then subsequently after one week the plaque and gingival index scores were taken for the 30 patients. The final data was sorted in excel and statistically analysed using IBM SPSS software analysis, unpaired t-test. The results were interpreted as graphs and tabulations.

**Results:** The group 1 patients were found to have a mean plaque index value of  $2.6\pm 0.49$  before mouthwash usage and post 1 week usage of chlorhexidine the plaque index was  $1.53\pm 0.51$ . The mean plaque index was  $2.53\pm 0.51$  prior to usage of mouthwash in group 2 and  $1.27\pm 0.45$  post usage of *Nigella sativa* mouthwash. The difference in plaque index between both the groups was found to be statistically non-significant ( $p=0.067$ ). The mean gingival index of patients who used chlorhexidine was  $2.53\pm 0.51$  before the mouthwash usage and  $1.33\pm 0.45$  after mouthwash usage. In group 2 the gingival index before the mouthwash usage was  $2.47\pm 0.51$  and after the mouthwash was  $1.47\pm 0.51$ . However the difference in gingival index between both the groups was not statistically significant ( $p=0.393$ )

**Conclusion:**

The present study reveals that the *Nigella sativa* mouthwash was more effective in plaque control when compared to the standard chlorhexidine mouthwash. Herbal mouthwash shows promising results to be used as an alternative for chlorhexidine mouthwash for management of gingivitis.

Keywords: Chlorhexidine, Gingival Health, Gingivitis, Green Synthesis, Innovative, *Nigella sativa*

## INTRODUCTION:

Periodontal disease affects the gingiva and the alveolar bone and the supporting connective tissues which anchors the tooth in the jaws (1). Periodontal disease is one of the most common chronic diseases which has affected humans for many centuries (2,3). In recent years the occurrence of dental caries has declined and periodontal disease has been the cause of tooth loss, since less teeth are lost due to caries (4). The tooth root, periodontal tooth and the alveolar bone socket are referred to as periodontium (5). Periodontal disease destroys the periodontium, it causes loss of attachment which progresses to alveolar bone loss which results in the loss of the affected teeth (6,7). The severity of the diseases depends upon the host risk factors both modifiable and non modifiable and the environment (8,9).

The two common periodontal diseases are gingivitis and periodontitis (10,11). Gingivitis is the inflammation of the gingiva which harbours the connective tissue attachment that lets the tooth remain in its position (12). This affects only the soft tissue of the gingival epithelium and its connective tissue (13). Gingivitis presents itself by swollen gums, pain and bleeding and if it is left untreated it leads to periodontitis (14–20). Periodontitis is a much more advanced form of periodontal disease and destroys both the soft tissue and the hard tissues of the tooth (21). It affects the supporting tissues and causes attachment loss of the tooth along with bone destruction (22,23). This leads to mobility of the tooth and loss of the tooth (24–28).

*Nigella sativa* is known to be a miraculous herb and it also has a religious and historical background (29). It is also known as black cumin seeds or *Kalonji*. It belongs to the family Ranunculaceae and is a flowering plant. It is native to Southwest Asia and in some southern parts of Europe and North America. It holds an important place in the practice of Unani and Ayurveda (30). The seeds of *Nigella sativa* are used in folk medicine all over the world for prevention and treating several diseases (31). It is a black boat shaped edible seed of size 2 to 4mm length .

In traditional medicine it is used to treat headaches, common colds, nasal congestion, warts, rheumatic diseases and many more (32). Many studies suggest that it helps with allergies, asthma and dyspnea (33–36). It has a very extensive and broad pharmacological use as it is an analgesic

(37), diuretic and antihypertensive (38), antidiabetic (39), antilipemic (40), antimicrobial (41), antifungal (42), antihelminthic (43), anticancer (44), anti-inflammatory (45), bronchodilator, spasmolytic and calcium antagonist (46). *Nigella sativa* is well known for its terpenoids especially thymoquinone and its fatty acids (47). Thymoquinone being an anti-inflammatory and antioxidant helps with the periodontal disease as inflammation and release of reactive oxygen species are the main offenders of periodontal disease pathogenesis (48).

Recently, the use of therapeutic herbs for the treatment of several diseases has been growing due to its promising results and rare side effects and cost effectiveness (49). Periodontal treatment along with adjunctive mouthwashes, irrigations and others improves the patient's hygiene significantly (50). Our team has extensive knowledge and research experience that has translated into high quality publications (51–70). In this context, the aim of the present study was to evaluate the efficacy of *Nigella sativa* on gingival health.

## **MATERIALS AND METHODS**

### **Study population**

The present double-blinded, parallel designed randomized clinical trial was carried out in the Department of Periodontics, Saveetha Dental College and Hospitals, Chennai, India. A total of 30 patients with gingivitis within the age group of 18-28 years were enrolled. The ethical clearance was obtained from the Institutional Ethical Committee and a written informed consent was obtained from all the study participants.

### **Inclusion criteria**

Participants within the age group of 18-28 years who were systemically healthy, presence of at least 20 teeth, probing depth of 1-3 mm, presence of bleeding on probing (BOP) in at least 30% of the sites were included in the study.

### **Exclusion criteria**

Participants allergic to herbal extracts, smokers, pregnant or lactating mothers, participants under long term medications, systemically compromised patients were excluded from the study.

### **Test solutions**

Group 1: Chlorhexidine Mouthwash (Clohex Plus® mouthwash, Dr Reddy's Lab Ltd., Hyderabad, India).

Group 2: *Nigella sativa* mouthwash formulated under laboratory conditions

### **Study design**

A pilot study was conducted using similar herbal mouthwashes to check the feasibility of the study. The prevalence of gingivitis was 80% in the pilot study. Considering the dropouts, the sample size was inflated by 20%, hence the sample size was 30 with 15 participants in each group [Group 1 (CHX mouthwash), Group 2 (*Nigella sativa* mouthwash)]. Participants were assigned to the groups by a person not involved in the study. All the subjects were provided with their assigned mouthrinses and were divided into Group 1 and Group 2 randomly using a simple lottery method with 15 participants in each group. All the mouthrinses were dispensed in identical bottles thereby ensuring subject masking. The examiner and the participants were also blinded with regard to the mouthrinse allocated to them thereby ensuring a double-blinded study. Subjects were instructed to use 10 ml of mouthwash for 1 min twice daily after tooth brushing for a period of 2 weeks.

### **Preparation of the novel herbal mouthwash :**

Dry powder of *Nigella sativa* seeds were suspended in 10 times its quantity of sterile distilled water in a flask and was kept undisturbed for 72 h at 4°C. The aqueous extract thus obtained was decanted and clarified by filtration through double-layered muslin cloth. This solution was then transferred to a porcelain dish and let it evaporate at 40°C.

The dried remnant obtained was stored for making the mouthwash solution. 200 g of the powder was suspended in polyethylene glycol, and distilled water of 800 ml and was allowed to evaporate to get the final concentrate. The final concentrate was then diluted with sterile distilled water to make a mouthwash of 20% (w/v) concentration. Two tablespoons of honey and peppermint extract were added as a natural sweetener and flavoring agent, respectively.

### **Clinical parameters**

- Silness and Loe plaque index (PI)
- Loe and Silness gingival index (GI)

Before recording the clinical parameters at baseline, thorough scaling and root planing was carried out using ultrasonic scaler, patients were advised not to use any oral hygiene aids for the next three days. After three days patients were recalled and gingival and plaque indices were recorded which was the baseline data of the study. Patients were asked to use the mouthwash for two weeks and were recalled after 14 days and both indices were recorded.

### **Statistical analysis**

The data was analyzed using Statistical Package for Social Sciences (SPSS Software, Version 23.0). Descriptive and inferential statistics were done for data summarization and presentation. Unpaired t-test was used to compare the mean values of plaque index and gingival index between the groups. The data were analyzed and interpreted as graphs and tabulations.

### **RESULTS:**


In this present study, a total of 30 gingivitis patients were enrolled for the study and were divided into two groups in which 15 were enrolled in group A (chlorhexidine mouthwash) and 15 were enrolled in group B (*Nigella sativa* mouthwash). Plaque index and gingival index of both the groups were compared at baseline and after 14 days.

In this present study we found that the plaque index had reduced in both standard chlorhexidine and *Nigella sativa* mouthwash. The mean plaque index present before mouthwash in the chlorhexidine group was  $2.6 \pm 0.49$  and in the *Nigella sativa* group was  $2.53 \pm 0.51$  before treatment. The mean plaque index which was recorded after the treatment of mouthwash was  $1.53 \pm 0.51$  in the chlorhexidine group and  $1.27 \pm 0.45$  in the *Nigella sativa* group. The plaque index values were found to be statistically non-significant ( $p=0.067$ ), unpaired t- test [Figure 1, Table 1]

Similarly we also found the gingival index of the study sample, which also had a reduction in both the groups. The gingival index which was present before the treatment was  $2.53 \pm 0.51$  in chlorhexidine group and  $2.47 \pm 0.51$  in the *Nigella sativa* group. The gingival index post treatment was noted to be  $1.33 \pm 0.45$  in chlorhexidine group and  $1.47 \pm 0.51$  in *Nigella sativa* group. The gingival index values were found to be statistically non-significant ( $p=0.393$ ), unpaired t-test [Figure 2, Table 1].


Table 1 - Comparison of mean plaque index and gingival index scores between the study groups using unpaired t test

	Levene's Test for Equality of Variances		t-test for Equality of Means						
	F	Sig	t	df	Sig(2-tailed)	Mean difference	Std. Error difference	95% Confidence Interval of the Difference	
								Lower	Upper
Mean gingival index Equal variances assumed Equal variances not assumed	3.646	.067	4.49	28	.000	-.800	.178	-1.165	-.435
			4.49	27.603	.000	-.800	.178	-1.165	-.435
Mean plaque index Equal variances assumed Equal variances not assumed	.753	.393	1.00	28	.326	.133	.133	-.140	.406
			1.00	27.858	.326	.133	.133	-.140	.407



Error Bars: +/- 1 SE

Figure 1 : Comparison of difference between chlorhexidine and *Nigella sativa* mouthwash in terms of mean plaque index scores. The x-axis depicts the types of mouthwash used in this study, chlorhexidine and *Nigella sativa* mouthwash and the y-axis represents the mean plaque index value. Blue bar depicts the preoperative plaque index scores, green bar depicts the postoperative plaque index scores. The reduction in plaque index scores was higher in the *Nigella sativa* group when compared to the chlorhexidine group. The difference between both the groups were found to be statistically not significant,  $p=0.067$  (unpaired t-test).



Error Bars: +/- 1 SE

Figure 2 : Comparison of difference between chlorhexidine and *Nigella sativa* mouthwash in terms of gingival index scores. The x-axis depicts the types of mouthwash used in this study, chlorhexidine and *Nigella sativa* mouthwash and the y-axis represents the mean gingival index value. Blue bar depicts the preoperative gingival index scores and the green bar depicts the postoperative gingival index scores. The reduction in gingival index scores was higher in the chlorhexidine group when compared to the *Nigella sativa* group. The difference between both the groups were found to be statistically not significant,  $p=0.393$  (unpaired t-test).

## DISCUSSION:

The present study was done to assess the efficacy of *Nigella sativa* on gingival health.

In this present study, we observed that the plaque index decreased in both groups. Khairnar et al found that the herbal mouthwash had a significant plaque control when compared to the standard chlorhexidine (71). A study by Balappanavar found that there was no difference of efficiency between herbal mouthwash and chlorhexidine for plaque control (72). Another study done on green tea also suggests that the herbal mouthwash helps in reducing plaque as effective as chlorhexidine mouthwash (73).

The present study also observed the gingival index reduced in both the groups post usage of the mouthwash. Al-Wafi et al found that the role of thymoquinone present in *Nigella sativa* reduced the gingival inflammation in rat models (74). Mahyari et al found that there was a significant improvement in gingival index from baseline to the end of the trial with herbal mouthwash and chlorhexidine (75). Another study by Cortelli found that there was significant reduction in gingival index post baseline time points in the case of herbal mouthrinse and that it was significantly superior when compared with cetylpyridinium chloride as standard control (76). An in vivo study also found that the herbal mouthrinse group provided the results of striking decrease in both gingival index and plaque index which was similar to our study (77).

It is found that the *Nigella sativa* mouthwash like other herbal mouthwashes are effective in reducing plaque and gingivitis. But there was no difference between the herbal and chlorhexidine mouthwashes. The natural botanical products provide a promising result in treatment of gingivitis. The exact mechanism behind the antiplaque and anti-gingival activities of it are not well known. It could be because of its components like thymoquinone or others like tannins, sterols or oils. Further molecular and clinical studies are needed to investigate the mechanism of action of *Nigella sativa* constituents, especially thymoquinone and to clarify the biomechanics behind its antiplaque and antigingivitis effect.

There are certain limitations to this present study. The duration of the study was only for two weeks and therefore the results of it cannot be generalized for a longer duration of the study.

There was a geographic limitation which had more of the South Indian population. The sample size and the duration of the study can be expanded for better results.

For future scope of the research larger sample size and inclusion of different ethnicity will provide better results. Longitudinal or periodic studies can be done to evaluate the efficiency of Nigella mouthwash on the gingival health.

## **CONCLUSION:**

The present study reveals that the *Nigella sativa* mouthwash was more effective in plaque control when compared to the standard chlorhexidine mouthwash(78) (79) (80) (81) (82) (83) (84) (85) (86) (87) (88) ((88,89) (90) (91). Herbal mouthwash shows promising results to be used as an alternative for chlorhexidine mouthwash for management of gingivitis.

## **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## **REFERENCES:**

1. Williams RC. Periodontal disease. N Engl J Med. 1990 Feb 8;322(6):373–82.

2. Evaluation of Antiplaque and Antigingivitis Effects of A Herbal Mouthwash [Internet]. Vol. 13, International Journal of Pharmaceutical Research. 2021. Available from: <http://dx.doi.org/10.31838/ijpr/2021.13.01.521>
3. Rajasekar A, Mathew MG, Others. Prevalence of Periodontal Disease among Individuals between 18-30 Years of Age: A Retrospective Study. Ann Med Health Sci Res [Internet]. 2021; Available from: <https://www.amhsr.org/abstract/prevalence-of-periodontal-disease-among-individuals-between-1830-years-of-age-a-retrospective-study-8496.html>
4. Shaw JH. Causes and control of dental caries. N Engl J Med. 1987 Oct 15;317(16):996–1004.
5. Williams RC, Zager NI. The periodontium. Textbook of oral biology WB Saunders, Philadelphia, USA. 1978;255–76.
6. Pihlstrom BL, Michalowicz BS, Johnson NW. Periodontal diseases. Lancet. 2005 Nov 19;366(9499):1809–20.
7. Kinane DF, Stathopoulou PG, Papapanou PN. Periodontal diseases [Internet]. Vol. 3, Nature Reviews Disease Primers. 2017. Available from: <http://dx.doi.org/10.1038/nrdp.2017.38>
8. Kinane DF, Peterson M, Stathopoulou PG. Environmental and other modifying factors of the periodontal diseases. Periodontol 2000. 2006;40:107–19.
9. Borrell LN, Papapanou PN. Analytical epidemiology of periodontitis. J Clin Periodontol. 2005;32 Suppl 6:132–58.
10. S TA, Thanish AS, Rajasekar A, Mathew MG. Assessment of tooth loss in chronic periodontitis patients with and without diabetes mellitus: A cross-sectional study [Internet]. Vol. 11, International Journal of Research in Pharmaceutical Sciences. 2020. p. 1927–31. Available from: <http://dx.doi.org/10.26452/ijrps.v11ispl3.3649>
11. Robo I, Heta S, Papa P, Sadiku E, Alliu N. THE IMPACT OF SMOKING ON THE HEALTH OF PERIODONTAL TISSUE [Internet]. RAD Conference Proceedings. 2017. Available from: <http://dx.doi.org/10.21175/radproc.2017.47>
12. Prevalence of Angular Bone Defects in Chronic Periodontitis Patients with and without Systemic Diseases [Internet]. Indian Journal of Forensic Medicine & Toxicology. 2020. Available from: <http://dx.doi.org/10.37506/ijfmt.v14i4.12487>
13. Newman MG, Takei H, Klokkevold PR, Carranza FA. Carranza's Clinical Periodontology. Elsevier Health Sciences; 2011. 872 p.
14. B G, Geethika B, Rajasekar A, Chaudary M. Comparison of periodontal status among pregnant and non-pregnant women [Internet]. Vol. 11, International Journal of Research in Pharmaceutical Sciences. 2020. p. 1923–6. Available from: <http://dx.doi.org/10.26452/ijrps.v11ispl3.3648>

15. Rajasekar A, Lecturer S, Department of Periodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, et al. Assessment Of Periodontal Status among Post Menopausal Women: A Retrospective Study [Internet]. *International Journal of Dentistry and Oral Science*. 2020. p. 1063–6. Available from: <http://dx.doi.org/10.19070/2377-8075-20000210>
16. Kandhan T, Rajasekar A. Prevalence of Periodontal Diseases Among Patients with And Without Systemic Diseases - A Retrospective Study [Internet]. Vol. 11, *Journal of Complementary Medicine Research*. 2020. p. 155. Available from: <http://dx.doi.org/10.5455/jcmr.2020.11.04.20>
17. Assessment of periodontal health among patients with diabetes mellitus: a retrospective study [Internet]. Vol. 26, *Journal of Contemporary Issues in Business and Government*. 2021. Available from: <http://dx.doi.org/10.47750/cibg.2020.26.02.004>
18. Assessment of Gender Based Difference in Occurrence of Periodontal Diseases: A Retrospective Study [Internet]. Vol. 27, *Journal of Contemporary Issues in Business and Government*. 2021. Available from: <http://dx.doi.org/10.47750/cibg.2021.27.02.064>
19. Evaluation of Periodontal Health Adjacent to Class V Restoration [Internet]. Vol. 27, *Journal of Contemporary Issues in Business and Government*. 2021. Available from: <http://dx.doi.org/10.47750/cibg.2021.27.02.040>
20. Prevalence of Periodontal Diseases Among Individuals Above 45 Years: A Retrospective Study [Internet]. Vol. 27, *Journal of Contemporary Issues in Business and Government*. 2021. Available from: <http://dx.doi.org/10.47750/cibg.2021.27.02.065>
21. Azodo C, Erhabor P. Management of tooth mobility in the periodontology clinic: An overview and experience from a tertiary healthcare setting. *Afr J Med Health Sci*. 2016;15(1):50.
22. Preber H. Cigarette Smoking and Periodontal Disease: Clinical and Therapeutic Aspects. 1986. 125 p.
23. Flemmig TF. Periodontitis. *Ann Periodontol*. 1999 Dec;4(1):32–8.
24. Rajeshwaran N, Rajasekar A, Kaarthikeyan G. Prevalence of Pathologic Migration in Patients with Periodontitis: A Retrospective Analysis [Internet]. Vol. 11, *Journal of Complementary Medicine Research*. 2020. p. 172. Available from: <http://dx.doi.org/10.5455/jcmr.2020.11.04.22>
25. Prevalence of Tooth Loss Among Chronic Periodontitis Patients - A Retrospective Study [Internet]. Vol. 12, *International Journal of Pharmaceutical Research*. 2020. Available from: <http://dx.doi.org/10.31838/ijpr/2020.sp2.239>
26. Murthykumar K, Rajasekar A, Kaarthikeyan G. Assessment of various treatment modalities for isolated gingival recession defect- A retrospective study [Internet]. Vol. 11, *International Journal of Research in Pharmaceutical Sciences*. 2020. p. 3–7. Available from:

<http://dx.doi.org/10.26452/ijrps.v11ispl3.2824>

27. Sabarathinam J, Rajasekar A, Madhulaxmi M. Prevalence of Furcation Involvement Among Patients with Periodontitis: A Cross Sectional Study [Internet]. Vol. 11, International Journal of Research in Pharmaceutical Sciences. 2020. p. 1483–7. Available from: <http://dx.doi.org/10.26452/ijrps.v11ispl3.3457>
28. Ong G. Periodontal disease and tooth loss. *Int Dent J*. 1998 Jun;48(3 Suppl 1):233–8.
29. Goreja WG. Black Seed: Nature's Miracle Remedy. Amazing Herbs Press; 2003. 54 p.
30. Sharma PC, Yelne MB, Dennis TJ. Database on medicinal plants used in ayurveda & siddha. Volume 5. 2008;
31. Yarnell E, Abascal K. *Nigella sativa*: Holy Herb of the Middle East. *Focus Altern Complement Ther*. 2011 Apr 1;17(2):99–105.
32. Kooti W, Hasanzadeh-Noohi Z, Sharafi-Ahvazi N, Asadi-Samani M, Ashtary-Larky D. Phytochemistry, pharmacology, and therapeutic uses of black seed (*Nigella sativa*). *Chin J Nat Med*. 2016 Oct;14(10):732–45.
33. Avicenna. Canon of Medicine Volume 2: Natural Pharmaceuticals. Great Books of the Islamic World, Incorporated; 2012. 1400 p.
34. Boskabady MH, Mohsenpoor N, Takaloo L. Antiasthmatic effect of *Nigella sativa* in airways of asthmatic patients. *Phytomedicine*. 2010 Aug;17(10):707–13.
35. Nikakhlagh S, Rahim F, Aryani FHN, Syahpoush A, Brougerdnya MG, Saki N. Herbal treatment of allergic rhinitis: the use of *Nigella sativa*. *Am J Otolaryngol*. 2011 Sep;32(5):402–7.
36. Kalus U, Pruss A, Bystron J, Jurecka M, Smekalova A, Lichius JJ, et al. Effect of *Nigella sativa* (black seed) on subjective feeling in patients with allergic diseases. *Phytother Res*. 2003 Dec;17(10):1209–14.
37. Abdel-Fattah M, Abdel-Fattah A-FM, Matsumoto K, Watanabe H. Antinociceptive effects of *Nigella sativa* oil and its major component, thymoquinone, in mice [Internet]. Vol. 400, *European Journal of Pharmacology*. 2000. p. 89–97. Available from: [http://dx.doi.org/10.1016/s0014-2999\(00\)00340-x](http://dx.doi.org/10.1016/s0014-2999(00)00340-x)
38. Zaoui A, Cherrah Y, Lacaille-Dubois MA, Settaf A, Amarouch H, Hassar M. [Diuretic and hypotensive effects of *Nigella sativa* in the spontaneously hypertensive rat]. *Therapie*. 2000 May;55(3):379–82.
39. Hassanin NI, Hassan FM. A preliminary study on the effect of *Nigella sativa* L. seeds on hypoglycemia. *Vet Med J* [Internet]. 1996; Available from: <https://agris.fao.org/agris-search/search.do?recordID=EG9702048>

40. Danladi J. Hepatoprotective Effect of Black Seed (*Nigella sativa*) oil on Carbon Tetrachloride (CCl<sub>4</sub>) Induced Liver Toxicity in Adult Wistar Rats [Internet]. Vol. 4, IOSR Journal of Dental and Medical Sciences. 2013. p. 56–62. Available from: <http://dx.doi.org/10.9790/0853-0435662>
41. Sokmen A, Jones BM, Erturk M. The in vitro antibacterial activity of Turkish medicinal plants. *J Ethnopharmacol.* 1999 Oct;67(1):79–86.
42. Khan MAU, Ashfaq MK, Zuberi HS, Mahmood MS, Gilani AH. The in vivo antifungal activity of the aqueous extract from *Nigella sativa* seeds. *Phytother Res.* 2003 Feb;17(2):183–6.
43. Akhtar MS, Riffat S. Field trial of *Saussurea lappa* roots against nematodes and *Nigella sativa* seeds against cestodes in children. *J Pak Med Assoc.* 1991 Aug;41(8):185–7.
44. Worthen DR, Ghosheh OA, Crooks PA. The in vitro anti-tumor activity of some crude and purified components of blackseed, *Nigella sativa* L. *Anticancer Res.* 1998 May;18(3A):1527–32.
45. Mutabagani A, El-Mahdy SAM. A study of the anti-inflammatory activity of *Nigella sativa* L. and thymoquinone in rats. *Saudi Pharmaceutical Journal.* 1997;5:110–3.
46. Gilani AH, Aziz N, Khurram IM, Chaudhary KS, Iqbal A. Bronchodilator, spasmolytic and calcium antagonist activities of *Nigella sativa* seeds (Kalonji): a traditional herbal product with multiple medicinal uses. *J Pak Med Assoc.* 2001 Mar;51(3):115–20.
47. Akram Khan M, Afzal M. Chemical composition of *Nigella sativa* Linn: Part 2 Recent advances. *Inflammopharmacology.* 2016 Jun;24(2-3):67–79.
48. Varela-López A, Bullón P, Giampieri F, Quiles JL. Non-Nutrient, Naturally Occurring Phenolic Compounds with Antioxidant Activity for the Prevention and Treatment of Periodontal Diseases. *Antioxidants (Basel).* 2015 Jun 24;4(3):447–81.
49. Eid Abdelmagyd HA, Ram Shetty DS, Musa Musleh Al-Ahmari DM. Herbal medicine as adjunct in periodontal therapies- A review of clinical trials in past decade. *J Oral Biol Craniofac Res.* 2019 Jul;9(3):212–7.
50. Herrera D, Matesanz P, Martín C, Oud V, Feres M, Teughels W. Adjunctive effect of locally delivered antimicrobials in periodontitis therapy: A systematic review and meta-analysis. *J Clin Periodontol.* 2020 Jul;47 Suppl 22(S22):239–56.
51. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol.* 2018 Oct;89(10):1241–8.
52. Paramasivam A, Priyadharsini JV, Raghunandhakumar S, Elumalai P. A novel COVID-19 and its effects on cardiovascular disease. *Hypertens Res.* 2020 Jul;43(7):729–30.

53. S G, T G, K V, Faleh A A, Sukumaran A, P N S. Development of 3D scaffolds using nanochitosan/silk-fibroin/hyaluronic acid biomaterials for tissue engineering applications. *Int J Biol Macromol*. 2018 Dec;120(Pt A):876–85.
54. Del Fabbro M, Karanxha L, Panda S, Bucchi C, Nadathur Doraiswamy J, Sankari M, et al. Autologous platelet concentrates for treating periodontal infrabony defects. *Cochrane Database Syst Rev*. 2018 Nov 26;11:CD011423.
55. Paramasivam A, Vijayashree Priyadharsini J. MitomiRs: new emerging microRNAs in mitochondrial dysfunction and cardiovascular disease. *Hypertens Res*. 2020 Aug;43(8):851–3.
56. Jayaseelan VP, Arumugam P. Dissecting the theranostic potential of exosomes in autoimmune disorders. *Cell Mol Immunol*. 2019 Dec;16(12):935–6.
57. Vellappally S, Al Kheraif AA, Divakar DD, Basavarajappa S, Anil S, Fouad H. Tooth implant prosthesis using ultra low power and low cost crystalline carbon bio-tooth sensor with hybridized data acquisition algorithm. *Comput Commun*. 2019 Dec 15;148:176–84.
58. Vellappally S, Al Kheraif AA, Anil S, Assery MK, Kumar KA, Divakar DD. Analyzing Relationship between Patient and Doctor in Public Dental Health using Particle Memetic Multivariable Logistic Regression Analysis Approach (MLRA2). *J Med Syst*. 2018 Aug 29;42(10):183.
59. Varghese SS, Ramesh A, Veeraiyan DN. Blended Module-Based Teaching in Biostatistics and Research Methodology: A Retrospective Study with Postgraduate Dental Students. *J Dent Educ*. 2019 Apr;83(4):445–50.
60. Venkatesan J, Singh SK, Anil S, Kim S-K, Shim MS. Preparation, Characterization and Biological Applications of Biosynthesized Silver Nanoparticles with Chitosan-Fucoidan Coating. *Molecules* [Internet]. 2018 Jun 12;23(6). Available from: <http://dx.doi.org/10.3390/molecules23061429>
61. Alsubait SA, Al Ajlan R, Mitwalli H, Aburaisi N, Mahmood A, Muthurangan M, et al. Cytotoxicity of Different Concentrations of Three Root Canal Sealers on Human Mesenchymal Stem Cells. *Biomolecules* [Internet]. 2018 Aug 1;8(3). Available from: <http://dx.doi.org/10.3390/biom8030068>
62. Venkatesan J, Rekha PD, Anil S, Bhatnagar I, Sudha PN, Dechsakulwatana C, et al. Hydroxyapatite from Cuttlefish Bone: Isolation, Characterizations, and Applications. *Biotechnol Bioprocess Eng*. 2018 Aug 1;23(4):383–93.
63. Vellappally S, Al Kheraif AA, Anil S, Wahba AA. IoT medical tooth mounted sensor for monitoring teeth and food level using bacterial optimization along with adaptive deep learning neural network. *Measurement*. 2019 Mar 1;135:672–7.
64. PradeepKumar AR, Shemesh H, Nivedhitha MS, Hashir MMJ, Arockiam S, Uma Maheswari TN, et al. Diagnosis of Vertical Root Fractures by Cone-beam Computed

Tomography in Root-filled Teeth with Confirmation by Direct Visualization: A Systematic Review and Meta-Analysis. *J Endod.* 2021 Aug;47(8):1198–214.

65. R H, Ramani P, Tilakaratne WM, Sukumaran G, Ramasubramanian A, Krishnan RP. Critical appraisal of different triggering pathways for the pathobiology of pemphigus vulgaris-A review. *Oral Dis* [Internet]. 2021 Jun 21; Available from: <http://dx.doi.org/10.1111/odi.13937>
66. Ezhilarasan D, Lakshmi T, Subha M, Deepak Nallasamy V, Raghunandhakumar S. The ambiguous role of sirtuins in head and neck squamous cell carcinoma. *Oral Dis* [Internet]. 2021 Feb 11; Available from: <http://dx.doi.org/10.1111/odi.13798>
67. Sarode SC, Gondivkar S, Sarode GS, Gadbail A, Yuwanati M. Hybrid oral potentially malignant disorder: A neglected fact in oral submucous fibrosis. *Oral Oncol.* 2021 Jun 16;105390.
68. Kavarthapu A, Gurumoorthy K. Linking chronic periodontitis and oral cancer: A review. *Oral Oncol.* 2021 Jun 14;105375.
69. Vellappally S, Abdullah Al-Kheraif A, Anil S, Basavarajappa S, Hassanein AS. Maintaining patient oral health by using a xeno-genetic spiking neural network. *J Ambient Intell Humaniz Comput* [Internet]. 2018 Dec 14; Available from: <https://doi.org/10.1007/s12652-018-1166-8>
70. Aldhuwayhi S, Mallineni SK, Sakhamuri S, Thakare AA, Mallineni S, Sajja R, et al. Covid-19 Knowledge and Perceptions Among Dental Specialists: A Cross-Sectional Online Questionnaire Survey. *Risk Manag Healthc Policy.* 2021 Jul 7;14:2851–61.
71. Khairnar MS, Pawar B, Marawar PP, Mani A. Evaluation of *Calendula officinalis* as an anti-plaque and anti-gingivitis agent. *J Indian Soc Periodontol.* 2013 Nov;17(6):741–7.
72. Balappanavar AY, Sardana V, Singh M. Comparison of the effectiveness of 0.5% tea, 2% neem and 0.2% chlorhexidine mouthwashes on oral health: a randomized control trial. *Indian J Dent Res.* 2013 Jan;24(1):26–34.
73. Priya BM, Anitha V, Shanmugam M, Ashwath B, Sylva SD, Vigneshwari SK. Efficacy of chlorhexidine and green tea mouthwashes in the management of dental plaque-induced gingivitis: A comparative clinical study. *Contemp Clin Dent.* 2015 Oct;6(4):505–9.
74. Al-Wafi HA. Tufts University school of Dental Medicine [Internet]. 2014. Available from: <https://search.proquest.com/openview/2046fe13104528afc21cd67dad7de375/1?pq-origsite=gscholar&cbl=18750>
75. Mahyari S, Mahyari B, Emami SA, Malaekheh-Nikouei B, Jahanbakhsh SP, Sahebkar A, et al. Evaluation of the efficacy of a polyherbal mouthwash containing *Zingiber officinale*, *Rosmarinus officinalis* and *Calendula officinalis* extracts in patients with gingivitis: A randomized double-blind placebo-controlled trial. *Complement Ther Clin Pract.* 2016 Feb;22:93–8.

76. Cortelli SC, Cortelli JR, Shang H, Costa R, Charles CA. Gingival health benefits of essential-oil and cetylpyridinium chloride mouthrinses: a 6-month randomized clinical study. *Am J Dent*. 2014 Jun;27(3):119–26.
77. Charles CA, McGuire JA, Qaqish J, Amini P. Increasing antiplaque/antigingivitis efficacy of an essential oil mouthrinse over time: an in vivo study. *Gen Dent*. 2013 Jan;61(1):23–8.
78. Danda AK. Comparison of a single noncompression miniplate versus 2 noncompression miniplates in the treatment of mandibular angle fractures: a prospective, randomized clinical trial. *J Oral Maxillofac Surg*. 2010 Jul;68(7):1565–7.
79. Robert R, Justin Raj C, Krishnan S, Jerome Das S. Growth, theoretical and optical studies on potassium dihydrogen phosphate (KDP) single crystals by modified Sankaranarayanan–Ramasamy (mSR) method [Internet]. Vol. 405, *Physica B: Condensed Matter*. 2010. p. 20–4. Available from: <http://dx.doi.org/10.1016/j.physb.2009.08.015>
80. Krishnan V, Lakshmi T. Bioglass: A novel biocompatible innovation. *J Adv Pharm Technol Res*. 2013 Apr;4(2):78–83.
81. Soh CL, Narayanan V. Quality of life assessment in patients with dentofacial deformity undergoing orthognathic surgery—A systematic review [Internet]. Vol. 42, *International Journal of Oral and Maxillofacial Surgery*. 2013. p. 974–80. Available from: <http://dx.doi.org/10.1016/j.ijom.2013.03.023>
82. Lekha L, Kanmani Raja K, Rajagopal G, Easwaramoorthy D. Schiff base complexes of rare earth metal ions: Synthesis, characterization and catalytic activity for the oxidation of aniline and substituted anilines [Internet]. Vol. 753, *Journal of Organometallic Chemistry*. 2014. p. 72–80. Available from: <http://dx.doi.org/10.1016/j.jorganchem.2013.12.014>
83. Dhinesh B, Isaac JoshuaRamesh Lalvani J, Parthasarathy M, Annamalai K. An assessment on performance, emission and combustion characteristics of single cylinder diesel engine powered by *Cymbopogon flexuosus* biofuel [Internet]. Vol. 117, *Energy Conversion and Management*. 2016. p. 466–74. Available from: <http://dx.doi.org/10.1016/j.enconman.2016.03.049>
84. PradeepKumar AR, Shemesh H, Jothilatha S, Vijayabharathi R, Jayalakshmi S, Kishen A. Diagnosis of Vertical Root Fractures in Restored Endodontically Treated Teeth: A Time-dependent Retrospective Cohort Study. *J Endod*. 2016 Aug;42(8):1175–80.
85. Vijayakumar GNS, Nixon Samuel Vijayakumar G, Devashankar S, Rathnakumari M, Sureshkumar P. Synthesis of electrospun ZnO/CuO nanocomposite fibers and their dielectric and non-linear optic studies [Internet]. Vol. 507, *Journal of Alloys and Compounds*. 2010. p. 225–9. Available from: <http://dx.doi.org/10.1016/j.jallcom.2010.07.161>
86. Kavitha M, Subramanian R, Narayanan R, Udhayabanu V. Solution combustion synthesis and characterization of strontium substituted hydroxyapatite nanocrystals [Internet]. Vol. 253, *Powder Technology*. 2014. p. 129–37. Available from:

<http://dx.doi.org/10.1016/j.powtec.2013.10.045>

87. Sahu D, Kannan GM, Vijayaraghavan R. Size-Dependent Effect of Zinc Oxide on Toxicity and Inflammatory Potential of Human Monocytes [Internet]. Vol. 77, *Journal of Toxicology and Environmental Health, Part A*. 2014. p. 177–91. Available from: <http://dx.doi.org/10.1080/15287394.2013.853224>
88. Neelakantan P, Cheng CQ, Mohanraj R, Sriraman P, Subbarao C, Sharma S. Antibiofilm activity of three irrigation protocols activated by ultrasonic, diode laser or Er:YAG laser in vitro [Internet]. Vol. 48, *International Endodontic Journal*. 2015. p. 602–10. Available from: <http://dx.doi.org/10.1111/iej.12354>
89. Lekha L, Kanmani Raja K, Rajagopal G, Easwaramoorthy D. Synthesis, spectroscopic characterization and antibacterial studies of lanthanide(III) Schiff base complexes containing N, O donor atoms [Internet]. Vols. 1056-1057, *Journal of Molecular Structure*. 2014. p. 307–13. Available from: <http://dx.doi.org/10.1016/j.molstruc.2013.10.014>
90. Gopalakannan S, Senthilvelan T, Ranganathan S. Modeling and Optimization of EDM Process Parameters on Machining of Al 7075-B4C MMC Using RSM [Internet]. Vol. 38, *Procedia Engineering*. 2012. p. 685–90. Available from: <http://dx.doi.org/10.1016/j.proeng.2012.06.086>
91. Parthasarathy M, Isaac Joshua Ramesh Lalvani J, Dhinesh B, Annamalai K. Effect of hydrogen on ethanol-biodiesel blend on performance and emission characteristics of a direct injection diesel engine. *Ecotoxicol Environ Saf*. 2016 Dec;134(Pt 2):433–9.