

Preparation of ethanolic extract of cassia auriculata and its anti-diabetic activity

Running title: cassia auriculata and its diabetic activities

Abstract:

Introduction:

Diabetes has caused a major burden to the health sector in the developing countries and has shown an increasing trend among the urban population. It is estimated that most patients are with type II diabetes which could be easily treated with dietary changes, exercise, and medication. Sri Lanka carries a long history of ayurvedic medicine where it uses the plant for treating many diseases. Therefore it is important to screen medicinal plants scientifically so they could be used safely and effectively in the traditional medical system and also be used for further investigations. cassia auriculata is a plant used in the Ayurvedic medical system in Sri Lanka for treating many diseases including diabetics. We evaluated the anti-diabetic properties and the antioxidant properties of cassia auriculata leaves.

Methods:

The methanol extract of the leaves was sequentially extracted with petroleum ether and thereafter was partitioned between EtOAc, and water. The α -amylase inhibition assay was performed using the 3,5- dinitrosalicylic acid method. The antioxidant activities were measured using the DPPH free radical scavenging activity and the total phenolic content using Folin-Ciocalteu's reagent. The cytotoxicity of the extract was evaluated using the Brine shrimp bioassay.

Results: The extract shows very good anti-diabetic activity for the *Cassia auriculata* extract by using BSA and EAA Assay.

Conclusion:

The leaf extracts of cassia auriculata exhibit remarkable α -amylase inhibitory activity in the crude methanolic extract. Hence leaves of cassia auriculata have a potential to be used as a regular green vegetable and also be investigated further in isolating pure compounds with anti-diabetic activity.

Keywords: Cassia auriculata, Diabetic Activities, Ethanolic extract, α -Amylase Assay, α -Glucosidase Assay.

Introduction:

Diabetes is a disorder of β -cells of langerhans that can be diagnosed by its symptoms like weight loss, excessive hunger, thirst and urination(Dua, Wadhwa, Singhvi, Rapalli, Shukla, Shastri, Gupta, Satija, Mehta, Khurana, Awasthi, Maurya, Thangavelu, Rajeshkumar, *et al.*, 2019),(Ramesh *et al.*, 2018a). It is a fifth leading cause of deaths as well the leading cause of adult blindness, responsible for heart attacks, strokes and gangrenous leg amputations. Cassia auriculata Linn (Kingdom : Plantae, Order : Fabales, Family : Fabaceae, Sub family : Caesalpinioideae, Tribe : Cassie,Genius : Cassia,Species : C. auriculata),(Ezhilarasan, 2018a),(A. C. Gomathi *et al.*, 2020a) commonly known as Tanners Senna, is distributed throughout hot deciduous forests of India and holds a very prestigious position in Ayurveda and Siddha systems of medicine(Rajeshkumar, Venkat Kumar, *et al.*, 2018),(Ezhilarasan, Sokal and Najimi, 2018a). The plant is used in the traditional system of medicine for urinary disorders, female antifertility, leprosy, worm infestation, diarrhoea, disease of pittam; leaves, flowers and fruits as anthelmintic; seeds for eye troubles, diabetes(Nandhini, Rajeshkumar and Mythili, 2019a).

Non-pharmacological and pharmacological methods to diabetes treatment are used(Veerasamy *et al.*, 2021). Exercise, food management, and surgery are examples of non-pharmacological approaches, whereas medicines like insulin and oral hypoglycemic treatments are examples of pharmacological approaches(Vairavel, Devaraj and Shanmugam, 2020a)s. Conventional medicines are not only expensive, but they also come with a slew of side effects(M. Gomathi *et al.*, 2020). For the treatment of diabetes, many herbal remedies have been advocated(Rajasekaran *et al.*, 2020a),(Ramesh *et al.*, 2018a). Medicinal plant components are considered to operate on a number of targets through a variety of modes and processes(2018 Conference on Signal Processing and Communication Engineering Systems (SPACES): 4-5 Jan. 2018, 2018),(Gheena and Ezhilarasan, 2019a).They have the potential to treat complex diseases such as diabetes and its consequences(R *et al.*, 2020),(Saravanan *et al.*, 2018a).

This research is needed to know the importance of cassia auriculata in anti- diabetic activity.The main deficiency it fulfill that the Cassia auriculata is related to histamine,kinn and prostaglandin inhibiting activity.Our team has extensive knowledge and research experience that has translate

into high quality publications(Rajeshkumar, Kumar, *et al.*, 2018; Nandhini, Rajeshkumar and Mythili, 2019b; M. Gomathi *et al.*, 2020; Rajasekaran *et al.*, 2020b; Vairavel, Devaraj and Shanmugam, 2020b),(Santhoshkumar *et al.*, 2019),(Raj R, D and S, 2020),(Saravanan *et al.*, 2018b),(Gheena and Ezhilarasan, 2019b),(Ezhilarasan, Sokal and Najimi, 2018b),(Ezhilarasan, 2018b),(Dua, Wadhwa, Singhvi, Rapalli, Shukla, Shastri, Gupta, Satija, Mehta, Khurana, Awasthi, Maurya, Thangavelu, S, *et al.*, 2019; A. C. Gomathi *et al.*, 2020b; Vairavel, Devaraj and Shanmugam, 2020b),(Ramesh *et al.*, 2018b; Duraisamy *et al.*, 2019; Ezhilarasan, Apoorva and Ashok Vardhan, 2019; Arumugam, George and Jayaseelan, 2021; Joseph and Prasanth, 2021) .,(Gnanavel, Roopan and Rajeshkumar, 2019),(Markov *et al.*, 2021)(Veerasamy *et al.*, 2021).The aim of this study is to determine anti-diabetic activity of cassia auriculata flower extract.

Materials and Methods:

In vitro α -amylase inhibitory studies The α -amylase inhibition assay was performed using the 3,5-dinitrosalicylic acid (DNSA) method. The leaf extract of *cassia auriculata*¹ was dissolved in minimum amount of 10% DMSO and was further dissolved in buffer ((Na₂HPO₄/NaH₂PO₄)² (0.02 M), NaCl (0.006 M) at pH 6.9) to give concentrations ranging from 10 to 1000 μ g/ml. A volume of 200 μ l of α -amylase solution (2 units/ml) was mixed with 200 μ l of the extract and was incubated for 10 min at 30 °C. Thereafter 200 μ l of the starch solution (1% in water (w/v)) was added to each tube and incubated for 3 min. The reactivity was terminated by the addition of 200 μ l DNSA reagent (12 g of sodium potassium tartrate tetrahydrate in 8.0 mL of 2 M NaOH and 20 mL of 96 mM of 3,5-dinitrosalicylic acid solution) and was boiled for 10 min in a water bath at 85–90 °C. The mixture was cooled to ambient temperature and was diluted with 5 ml of distilled water, and the absorbance was measured at 540 nm using a UV-Visible spectrophotometer. The blank with 100% enzyme activity was prepared by replacing the plant extract with 200 μ l of buffer. A blank reactivity was similarly prepared using the plant extract at each concentration in the absence of the enzyme solution. A positive control sample was prepared using Acarbose (100 μ g/ml–2 μ g/ml) and the reactivity was performed similarly to the reactivity with plant extract as mentioned above. The α -amylase inhibitory activity was expressed as percent inhibition and was calculated using the equation given below: The % α -amylase inhibition was plotted against the extract concentration and the IC₅₀ values were obtained from the graph. The test was realized following the protocol of Pistia-Brueggeman and Hollingsworth. An amount³ of 50 μ L of WFLE extract was prepared at various concentrations (1, 3, 7, 15, 31, 62, 125, 250, 500 and 1000 μ g/mL) and incubated with the solution containing 10 μ L of α -glucosidase (maltase) 1 U/mL and 125 μ L of 0.1 M phosphate buffer (pH 6.8) for 20 min at 37 °C. To start the reaction, a solution of 20 μ L of 1 M pNPG (4-Nitrophenyl β -D-glucopyranoside) (substrate) was added then incubated for half-hour. To terminate the reaction, 50 μ L of 0.1 N Na₂CO₃ was added. The optical density was measured at 405 nm using a spectrophotometer. For this assay, acarbose was used as a positive control (α -amylase inhibitor).

The experiments were repeated three times for each concentration. The α -Glucosidase inhibitory activity was calculated by using the following formula: Extract inhibitory activity = $[(XA - XB)/XA] \times 100$. XA is the absorbance of the control (100% enzyme activity) and XB is the absorbance of the sample. After determining the α -glucosidase inhibitory activity of the different concentrations, the IC₅₀ values were determined for the acarbose and WFLE extracts (concentration required to inhibit 50% of α -glucosidase).

Results:




Image 1: *Cassia auriculata* ethanolic extract

The results of anti-diabetic activity and egg albumin assay were depicted in (Figures 1-2). In the present study, the total anti-diabetic of *Cassia auriculata ethanolic* extract (CAE) was determined using the egg albumin assay method. CAE Ext showed anti-diabetic property in a concentration dependent manner. The result indicated that the CAE Ext significantly (<0.05) inhibited albumin Denaturation Assay method. Egg albumin assay is an easy, rapid and sensitive method for the anti-diabetic screening of plant extracts. The present study investigated the anti-diabetic activity of CAE Ext, and expressed the inhibition of albumin denaturation Assay using BSA as standard reference.




Figure 1: The above graph depicts the anti-diabetic activity with an increased percentage of inhibition with a concentration in microlitres. X axis denotes concentration and Y axis denotes percentage of inhibition of *Cassia auriculata*.⁴




Figure 2: The above graph depicts the anti-diabetic activity of *Cassia auriculata* ethanolic extract on egg albumin assay increased percentage of inhibition with a concentration in microlitres. X axis denotes concentration and Y axis denotes percentage of inhibition of *Cassia auriculata*.⁵

Discussion:

The *Cassia auriculata* leaves were used as a sample to evaluate the anti-diabetic activity for the reason, increased diabetic patients and there were less treatment procedures and also less drugs. They are more resistant when compared to *cassia auriculata* extracts. Hence plaque samples were used rather than standard strains. Other study done by anti-diabetic effect of *Cassia auriculata*, which studied the anti-diabetic property of *Cassia auriculata* has been evaluated against *Staphylococcus aureus*, *Enterococcus faecalis*, *Bacillus subtilis*, *Salmonella typhi*, *Salmonella paratyphi A*, *Escherichia coli*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Vibrio cholerae* and *Shigella dysenteriae*. But this study did not evaluate the effect of *Cassia auriculata* against diabetes. Study was done to evaluate the anti-diabetic activity of *Cassia auriculata*. In this study, it was shown that anti-diabetic activity.⁶

Conclusion:

From the present study, we can conclude that ethanolic extract of *Cassia auriculata* flower extract has anti-diabetic activity. But for more precise effect in a specific way, more studies with specific strain should be done.

NOTE:

The study highlights the efficacy of " ayurvedic medicine " which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

Reference:

2018 Conference on Signal Processing and Communication Engineering Systems (SPACES): 4-5 Jan. 2018 (2018).

Arumugam, P., George, R. and Jayaseelan, V. P. (2021) 'Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma', *Archives of oral biology*, 122, p. 105030.

Dua, K., Wadhwa, R., Singhvi, G., Rapalli, V., Shukla, S. D., Shastri, M. D., Gupta, G., Satija, S., Mehta, M., Khurana, N., Awasthi, R., Maurya, P. K., Thangavelu, L., Rajeshkumar, S., *et al.* (2019) 'The potential of siRNA based drug delivery in respiratory disorders: Recent advances and progress', *Drug Development Research*, pp. 714–730. doi: 10.1002/ddr.21571.

Dua, K., Wadhwa, R., Singhvi, G., Rapalli, V., Shukla, S. D., Shastri, M. D., Gupta, G., Satija, S., Mehta, M., Khurana, N., Awasthi, R., Maurya, P. K., Thangavelu, L., S, R., *et al.* (2019) 'The potential of siRNA based drug delivery in respiratory disorders: Recent advances and progress', *Drug development research*, 80(6), pp. 714–730.

Duraisamy, R. *et al.* (2019) 'Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments', *Implant dentistry*, 28(3), pp. 289–295.

Ezhilarasan, D. (2018a) 'Oxidative stress is bane in chronic liver diseases: Clinical and experimental perspective', *Arab Journal of Gastroenterology*, pp. 56–64. doi: 10.1016/j.ajg.2018.03.002.

Ezhilarasan, D. (2018b) 'Oxidative stress is bane in chronic liver diseases: Clinical and experimental perspective', *Arab journal of gastroenterology: the official publication of the Pan-Arab Association of Gastroenterology*, 19(2), pp. 56–64.

Ezhilarasan, D., Apoorva, V. S. and Ashok Vardhan, N. (2019) 'Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(2), pp. 115–121.

Ezhilarasan, D., Sokal, E. and Najimi, M. (2018a) 'Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets', *Hepatobiliary & Pancreatic Diseases International*, pp. 192–197. doi: 10.1016/j.hbpd.2018.04.003.

Ezhilarasan, D., Sokal, E. and Najimi, M. (2018b) 'Hepatic fibrosis: It is time to go with hepatic

stellate cell-specific therapeutic targets', *Hepatobiliary & pancreatic diseases international: HBPD INT*, 17(3), pp. 192–197.

Gheena, S. and Ezhilarasan, D. (2019a) 'Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells', *Human & Experimental Toxicology*, pp. 694–702. doi: 10.1177/0960327119839173.

Gheena, S. and Ezhilarasan, D. (2019b) 'Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells', *Human & experimental toxicology*, 38(6), pp. 694–702.

Gnanavel, V., Roopan, S. M. and Rajeshkumar, S. (2019) 'Aquaculture: An overview of chemical ecology of seaweeds (food species) in natural products', *Aquaculture*, 507, pp. 1–6.

Gomathi, A. C. *et al.* (2020a) 'Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of Tamarindus indica on MCF-7 human breast cancer cell line', *Journal of Drug Delivery Science and Technology*, p. 101376. doi: 10.1016/j.jddst.2019.101376.

Gomathi, A. C. *et al.* (2020b) 'Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of Tamarindus indica on MCF-7 human breast cancer cell line', *Journal of drug delivery science and technology*, 55, p. 101376.

Gomathi, M. *et al.* (2020) 'Green synthesis of silver nanoparticles using Gymnema sylvestre leaf extract and evaluation of its antibacterial activity', *South African Journal of Chemical Engineering*, pp. 1–4. doi: 10.1016/j.sajce.2019.11.005.

Joseph, B. and Prasanth, C. S. (2021) 'Is photodynamic therapy a viable antiviral weapon against COVID-19 in dentistry?', *Oral surgery, oral medicine, oral pathology and oral radiology*, pp. 118–119.

Markov, A. *et al.* (2021) 'Mesenchymal stem/stromal cells as a valuable source for the treatment of immune-mediated disorders', *Stem cell research & therapy*, 12(1), p. 192.

Nandhini, N. T., Rajeshkumar, S. and Mythili, S. (2019a) 'The possible mechanism of eco-friendly synthesized nanoparticles on hazardous dyes degradation', *Biocatalysis and Agricultural Biotechnology*, p. 101138. doi: 10.1016/j.bcab.2019.101138.

Nandhini, N. T., Rajeshkumar, S. and Mythili, S. (2019b) 'The possible mechanism of eco-friendly synthesized nanoparticles on hazardous dyes degradation', *Biocatalysis and agricultural biotechnology*, 19, p. 101138.

Rajasekaran, S. *et al.* (2020a) 'Collective influence of 1-decanol addition, injection pressure and EGR on diesel engine characteristics fueled with diesel/LDPE oil blends', *Fuel*, p. 118166. doi: 10.1016/j.fuel.2020.118166.

Rajasekaran, S. *et al.* (2020b) 'Collective influence of 1-decanol addition, injection pressure and EGR on diesel engine characteristics fueled with diesel/LDPE oil blends', *Fuel*, 277, p. 118166.

Rajeshkumar, S., Venkat Kumar, S., *et al.* (2018) 'Biosynthesis of zinc oxide nanoparticles

using *Mangifera indica* leaves and evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells', *Enzyme and Microbial Technology*, pp. 91–95. doi: 10.1016/j.enzmictec.2018.06.009.

Rajeshkumar, S., Kumar, S. V., *et al.* (2018) 'Biosynthesis of zinc oxide nanoparticles using *Mangifera indica* leaves and evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells', *Enzyme and microbial technology*, 117, pp. 91–95.

Raj R, K., D, E. and S, R. (2020) ' β -Sitosterol-assisted silver nanoparticles activates Nrf2 and triggers mitochondrial apoptosis via oxidative stress in human hepatocellular cancer cell line', *Journal of biomedical materials research. Part A*, 108(9), pp. 1899–1908.

Ramesh, A. *et al.* (2018a) 'Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study', *Journal of Periodontology*, pp. 1241–1248. doi: 10.1002/jper.17-0445.

Ramesh, A. *et al.* (2018b) 'Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study', *Journal of periodontology*, 89(10), pp. 1241–1248.

R, K. R. *et al.* (2020) ' β - Sitosterol - assisted silver nanoparticles activates Nrf2 and triggers mitochondrial apoptosis via oxidative stress in human hepatocellular cancer cell line', *Journal of Biomedical Materials Research Part A*, pp. 1899–1908. doi: 10.1002/jbm.a.36953.

Santhoshkumar, J. *et al.* (2019) 'Toxicology evaluation and antidermatophytic activity of silver nanoparticles synthesized using leaf extract of *Passiflora caerulea*', *South African Journal of Chemical Engineering*, 29, pp. 17–23.

Saravanan, M. *et al.* (2018a) 'Synthesis of silver nanoparticles from *Phenerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria', *Microbial Pathogenesis*, pp. 68–72. doi: 10.1016/j.micpath.2018.02.008.

Saravanan, M. *et al.* (2018b) 'Synthesis of silver nanoparticles from *Phenerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria', *Microbial pathogenesis*, 117, pp. 68–72.

Vairavel, M., Devaraj, E. and Shanmugam, R. (2020a) 'An eco-friendly synthesis of *Enterococcus* sp.-mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells', *Environmental Science and Pollution Research*, pp. 8166–8175. doi: 10.1007/s11356-019-07511-x.

Vairavel, M., Devaraj, E. and Shanmugam, R. (2020b) 'An eco-friendly synthesis of *Enterococcus* sp.-mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells', *Environmental Science and Pollution Research*, 27(8), pp. 8166–8175.

Veerasamy, R. *et al.* (2021) 'Structure–Activity Relationship Analysis of Benzimidazoles as Emerging Anti-Inflammatory Agents: An Overview', *Pharmaceuticals*, 14(7), p. 663.

UNDER PEER REVIEW