

### **A SYSTEMATIC REVIEW: THE RELATIONSHIP BETWEEN FIXED PROSTHODONTICS AND DENTAL SENSITIVITY.**

---

#### **ABSTRACT:**

Dental prosthetics that are permanently attached (fixed) to the teeth are called fixed prosthetics. Crowns, bridges (fixed dentures), onlays, inlays, and veneers are also referred to as indirect restorations. Professionals in prosthodontics have trained in this area from academic institutions. With fixed prosthodontics, a single tooth or multiple teeth can be restored over areas where teeth are missing. A common oral health issue is trauma-induced tooth hypersensitivity following [the](#) preparation of the tooth. [A](#) dental hypersensitivity is caused by a sudden onset of sharp pain on contact with exposed dentin, usually in response to heat, evaporation, pressure, or cold and is not attributed to any other dental problem or condition." Despite a lack of literature on the subject, post-cementation sensitivity has not been definitively studied. When dentinal tubules are separated during [the](#) preparation of teeth, a certain degree of pulpal trauma will result. An important decision is whether to carry out elective endodontic treatment for the tooth's vital abutments or [to](#) proceed with the pulp preservation procedure.

**Keywords:** Fixed prosthodontics, prosthesis, dentin, tooth hyper-sensitivity, dental restoration.

## INTRODUCTION:

The specialty of prosthodontics is most commonly associated with replacing ~~the~~ missing teeth. There are generally two types of prosthodontics: fixed and removable. Removable prosthodontics applies to treatments involving prosthetic devices that can be removed from the mouth for cleaning and maintenance. Fixed prosthodontics includes all kinds of treatment and procedures in which replacement teeth are permanently installed in the mouth [1].

In the United States, fixed prosthodontics has become a major element of general dentistry. Crowns and fixed prostheses offer predictability and reliability, and most dentists can place them with relative ease. It is often said that a crown or short-span fixed prosthesis will last for many years; so that the expense will be worth it [2].

Direct prosthetic restorations offer two major advantages over fixed prosthetic restorations: superior strength and aesthetic appearance when ~~the~~ large restorations are used. Besides the amount and type of an appropriate dental restoration material, it is important to consider the extent of destruction of an adjacent tooth, its position and orientation, and which teeth are adjacent to the one being restored.

Generally, dental hypersensitivity is found in canine and premolar teeth, at sites that are susceptible also to gingival recession. Sometimes, dentin can be irritated, but daily activities like drinking iced tea do not ~~tend to~~ cause pain. It is more accurate to refer to dental hypersensitivity as sensitive dentin ~~rather~~ than dentin sensitivity [3].

Dentine hypersensitivity is subject to debate. In an attempt to explain dentine hypersensitivity, several theories have been proposed, including the "odontoblastic transduction theory", "neural theory", and "hydrodynamic theory". According to Brannstrom, the model most widely accepted is the hydrodynamic or fluid movement theory of 1964 [4]. This theory predicts that when the exposed dentine surface is exposed to temperatures, chemicals, tactiles or evaporative forces, the flow of fluid within the tubules increases [5].

Preparation of the pulp is associated with pulpal damage depending on the amount of remaining dentin, ~~the~~ density of that dentin, methods used to construct provisional crowns, quality of the

adhesives used for temporary and final cementation and degree of marginal penetration [6]. ~~Number-~~The number and diameter of exposed and open dentin tubules strongly influence sensitivity. In spite of the fact that dentin hypersensitivity is a common complication in dentistry, it is rarely documented, and few epidemiological data are available [7].

A dentist's role in prosthodontics includes all phases of prosthesis fabrication. As part of this, all phases of preparation, fabrication, insertion, and even follow up are included, despite the fact that manufacturers of the prosthetics may not be able to handle certain phases [8]. It is common for dentists to neglect even the most basic principles of prosthetic construction, such as making a master cast and sending it to a lab for fabrication of removable prostheses without preparing the mouth beforehand or explaining the job to them appropriately [9].

A few studies reported the evidence that some risk factors related to postoperative sensitivity occurs in three stages of the dental prosthesis process; however, those findings do not provide conclusive information regarding its origin or prevention. Therefore, it is suggested that review studies be performed ~~that could help us in understanding~~go help us understand the relationship between fixed prosthodontics and dental sensitivity.

### **AIM AND OBJECTIVES:**

The main aim of this study is to review the literature related to the dental sensitivity associated with fixed prosthodontics and understand the relationship between them.

#### **Specific objectives:**

1. To study ~~about the~~ fixed prosthodontics and the different types available.
2. Determining the significance of fixed prosthodontics in treating dental problems.
3. Carrying the online literature review using ~~the different databases to know~~various databases to-learn about dental hypersensitivity.
4. Performing a systematic review of all the articles understanding the relationship between fixed prosthodontics and dental sensitivity.

### **METHODOLOGY:**

An online based, web search was made using various databases like PubMed, Scopus, Google Scholar, Medline, and Cochrane electronic databases. The A systematic review of literature related to the relationship between fixed prosthodontics and dental sensitivity was carried out in the English language. Initially no filters were applied to the search, and all the studies were available for successive screening. On further research, only adult studies and only human studies filters were added. The research was done in a time range of 6 months to study and review all the literature.

### **DENTAL HYPERSENSITIVITY AND FIXED PROSTHODONTICS:**

In order to benefit patients, prosthetic treatments must provide comfort, function, and health. It is vital not to damage the periodontal structure. Prior to prosthodontic treatment, it is crucial that the dental tissues are strong and the dental sensitivity treatment may be required [8]. When periodontists and prosthodontists work together, the tooth or teeth that are being repaired can be cleaned in order to ensure that the prosthesis will last for as long as possible. Prosthetic outcomes are also dependent on healthy dentin. Periodontal treatment to reduce sensitivity may be required to promote this process [9].

Sensitivity to fixed dental prosthetics is caused by exposed dentinal tubules after the prosthetic is placed. The permeability of dentin increases with deeper preparations and more exposed surface area [10]. Dental sensitivity has to be prevented by a number of several methods. A technique called "immediate dentin sealing" (IDS) involves applying an adhesive system immediately to the dentin and photo polymerizing it, before an impression is taken. In this way, the dentinal tubules are closed, and bacterial invasions and dentin circulation are stopped, resulting in pain. It has been reported that this procedure decreases infection risk and reduces postoperative sensitivity and adhesion strength [11].

There is evidence that provisionalization increases pulp chamber temperature, which may be the main cause of postoperative sensitivity. It has been demonstrated<sup>24-25</sup> that polymethylmethacrylate (PMMA), polyethylmethacrylate (PEMA), polyvinyl methacrylate (PVMA), and Bisacryl resin cause temperature increases. In research studies, self-curing sets have demonstrated a lower thermal change than dual curing sets, and the latter results in a lower temperature rise [12].

A dentin sensitivity is caused by acid etching because it removes the smear layer, which allows bacteria to infiltrate the dentinal tubules. The exposure of collagen by mineralized dentin and resin infiltration can also lead to dentin sensitization. Therefore, self-etching cements should be used in order to avoid removing the smear layer [13]. The pH value affects dental sensitivity as well. In comparison to resin cements, glass ionomers and zinc phosphate have an inverse correlation between pH value and the compound. In Lam and Wilson's findings, the removal of the smear layer can lead to significant increases in pressure transmission to the pulp chamber. Immediate dentin sealing (IDS) is a technique for creating a barrier after adhering the smear layer to the dentin.

The mechanism ~~that is~~ involved in ~~the~~ dentin hypersensitivity is explained by several theories like: Neural theory, Odontoblastic transduction theory, hydrodynamic theory. In accordance with neural theory, the connection between pulpal nerve endings and dentine tubules may cause direct stimulation of nerve endings in dentine tubules. A theory related to odontoblast transduction states that odontoblasts function as receptor cells, transmitting synaptic impulses to nerve terminals that produce a sensation of pain due to nerve endings located at the pulpodentine border. The hydrodynamic theory, proposed by Brännström in 1964, is currently accepted as the explanation for dentine hypersensitivity [14]. Theoretically, when the surface of the exposed dentin is exposed to thermal, chemical, tactile or evaporative stimuli, then the flow of fluid within the tubules of the dentine will increase. Fluid movement affects the pressure in dentine tubules, which in turn activates pressure-sensitive nerve receptors. The response of excited pulpal nerves, especially those in the intradentine fibers, will be determined by the intensity of pain stimuli. Exposed dentine in all cases is not sensitive. Dentine Hypersensitivity is the result of lesion localization. Upon the removal of the protective covering of the smear layer, the dentinal tubules become exposed.

The composition of provisional restorations, preparation under high volume spray and tooth reduction were considered to have significant impacts on the incidence of dentin sensitivity. In order to reduce postoperative sensitivity, several methods have been used, including the choice of operative technique and a plentiful use of water cooling before and after tooth extraction [12].

In order to achieve optimal seal and adhesion, dentin surfaces must be thoroughly cleaned. Clean and freshly cut dentin is more likely to be susceptible to resin infiltration. Dentin is protected by

immediate sealing from contamination caused by bacterial leakage or remnants of expired temporary cements [15]. It is possible to eliminate ~~the concern for~~ gap formation and ill-fitting restorations by including the hybrid layer in an impression. A new approach to dentin sealing is called immediate dentin sealing (IDS), in which the dentin is sealed immediately after preparation of the tooth and prior to impression taking. In scanning electron microscopy, the continuity between the hybrid layer and dentin appeared ~~to be~~ more evident when the dentin was sealed with a dentin bonding agent.

To reduce the risk of vital abutment sensitivity, the exposed dentin can be desensitized with desensitizing agents after tooth preparation and before restoration is cemented. Desensitizing agents have been found to reduce dentin sensitivities when ~~they are~~ used on vital abutment teeth to receive full coverage restorations. Desensitizers occlude the dentinal tubules on the surface and under the surface, therefore decreasing pain by counteracting the hydrodynamic mechanism of dentinal hypersensitivity. An important decision is whether to carry out elective endodontic treatment for the tooth's vital abutments or ~~to~~ proceed with the pulp preservation procedure. When dentinal tubules are separated during the preparation of teeth, a certain degree of pulpal trauma will result.

## **CONCLUSION:**

Occasionally, ~~an~~ increased sensitivity to hot or cold stimulation can be a perplexing result of a newly cemented crown or fixed partial denture. There is no way to completely avoid sensitivity. Despite a lack of literature on the subject, post-cementation sensitivity has not been definitively studied. Management of post cementation hypersensitivity can be improved by better understanding the causes and precautions.

## **ETHICAL CONSIDERATIONS:**

### **Compliance with ethical standards**

Ethical approval: This is a systematic review about of the relationship between the fixed prosthodontics and dental sensitivity.

#### REFERENCES:

1. Chandrasekhar V. Post cementation sensitivity evaluation of glass Ionomer, zinc phosphate and resin modified glass Ionomer luting cements under class II inlays: An in vivo comparative study. *Journal of conservative dentistry: JCD*. 2010 Jan; 13(1):23.
2. Lam CW, Wilson PR. Crown cementation and pulpal health. *International Endodontic Journal*. 1999 Jul;32(4):249-56.
3. Mullaney TP, Laswell HR. Iatrogenic blushing of dentin following full crown preparation. *The Journal of prosthetic dentistry*. 1969 Sep 1; 22(3):354-9.
4. Brännström M. Reducing the risk of sensitivity and pulpal complications after the placement of crowns and fixed partial dentures. *Quintessence International*. 1996 Oct 1; 27(10)
5. Sivaramakrishnan G, Sridharan K. Fluoride varnish versus glutaraldehyde for hypersensitive teeth: a randomized controlled trial, meta-analysis and trial sequential analysis. *Clin Oral Investig*. 2019 Jan; 23(1):209-220.
6. Murray LE, Roberts AJ. The prevalence of self-reported hypersensitive teeth. *Arch Oral Biol*. 1994;39:1295.
7. West NX. Dentine hypersensitivity: Preventive and therapeutic approaches to treatment. *Periodontol 2000*. 2008;48:31–41.
8. France CR. Decreased pain perception and risk for hypertension: Considering a common physiological mechanism. *Psychophysiology*. 1999;36:683–92.
9. Gijssbers K, Nicholson F. Experimental pain thresholds influenced by sex of experimenter. *Percept Mot Skills*. 2005;101:803–7.
10. Holland GR, Narhi MN, Addy M, Gangarosa L, Orchardson R. Guidelines for the design and conduct of clinical trials on dentine hypersensitivity. *J Clin Periodontol*. 1997;24:808–13.
11. Chun JS, Har A, Lim HP, Lim HJ. The analysis of cost-effectiveness of implant and conventional fixed dental prosthesis. *J Adv Prosthodont*. 2016 Feb;8(1):53-61.

12. McGuire MK, Nunn ME. Prognosis versus actual outcome. II. The effectiveness of clinical parameters in developing an accurate prognosis. *J Periodontol.* 1996 Jul;67(7):658-65.
13. Korenori A, Koji K, Yuki T, Murata T, Sachiko TM, Shunsuke B. Cost-effectiveness of molar single-implant versus fixed dental prosthesis. *BMC Oral Health.* 2018 Aug 20;18(1):141.
14. Brännström M. Reducing the risk of sensitivity and pulpal complications after the placement of crowns and fixed partial dentures. *Quintessence International.* 1996 Oct 1; 27(10).
15. Gupta N, Reddy UN, Vasundhar PL, Ramarao KS, Varma KP, Vinod V. Effectiveness of desensitizing agents in relieving the pre-and post-cementation sensitivity for full coverage restorations: A clinical evaluation. *J. Contemp. Dent. Pract.* 2013 Sep 1;14:858-65.