

QUANTITATIVE ANALYSIS OF **TANNERELLA FORSYTHIA** LEVEL IN PERIODONTITIS PATIENTS WITH OR WITHOUT DIABETES **MELLITUS**

ABSTRACT

INTRODUCTION- Periodontitis is a systemic chronic inflammatory disease resulting from a complex polymicrobial infection causing tissue destruction as a consequence of perturbation of the homeostasis between sub gingival microflora. **T.forsythia** has been strongly implicated in the onset of periodontitis. The level of glycemic control is the key to determining increased risk. In adults, **HbA1C** levels >9% have a higher prevalence rate for periodontitis than those without diabetes. In adults, random blood sugar levels >200mg/dl have higher susceptibility to periodontitis.

MATERIALS AND METHOD- This study was carried out in a university setting at saveetha dental college. Total of 8 samples were collected from November 2020 to February 2021. The subgingival plaque samples containing bacterial cells were pelleted by centrifugation at room temperature. Following centrifugation with all buffer solutions, quantitative RT-PCR was performed with stratagene MX3000P and the relative gene count was calculated using relative CT method. The purified DNA molecules were identified by agarose gel electrophoresis and **Tanerella forsythia** was identified using this method.

RESULTS- The assessment of mRNA expression of T.forsythia assessed by RT-PCR showed that **T.forsythia** was seen with more prevalence in patients with periodontitis with diabetes mellitus (n=1.388), than in patients with periodontitis without diabetes mellitus (n=1) (P=0.0028 < 0.005). The study was statistically significant.

CONCLUSION- This study shows that **Tannerella forsythia** is seen with more prevalence in patients with periodontitis with diabetes mellitus when compared to periodontitis without diabetes mellitus.

Keywords: Diabetes mellitus, Innovative technology, **Tannerella.forsythia**, periodontitis.

1. **INTRODUCTION**

Periodontitis is a systemic chronic inflammatory disease resulting from a complex polymicrobial infection causing tissue destruction as a consequence of perturbation of the homeostasis between sub gingival microflora.[1] It can lead to root caries, eating disabilities, tooth mobility and systemic diseases in the human population[2][3]. The major microflora causing periodontitis in the human community belong to the red complex species which includes *A.actinomycetemcomitans*, *P.gingivalis*, *T. forsythia*, *P.nigrescens*, *P.intermedia*, *T.denticola*.[4]

T.forsythia has been strongly implicated in the onset of periodontitis.[5,6] *T.forsythia* is a anaerobic, gram negative bacteria belonging to the cytophaga-bacteroides family which was initially described as **Bacteroides forsythus**. [7,8] Diabetes mellitus increases the susceptibility to periodontitis by a threefold.[9–11] The level of glycemic control is the key to determining increased risk].In adults, HbA1C levels >9% have a higher prevalence rate for periodontitis than those without diabetes.[14,15] In adults, random

Comment [U1]: Microorganisms in cursive letter

Before HbA1C include the mean

Comment [U2]: Saaveetha S mayor

Comment [U3]: Without 1.

Comment [U4]: Gram

Comment [U5]: Cursive letter

blood sugar levels >200mg/dl have higher susceptibility to periodontitis.[16,17].The present study aims to quantitatively analyse *T.forsythia* levels in periodontitis patients with or without diabetes mellitus.

2. MATERIALS AND METHOD

This study was carried out in a university setting at Saveetha dental college.

Sample collection: Patients aged between 30-60 reported to the department of periodontics. Total of 8 samples were collected from November 2020 to February 2021. The samples were selected from patients diagnosed with generalised periodontitis with a history or without diabetes mellitus.

Inclusion criteria:

1. No >2 missing teeth per quadrant
2. >30% of sites with PD greater than or equal to 4mm
3. >20% of sites with CAL greater than 2mm
4. Bleed on probing greater than or equal to 30% and radiographic indication of bone loss.

Exclusion criteria:

1. If the patient had undergone and periodontal treatment in the last 6 months
2. Tobacco use and smoking
3. Alcoholism
4. Conditions that could alter healing response like periodontal abscess, acute necrotising gingivitis, oral mucosal lesions

Sample collection:

Supragingival and subgingival plaque samples were collected for 8 patients of the respective group with the help of curettes

Isolation and Quantification of *T.forsythia* by Real time-PCR:

The subgingival plaque samples containing bacterial cells were pelleted by centrifugation at room temperature. Following centrifugation with all buffer solutions, quantitative RT-PCR was performed with stratagene MX3000P and the relative gene count was calculated using relative CT method. The purified DNA molecules were identified by agarose gel electrophoresis and *Tanerella forsythia* was identified using this method.

3. RESULTS

The assessment of mRNA expression of *T.forsythia* assessed by RT-PCR showed that *T.forsythia* was seen with more prevalence in patients with periodontitis with diabetes mellitus (n=1.388), than in patients with periodontitis without diabetes mellitus (n=1) (P=0.0028 < 0.005). The study was statistically significant.

Comment [U6]: Methodology just omit the number 2

Comment [U7]: Saveetha

Comment [U8]: generalized

Comment [U9]: Cursive letter




Fig1- This graph shows the expression of mRNA of *T.forsythia* in patients with periodontitis without diabetes mellitus and patients with periodontitis with diabetes mellitus.

Comment [U10]: I suggest to avoid the title of the graph up. Only in the bottom

group	periodontitis	periodontitis +DM	P VALUE
<i>T.forsythia</i> mRNA	1 +/- 0.00	1.350 +/- 0.057	0.0067

Comment [U11]: Cursive

4.DISCUSSION

Tannerella forsythia is a Gram-negative anaerobic organism that inhabits the subgingival cavity and initiates connective tissue destruction and alveolar bone resorption in periodontal disease (PD).[18,19].

This study showed that *T.forsythia* was seen with more prevalence in patients with periodontitis with diabetes mellitus.[20,21] Investigations have demonstrated associations between periodontitis and various systemic diseases such as cardiovascular disorders, respiratory diseases, osteoporosis, immunodeficiencies and also diabetes mellitus.[22,23] Longitudinally studies have demonstrated a two-way relationship between diabetes and periodontitis[24], with more severe periodontal tissue destruction in diabetic patients and poorer glycemic control in diabetic subjects with periodontal disease. [14]

Glycemic control was the most important risk factor related to severity and extent of periodontitis.[25,26] Diabetes leads to worsening of periodontal disease, and a significant association between diabetes and periodontitis has been demonstrated.[27] Periodontal disease has a higher incidence in diabetic patients, and it is more prevalent and severe if compared with a healthy population. The risk of periodontitis is 3-fold times higher among diabetic patients, being its prevalence and severity even greater in diabetic patients presenting elevated HbA1c levels.[25,28] Periodontal inflammation, as any other infections, can have an adverse effect on diabetes glycemic control, compromising diabetes management in these individuals.[16] Most evidence on this issue is derived from interventional and observational studies, indicating that periodontitis affects the glycemic control of diabetic patients.[29] HbA1c values < 7% are

Comment [U12]: Cursive

Comment [U13]: Include a dot.

related with proper glycemic levels whilst > 8% values represents poorly controlled glycemia.[12] Longitudinal studies have demonstrated that severe periodontitis is associated with poorly controlled glycemia, higher HbA1c levels and development of diabetic systemic complications.

5.CONCLUSION

This study shows that *T. forsythia* is seen with more prevalence in patients with periodontitis with diabetes mellitus when compared to periodontitis without diabetes mellitus.

Comment [U14]: If is possible include more conclusions found in the research

Comment [U15]: A section of Ethical aspects as Conflict of interest, informed consent and also a section of Acknowledgments

REFERENCES

1. Bahekar AA, Singh S, Saha S, Molnar J, Arora R. The prevalence and incidence of coronary heart disease is significantly increased in periodontitis: a meta-analysis. *Am Heart J* [Internet]. 2007 Nov;154(5):830–7. Available from: <http://dx.doi.org/10.1016/j.ahj.2007.06.037>
2. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol* [Internet]. 2018 Oct;89(10):1241–8. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/JPER.17-0445>
3. Del Fabbro M, Karanxha L, Panda S, Bucchi C, Nadathur Doraiswamy J, Sankari M, et al. Autologous platelet concentrates for treating periodontal infrabony defects. *Cochrane Database Syst Rev* [Internet]. 2018 Nov 26;11:CD011423. Available from: <http://dx.doi.org/10.1002/14651858.CD011423.pub2>
4. Borody TJ, Khoruts A. Fecal microbiota transplantation and emerging applications. *Nat Rev Gastroenterol Hepatol* [Internet]. 2011 Dec 20;9(2):88–96. Available from: <http://dx.doi.org/10.1038/nrgastro.2011.244>
5. Cossart P, Sansonetti PJ. Bacterial invasion: the paradigms of enteroinvasive pathogens. *Science* [Internet]. 2004 Apr 9;304(5668):242–8. Available from: <http://dx.doi.org/10.1126/science.1090124>
6. Paramasivam A, Priyadharsini JV, Raghunandhakumar S, Elumalai P. A novel COVID-19 and its effects on cardiovascular disease [Internet]. Vol. 43, *Hypertension Research*. 2020. p. 729–30. Available from: <http://dx.doi.org/10.1038/s41440-020-0461-x>
7. S. G, T. G, K. V, A. AF, Sukumaran A, Sudha PN. Development of 3D scaffolds using nanochitosan/silk-fibroin/hyaluronic acid biomaterials for tissue engineering applications [Internet]. Vol. 120, *International Journal of Biological Macromolecules*. 2018. p. 876–85. Available from: <http://dx.doi.org/10.1016/j.ijbiomac.2018.08.149>
8. Paramasivam A, Vijayashree Priyadharsini J. MitomiRs: new emerging microRNAs in mitochondrial dysfunction and cardiovascular disease. *Hypertens Res* [Internet]. 2020 Aug;43(8):851–3. Available from: <http://dx.doi.org/10.1038/s41440-020-0423-3>
9. Salvi GE, Carollo-Bittel B, Lang NP. Effects of diabetes mellitus on periodontal and peri-implant conditions: update on associations and risks [Internet]. Vol. 35, *Journal of Clinical Periodontology*. 2008. p. 398–409. Available from: <http://dx.doi.org/10.1111/j.1600-051x.2008.01282.x>
10. Vellappally S, Al Kheraif AA, Anil S, Assery MK, Kumar KA, Divakar DD. Analyzing Relationship between Patient and Doctor in Public Dental Health using Particle Memetic Multivariable Logistic Regression Analysis Approach (MLRA2). *J Med Syst* [Internet]. 2018 Aug 29;42(10):183. Available

from: <http://dx.doi.org/10.1007/s10916-018-1037-z>

11. Varghese SS, Ramesh A, Veeraiyan DN. Blended Module-Based Teaching in Biostatistics and Research Methodology: A Retrospective Study with Postgraduate Dental Students [Internet]. Vol. 83, Journal of Dental Education. 2019. p. 445–50. Available from: <http://dx.doi.org/10.21815/jde.019.054>
12. Chávary NGM, Vettore MV, Sansone C, Sheiham A. The relationship between diabetes mellitus and destructive periodontal disease: a meta-analysis. Oral Health Prev Dent [Internet]. 2009;7(2):107–27. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19583037>
13. Venkatesan J, Singh SK, Anil S, Kim S-K, Shim MS. Preparation, Characterization and Biological Applications of Biosynthesized Silver Nanoparticles with Chitosan-Fucoidan Coating. Molecules [Internet]. 2018 Jun 12;23(6). Available from: <http://dx.doi.org/10.3390/molecules23061429>
14. Mealey BL, Ocampo GL. Diabetes mellitus and periodontal disease. Periodontol 2000 [Internet]. 2007;44:127–53. Available from: <http://dx.doi.org/10.1111/j.1600-0757.2006.00193.x>
15. Alsubait SA, Al Ajlan R, Mitwalli H, Aburaisi N, Mahmood A, Muthurangan M, et al. Cytotoxicity of Different Concentrations of Three Root Canal Sealers on Human Mesenchymal Stem Cells. Biomolecules [Internet]. 2018 Aug 1;8(3). Available from: <http://dx.doi.org/10.3390/biom8030068>
16. Tsai C, Hayes C, Taylor GW. Glycemic control of type 2 diabetes and severe periodontal disease in the US adult population. Community Dent Oral Epidemiol [Internet]. 2002 Jun;30(3):182–92. Available from: <http://dx.doi.org/10.1034/j.1600-0528.2002.300304.x>
17. Vellappally S, Al Kheraif AA, Anil S, Wahba AA. IoT medical tooth mounted sensor for monitoring teeth and food level using bacterial optimization along with adaptive deep learning neural network. Measurement [Internet]. 2019 Mar 1;135:672–7. Available from: <https://www.sciencedirect.com/science/article/pii/S0263224118311333>
18. Holt SC, Ebersole JL. Porphyromonas gingivalis, Treponema denticola, and Tannerella forsythia: the “red complex”, a prototype polybacterial pathogenic consortium in periodontitis. Periodontol 2000 [Internet]. 2005;38(1):72–122. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/j.1600-0757.2005.00113.x>
19. Venkatesan J, Rekha PD, Anil S, Bhatnagar I, Sudha PN, Dechsakulwatana C, et al. Hydroxyapatite from Cuttlefish Bone: Isolation, Characterizations, and Applications. Biotechnol Bioprocess Eng [Internet]. 2018 Aug 1;23(4):383–93. Available from: <https://doi.org/10.1007/s12257-018-0169-9>
20. R H, Ramani P, Tilakaratne WM, Sukumaran G, Ramasubramanian A, Krishnan RP. Critical appraisal of different triggering pathways for the pathobiology of pemphigus vulgaris-A review. Oral Dis [Internet]. 2021 Jun 21; Available from: <http://dx.doi.org/10.1111/odi.13937>
21. Ezhilarasan D, Lakshmi T, Subha M, Deepak Nallasamy V, Raghunandhakumar S. The ambiguous role of sirtuins in head and neck squamous cell carcinoma. Oral Dis [Internet]. 2021 Feb 11; Available from: <http://dx.doi.org/10.1111/odi.13798>
22. Bodet C, Chandad F, Grenier D. Potentiel pathogénique de Porphyromonas gingivalis, Treponema denticola et Tannerella forsythia, le complexe bactérien rouge associé à la parodontite. Pathol Biol [Internet]. 2007 Apr;55(3-4):154–62. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0369811406001751>
23. PradeepKumar AR, Shemesh H, Nivedhitha MS, Hashir MMJ, Arockiam S, Uma Maheswari TN, et al. Diagnosis of Vertical Root Fractures by Cone-beam Computed Tomography in Root-filled Teeth

with Confirmation by Direct Visualization: A Systematic Review and Meta-Analysis. *J Endod* [Internet]. 2021 Aug;47(8):1198–214. Available from: <http://dx.doi.org/10.1016/j.joen.2021.04.022>

24. Kavarthapu A, Gurumoorthy K. Linking chronic periodontitis and oral cancer: A review. *Oral Oncol* [Internet]. 2021 Jun 14;105375. Available from: <http://dx.doi.org/10.1016/j.oraloncology.2021.105375>
25. Emrich LJ, Shlossman M, Genco RJ. Periodontal disease in non-insulin-dependent diabetes mellitus. *J Periodontol* [Internet]. 1991 Feb;62(2):123–31. Available from: <http://dx.doi.org/10.1902/jop.1991.62.2.123>
26. Sarode SC, Gondivkar S, Sarode GS, Gadail A, Yuwanati M. Hybrid oral potentially malignant disorder: A neglected fact in oral submucous fibrosis. *Oral Oncol* [Internet]. 2021 Jun 16;105390. Available from: <http://dx.doi.org/10.1016/j.oraloncology.2021.105390>
27. Cianciola LJ, Park BH, Bruck E, Mosovich L, Genco RJ. Prevalence of periodontal disease in insulin-dependent diabetes mellitus (juvenile diabetes). *J Am Dent Assoc* [Internet]. 1982 May;104(5):653–60. Available from: <http://dx.doi.org/10.14219/jada.archive.1982.0240>
28. Vellappally S, Abdullah Al-Kheraif A, Anil S, Basavarajappa S, Hassanein AS. Maintaining patient oral health by using a xeno-genetic spiking neural network. *J Ambient Intell Humaniz Comput* [Internet]. 2018 Dec 14; Available from: <https://doi.org/10.1007/s12652-018-1166-8>
29. Aldhuwayhi S, Mallineni SK, Sakhamuri S, Thakare AA, Mallineni S, Sajja R, et al. Covid-19 Knowledge and Perceptions Among Dental Specialists: A Cross-Sectional Online Questionnaire Survey. *Risk Manag Healthc Policy* [Internet]. 2021 Jul 7;14:2851–61. Available from: <http://dx.doi.org/10.2147/RMHP.S306880>