

## Original Research Article

### **Gonial angle as a determinant of gender - a panoramic study among South Indian Population**

#### **ABSTRACT:**

**INTRODUCTION:** Sex determination is a valuable and important factor in the forensic dentistry. Mandible has several useful traits for sex determination. Forensic practitioners study this by two methods: morphological and metrical analyses. Among various methods, the gonial angle may be used to differentiate male and female strongly to express sexual dimorphism.

**AIM:** To analyse the gender determination using gonial angle as panoramic study among south indian population.

**MATERIALS AND METHODS:** This is a retrospective study conducted among 200 individuals. Gonial angle is a measurement taken by measuring the angle between the 2 tangents from the gonion. As a standard procedure, measurement of right gonial angle and left gonial angle were carried out by Adobe Photoshop and the results were recorded and analysed statistically using SPSS tool.

**RESULT:** The mean right gonial angle for males is 95.25 and for females is 95.22. The mean left gonial angle for males is 95.39 and for females is 95.15. The significance of the right gonial angle is  $P=0.874$  and significance of the left gonial angle is  $P=0.147$ . The total mean value for right gonial angle was found to be 95.23 and for left gonial angle is 95.27. Therefore, there was no significant difference found between the right and left gonial angle.

**CONCLUSION:** Within the limitations of the study, there is no statistically significant difference observed between males and females in the gonial angle measurements. The Gonial angle measurements can be used to determine the gender and the growth pattern of the mandible. Measurement of gonial angle has its own surgical importance too in the field of orthodontics and therefore future studies should be carried out using larger samples to get reliable results.

**KEYWORD:** Gonial angle; sex determination; sexual dimorphism; mandible; Innovative technique.

**Running title:** Gonial angle and gender determination

## INTRODUCTION:

Sex determination is a valuable and important factor in the forensic dentistry. Mandible has several useful traits for sex determination. Forensic practitioners study this by two methods: morphological and metrical analyses. Other than the mandibular measurements, pelvis plays an important role in determination of sex as it is the most sexually dimorphic region. Apart from gender determination, the gonial angle in Orthopantomogram is an important parameter for determining the growth pattern of an individual. It provides a significant amount of information about the dentition and the supporting bone(1).

Gonial angle can be used to differentiate male and female to express sexual dimorphism. The approximate measurement of the gonial angle reported in males is 100 degree to 148 degree whereas in females it is 103 degree to 153 degree with development and function, the mandibular angle has shown some changes in size, shape and length(2). The mandible is the largest and strongest bone present in the face. The occurrence of Sexual dimorphism in the mandible may be due to differences in development of the musculoskeletal system (2,3).the differences between ramus of the mandible among gender is appreciated based on the different stages of development and growth rates between male and female(2–4). Assessment of gonial angle for sex determination according to age and gender are evidenced in certain studies(2–5). The experience from our previous studies conducted among different fields (6) (7,8) (7)(9)(10)(11)(12)(10,12)(13)(14) (15) have led us to focus on the current topic.Measurement of gonial angle is essential for the treatment and surgery in orthodontics. Therefore, accurate determination of the gonial angle is important for assessing the orthodontic cases(16).

To evaluate the morphology of the mandible gonial angle ramus height and gonial width are measured. A wider gonial angle is found in edentulous individuals when compared to dentulous individuals. These factors are correlated with the function and architecture of the muscle of mastication. Aging causes changes in masticatory function and changes the contractile activity of individuals. Our team has extensive knowledge and research experience that has translated into high quality publications (17–24),(25),(26),(27),(28,29),(30),(31),(32–36). The aim of the present study is to assess the gender determination using gonial angle in OPGs of south indian population.

Comment [is1]: Too long paragraph

Comment [is2]: Citation number should be corrected


Comment [is3]: Citation number should be corrected

**MATERIALS AND METHODS:** This was a retrospective study conducted in a private dental college and hospital in Chennai. The study was approved by the institutional review board. There were 200 randomly selected OPGs of population aged between 5-60 years that were used as samples. The measurement of the gonial angle was taken by Adobe Photoshop. Gonial angle measurements were taken by measuring the angle between the 2 tangents from the gonion. Panoramic radiographs with intact dentition were included for the study. Panoramic radiographs with fracture or deformation were excluded. The collected data was tabulated and analysed using SPSS software. The mean value was calculated and the comparison analysis carried for gender using independent t-test.

**Comment [is4]:** More detail about the material and method


**Comment [is5]:** Title should be modified 200 can't represent whole south Indian population

**RESULT:** The mean value of gonial angle in the right side for males is 95.25 and for females is 95.22 (fig-1). The mean value of gonial angle in the left side for males is 95.35 and for females is 95.15 (fig-2). The mean value of left gonial angle is not statistically significant  $p=0.147$  ( $p>0.05$ ) and the mean value of right gonial angle is not statistically significant  $p=0.874$  ( $p>0.05$ ). The total mean value for right gonial angle was found to be 95.23 (fig-1) and for left gonial angle is 95.27 (fig-2). Therefore, there was no significant difference found between the right gonial angle and left gonial angle.



**Figure 1:** Bar graph shows the comparison between gender. The X-axis represents the gender and mean right gonial angle. Y-axis represents the mean right gonial angle measurement. The red colour represents male and the green colour represents female. There was statistically no significant difference observed between the genders with p value>0.05.

UNDER  
REVIEW



**Figure 2:** This graph shows the comparison between gender and mean left gonial angle. The X-axis represents the gender. Y-axis represents the mean left gonial angle measurement. The red colour represents male and the green colour represents female. There was statistically no significant difference observed between genders with  $p \text{ value} > 0.05$ .

## **DISCUSSION:**

From the results of the current study, it was observed that the mean value of both right and left gonial angle has no significant difference between male and female. In this context, we studied how gonial angle helps to determine sex. Gonial angle has been used as a forensic parameter. A variation in mandibular angle with age and gender can be observed. Evidence reported the relationship between complete loss of teeth and changes in gonial angle and intends to evaluate variation in gonial angle with gender(37), .

In case of females under 10 years of age, the gonial angle values are observed to be decreased. There were no significant gender differences in development and function, and the mandible angle has shown some changes in size and shape, which is supported by radiographs and anthropometric studies(38) .Gender determination can help establish a biological profile of the human body. All the linear measurements such as facial height, mandibular ramus height, mandibular plane, frontal sinus width were significantly larger in males except for angular variation which showed no significant differences between the two genders which is in agreement with the present study. (3,38).

Gonial angle is formed from two lines, the inferior border of the mandible and the posterior border of the mandible ramus(39). A study reporting the mean of gonial angle measurement confirmed that there was no difference in terms of data ( $p>0.05$ ) which is in agreement with the present study. It has been shown that the shape of the mandible correlates with condition and function of the masticatory muscles. A study evaluated the association of tooth loss on the shape of mandibles in subjects aged 60 years. The gonial angle of the mandible and the mandibular and condylar height were measured using panoramic radiographs. The morphology of the mandible changes as a consequence of tooth loss, which can be expressed as a widening of the gonial angle and shorting of the ramus and condylar height(40). Cross sectional studies indicate an increase in the size of the gonial angle from early embryonic stages to the time of birth and a continuous decrease from birth to old age. A complete loss of teeth may reverse the usual age changes and the gonial angle becomes more obtuse again. The size of the gonial angle is associated with the proportion between facial height and ramus height. The size of the gonial angle is rather independent of variations in facial development seen in the sagittal plane(41).

Mandibular ramus can differentiate between sexes, as the stages of mandibular development, growth rates, and duration are distinctly different in both sexes. Furthermore, the masticatory forces exerted by males and females differ, which influences the shape and size of the mandibular, which is thought to be influenced by the subject's age, where longitudinal studies have shown that remodeling of the mandibular bone occurs with age. Comparison between males and females revealed that males showed statistically significant higher mean ramus linear measurements and lower mean gonial angle values than females(4) The limitations of the study is that this is a retrospective study and future studies with larger sample size should be conducted to get definite results that it can be generalized.

**CONCLUSION:** Within the limitations of the study, there is no statistically significant difference observed between males and females in the gonial angle measurements. The Gonial angle measurements can be used to determine the gender and the growth pattern of the mandible. Measurement of gonial angle has its own surgical importance too in the field of orthodontics and therefore future studies should be carried out using larger samples to get reliable results.

#### REFERENCES:

1. Alfawzan AA. Gonial Angle as a Determinant of Gender, a Panoramic Study in a Sample of Saudi Population [Internet]. Vol. 11, Indian Journal of Public Health Research & Development. 2020. p. 1689. Available from: <http://dx.doi.org/10.37506/v11/i1/2020/ijphrd/194092>
2. Upadhyay R, Upadhyay J, Agrawal P, Rao N. Analysis of gonial angle in relation to age, gender, and dentition status by radiological and anthropometric methods [Internet]. Vol. 4, Journal of Forensic Dental Sciences. 2012. p. 29. Available from: <http://dx.doi.org/10.4103/0975-1475.99160>
3. Sharma M, Gorea RK, Gorea A, Abuderman A. A morphometric study of the human mandible in the Indian population for sex determination [Internet]. Vol. 6, Egyptian Journal of Forensic Sciences. 2016. p. 165–9. Available from: <http://dx.doi.org/10.1016/j.ejfs.2015.01.002>
4. Abu-Taleb NS, El Beshlawy DM. Mandibular Ramus and Gonial Angle Measurements as Predictors of Sex and Age in an Egyptian Population Sample: A Digital Panoramic Study [Internet]. Vol. 06, Journal of Forensic Research. 2015. Available from: <http://dx.doi.org/10.4172/2157-7145.1000308>
5. Abuhijleh E, Warreth A, Qawadi M, Abdulrida E, Radaideh A, Al Taki A, et al. Mandibular Gonial Angle Measurement as a Predictor of Gender-A Digital Panoramic Study [Internet]. Vol. 13, The Open Dentistry Journal. 2019. p. 399–404. Available from: <http://dx.doi.org/10.2174/1874210601913010399>

6. Shruthi M, Preetha S. Effect of Simple Tongue Exercises in Habitual Snorers [Internet]. Vol. 11, Research Journal of Pharmacy and Technology. 2018. p. 3614. Available from: <http://dx.doi.org/10.5958/0974-360x.2018.00665.0>
7. Preetha S, Packyanathan J. Comparison of the effect of Yoga, Zumba and Aerobics in controlling blood pressure in the Indian population [Internet]. Vol. 9, Journal of Family Medicine and Primary Care. 2020. p. 547. Available from: [http://dx.doi.org/10.4103/jfmpc.jfmpc\\_607\\_19](http://dx.doi.org/10.4103/jfmpc.jfmpc_607_19)
8. J SK, Saveetha Dental College and Hospitals, Road PH, Chennai, Tamilnadu, Preetha S, et al. Effect of aerobics exercise and yoga on blood pressure in hypertensives [Internet]. Vol. 6, International Journal of Current Advanced Research. 2017. p. 3124–6. Available from: <http://dx.doi.org/10.24327/ijcar.2017.3126.0200>
9. Prathap L, Suganthirababu P, Ganesan D. Fluctuating Asymmetry of Dermatoglyphics and DNA Polymorphism in Breast Cancer Population [Internet]. Vol. 10, Indian Journal of Public Health Research & Development. 2019. p. 3574. Available from: <http://dx.doi.org/10.5958/0976-5506.2019.04141.x>
10. Lavanya J, Prathap S, Alagesan J. Digital and palmar dermal ridge patterns in population with breast carcinoma. Biomedicine. 2014 Jul 1;34(3):315–21.
11. Prathap L, Jagadeesan V. Association of quantitative and qualitative dermatoglyphic variable and DNA polymorphism in female breast cancer population. Online J Health [Internet]. 2017; Available from: [https://www.researchgate.net/profile/Prathap\\_Suganthirababu/publication/321606278\\_Association\\_of\\_Quantitative\\_and\\_Qualitative\\_Dermatoglyphic\\_Variable\\_and\\_DNA\\_Polymorphism\\_in\\_Female\\_Breast\\_Cancer\\_Population/links/5a28c8f1a6fdcc8e8671c0cd/Association-of-Quantitative-and-Qualitative-Dermatoglyphic-Variable-and-DNA-Polymorphism-in-Female-Breast-Cancer-Population.pdf](https://www.researchgate.net/profile/Prathap_Suganthirababu/publication/321606278_Association_of_Quantitative_and_Qualitative_Dermatoglyphic_Variable_and_DNA_Polymorphism_in_Female_Breast_Cancer_Population/links/5a28c8f1a6fdcc8e8671c0cd/Association-of-Quantitative-and-Qualitative-Dermatoglyphic-Variable-and-DNA-Polymorphism-in-Female-Breast-Cancer-Population.pdf)
12. Lavanya J, Kumar VJ, Sudhakar N, Prathap S. Analysis of DNA repair genetic polymorphism in breast cancer population. Int J Pharma Bio Sci [Internet]. 2015; Available from: [https://scholar.google.ca/scholar?cluster=8949053652564257518&hl=en&as\\_sdt=0,5&scio dt=0,5](https://scholar.google.ca/scholar?cluster=8949053652564257518&hl=en&as_sdt=0,5&scio dt=0,5)
13. Prathap L, Suganthirababu P. Estrogen Exposure and its Influence in DNA Repair Genetic Variants in Breast Cancer Population [Internet]. Vol. 13, Biomedical and Pharmacology Journal. 2020. p. 1321–7. Available from: <http://dx.doi.org/10.13005/bpj/2001>
14. Ravikumar H, Prathap L, Preetha S. ANALYSIS OF PALMAR ATD ANGLE IN POPULATION WITH MALOCCLUSION. 2020 Jan 1;1174–82.
15. Prathap L. INTERPLAY OF OXIDATIVE STRESS AND LIPOPROTEINS IN BREAST CARCINOMA INITIATION, PROMOTION AND PROGRESSION -A SYSTEMATIC REVIEW. PalArch's Journal of Archaeology of Egypt/ Egyptology [Internet]. 2021 Jan 7 [cited 2021 Mar 9];17(7). Available from: <http://dx.doi.org/>
16. Lim TW, Sabri BAM, Ahmad R. The effect of mandibular angulation on gonial angle and tooth length measurement of dental panoramic radiographs [Internet]. Bioresources Technology in Sustainable Agriculture. 2018. p. 167–75. Available from:

<http://dx.doi.org/10.1201/9781315365961-12>

17. Sekar D, Lakshmanan G, Mani P, Biruntha M. Methylation-dependent circulating microRNA 510 in preeclampsia patients. *Hypertens Res.* 2019 Oct;42(10):1647–8.
18. Princeton B, Santhakumar P, Prathap L. Awareness on Preventive Measures taken by Health Care Professionals Attending COVID-19 Patients among Dental Students. *Eur J Dent.* 2020 Dec;14(S 01):S105–9.
19. Logeshwari R, Rama Parvathy L. Generating logistic chaotic sequence using geometric pattern to decompose and recombine the pixel values. *Multimed Tools Appl.* 2020 Aug;79(31-32):22375–88.
20. Johnson J, Lakshmanan G, M B, R M V, Kalimuthu K, Sekar D. Computational identification of MiRNA-7110 from pulmonary arterial hypertension (PAH) ESTs: a new microRNA that links diabetes and PAH. *Hypertens Res.* 2020 Apr;43(4):360–2.
21. Paramasivam A, Priyadharsini JV, Raghunandhakumar S, Elumalai P. A novel COVID-19 and its effects on cardiovascular disease. *Hypertens Res.* 2020 Jul;43(7):729–30.
22. Pujari GRS, Subramanian V, Rao SR. Effects of *Celastrus paniculatus* Willd. and *Sida cordifolia* Linn. in Kainic Acid Induced Hippocampus Damage in Rats. *Ind J Pharm Educ.* 2019 Jul 3;53(3):537–44.
23. Rajkumar KV, Lakshmanan G, Sekar D. Identification of miR-802-5p and its involvement in type 2 diabetes mellitus. *World J Diabetes.* 2020 Dec 15;11(12):567–71.
24. Ravisankar R, Jayaprakash P, Eswaran P, Mohanraj K, Vinitha G, Pichumani M. Synthesis, growth, optical and third-order nonlinear optical properties of glycine sodium nitrate single crystal for photonic device applications. *J Mater Sci: Mater Electron.* 2020 Oct;31(20):17320–31.
25. Wu S, Rajeshkumar S, Madasamy M, Mahendran V. Green synthesis of copper nanoparticles using *Cissus vitiginea* and its antioxidant and antibacterial activity against urinary tract infection pathogens. *Artif Cells Nanomed Biotechnol.* 2020 Dec;48(1):1153–8.
26. Vikneshan M, Saravanakumar R, Mangaiyarkarasi R, Rajeshkumar S, Samuel SR, Suganya M, et al. Algal biomass as a source for novel oral nano-antimicrobial agent. *Saudi J Biol Sci.* 2020 Dec;27(12):3753–8.
27. Alharbi KS, Fuloria NK, Fuloria S, Rahman SB, Al-Malki WH, Javed Shaikh MA, et al. Nuclear factor-kappa B and its role in inflammatory lung disease. *Chem Biol Interact.* 2021 Aug 25;345:109568.
28. Rao SK, Kalai Priya A, Manjunath Kamath S, Karthick P, Renganathan B, Anuraj S, et al. Unequivocal evidence of enhanced room temperature sensing properties of clad modified Nd doped mullite Bi<sub>2</sub>Fe<sub>4</sub>O<sub>9</sub> in fiber optic gas sensor [Internet]. Vol. 838, *Journal of Alloys and Compounds.* 2020. p. 155603. Available from: <http://dx.doi.org/10.1016/j.jallcom.2020.155603>
29. Bhavikatti SK, Karobari MI, Zainuddin SLA, Marya A, Nadaf SJ, Sawant VJ, et al. Investigating the Antioxidant and Cytocompatibility of *Mimusops elengi* Linn Extract over

Human Gingival Fibroblast Cells. *Int J Environ Res Public Health* [Internet]. 2021 Jul 4;18(13). Available from: <http://dx.doi.org/10.3390/ijerph18137162>

30. Marya A, Karobari MI, Selvaraj S, Adil AH, Assiry AA, Rabaan AA, et al. Risk Perception of SARS-CoV-2 Infection and Implementation of Various Protective Measures by Dentists Across Various Countries. *Int J Environ Res Public Health* [Internet]. 2021 May 29;18(11). Available from: <http://dx.doi.org/10.3390/ijerph18115848>
31. Barma MD, Muthupandiyam I, Samuel SR, Amaechi BT. Inhibition of Streptococcus mutans, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish. *Arch Oral Biol*. 2021 Jun;126:105132.
32. Vijayashree Priyadharsini J. In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens. *J Periodontol*. 2019 Dec;90(12):1441–8.
33. Priyadharsini JV, Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species [Internet]. Vol. 94, *Archives of Oral Biology*. 2018. p. 93–8. Available from: <http://dx.doi.org/10.1016/j.archoralbio.2018.07.001>
34. Uma Maheswari TN, Nivedhitha MS, Ramani P. Expression profile of salivary micro RNA-21 and 31 in oral potentially malignant disorders. *Braz Oral Res*. 2020 Feb 10;34:e002.
35. Gudipani RK, Alam MK, Patil SR, Karobari MI. Measurement of the Maximum Occlusal Bite Force and its Relation to the Caries Spectrum of First Permanent Molars in Early Permanent Dentition. *J Clin Pediatr Dent*. 2020 Dec 1;44(6):423–8.
36. Chaturvedula BB, Muthukrishnan A, Bhuvanaraghan A, Sandler J, Thiruvengkatachari B. Dens invaginatus: a review and orthodontic implications. *Br Dent J*. 2021 Mar;230(6):345–50.
37. Abbas B, Ministry of Health -Baghdad-Iraq, Najm AA. Evaluation of Gonial Angle, Ramus Height and Bigonial width in Relation to Age and Gender using Digital Panoramic Radiograph [Internet]. Vol. 18, *Diyala Journal of Medicine*. 2020. p. 55–61. Available from: <http://dx.doi.org/10.26505/djm.18024950922>
38. Larrazabal-Moron C, Sanchis-Gimeno JA. Gonial angle growth patterns according to age and gender. *Ann Anat*. 2018 Jan;215:93–6.
39. Hariemmy M, Boedi RM, Utomo H, Margaretha MS. Sex determination using gonial angle during growth spurt period: a direct examination [Internet]. Vol. 1, *Indonesian Journal of Dental Medicine*. 2018. p. 86. Available from: <http://dx.doi.org/10.20473/ijdm.v1i2.2018.86-89>
40. Huumonen S, Sipilä K, Haikola B, Tapio M, Söderholm A, Remes-lyly T, et al. Influence of edentulousness on gonial angle, ramus and condylar height [Internet]. Vol. 37, *Journal of Oral Rehabilitation*. 2010. p. 34–8. Available from: <http://dx.doi.org/10.1111/j.1365-2842.2009.02022.x>
41. Jensen E, Palling M. The gonial angle [Internet]. Vol. 40, *American Journal of Orthodontics*. 1954. p. 120–33. Available from: [http://dx.doi.org/10.1016/0002-9416\(54\)90127-x](http://dx.doi.org/10.1016/0002-9416(54)90127-x)

UNDER PEER REVIEW

