

Phytochemical Screening and *In-vitro* Free radical scavenging activity of Unani formulation Habb-e-asgand

ABSTRACT

Aim: Phytochemical Screening and *In-vitro* Free radical scavenging activity of Unani formulation Habb-e-asgand

Study Design: The investigation of the Phytochemical Screening and *In-vitro* Free radical scavenging activity of Unani formulation Habb-e-asgand was carried out.

Place and Duration of the Study: The present work has been carried out at Ali-Allana College of Pharmacy, Akkalkuwa, and duration February-2020 to November-2020.

Methodology: Habb-e-Asgand, a Unani polyherbal drug used in arthritis, gout, and joint pain, is a blend of many herbal medicinal plants. Scientific attempts to test and validate its effectiveness are scarce. The given Unani formulation was evaluated for DPPH radical scavenging activity, hydroxyl radical scavenging activity, and nitric oxide scavenging assay. The given formulation also evaluated the total antioxidant capacity and the reducing antioxidant capacity of ferrous metals.

Results: It was found that DPPH radical scavenging activity of HEA and BHT at a concentration of 100 μ g/ml was 95 \pm 0.74 and 59 \pm 0.94 μ g/ml. Hydroxyl radical scavenging activity of HEA, at the concentration of 100 μ g/ml, was 55 \pm 0.64, whereas at the same concentration catechin was 63 \pm 0.84 μ g/ml. The IC₅₀ value for nitric acid scavenging activity was found to be 49.60 \pm 1.57 for HEA, and 186.34 \pm 3.28 μ g/ml for ascorbic acid. The result obtained from TAC and FRAC activity demonstrates that HEA possessed the highest TAC and FRAC values in comparison to standards catechin and ascorbic acid.

Conclusion: From the previous results it was concluded that HEA has a good and potent strong antioxidant effect. HEA showed moderate to high TAC and FRAC with increased concentrations of the drugs in of standards catechin and ascorbic acid. HEA shows dose-dependent DPPH and hydroxyl radical scavenging activity. Nitric oxide generation was

observed after the incubation of solutions of sodium nitroprusside in PBS at 25°C for 150 min. HEA effectively reduced the generation of nitric oxide.

Keywords: Habb-e-Asgand, DPPH scavenging activity, Polyherbal, Phytochemical screening, Unani, etc.

ABBREVIATIONS

HEA: Habb-E-Asgand

DPPH: 2, 2-Diphenyl-1-Picrylhydrazyl

TAC: Total Antioxidant Capacity

FRAC: Ferrous Reducing Antioxidant Capacity.

BHT: Butylated hydroxytoluene

AA: Ascorbic Acid

CC: Catechin

NO: Nitric Oxide

1. INTRODUCTION

Most of Unani's medications are herbal, mineral, or a combination of the two. Most of these drugs are developed for specific indications, but some have general tonic and rejuvenating properties [1, 2]. In the absence of effective treatment for liver ailments. Unani drugs have been extensively studied to develop innovative therapies for liver disease and drug-induced liver toxicity [3, 4]. Most Unani drugs are safe, but toxicity, including liver injury, has also been observed [5, 6]. Furthermore, there are very few scientific attempts to certify and validate the safety of Unani's drugs [7, 8]. Therefore, the numerous studies reported have evaluated the hepatoprotective and antioxidant potential of an Unani drug, known as Habb-e-Asgand.

Habb-e-Asgand is a popular herbal preparation prescribed for arthritis, gout, and joint pain [9]. It is also known to have aphrodisiac properties [9]. Habb-e-Asgand contains *Withania somnifera* (L) Dunal (Solanaceae) (English name: winter cherry, Hindi name: ashwagandha) as the main constituent of **Table 1**, known for its various medicinal uses [10]. Briefly, Habb-e-Asgand Therapy comes in small, round, uniformly shaped pills made with the ingredients of the formulation composition shown in **Table 1**. The blend included Ajwain Desi [*Ptychotis ajowan* DC (Apiaceae) seeds], Asgand Nagauri (*W. somnifera*), Chob Bidhara [wood of *Gmelina Asiatica* L. (Lamiaceae)], Pippl Kalan-Desi [Fruit of *Piper longum* L. (Piperaceae), dried immature], Pipla Mool (the root of *P longum*) Moosli Siyah [*Curculigo orchoides* Gaertn. (Hypoxidaceae) stem], Satawar [*Asparagus racemosus* Willd. (Asparagaceae)], Zanjabeel-Khushk [rhizome of *Zingiber officinalis* Roscoe (Zingiberaceae)] and Qand Siyah Kohna [*Saccharum officinarum* L. (Poaceae)] as basic ingredients [9]. The high content of Qand Siyah Kohna is attributed to its use as a coating material in the preparation of pills.

A review of the literature revealed that the scientific study of the *In-vitro* Free radical scavenging activity of Unani formulation Habb-e-asgand was not previously conducted to evaluate the formulation traditional claim as an antioxidant. The main objective of the present research work is to study phytochemical screening and *in-vitro* free radical scavenging activity of Unani formulation Habb-e-asgand.

Table 1: Constituents of Habb-e-asgand

Sr. No	Plant Common Name	Plant Botanical Name
1	Ajwain Desi	<i>Ptychotis ajowan</i>
2	Asgand Nagauri	<i>Withania somnifera</i>
3	Chob Bidhara	<i>Gmelina Asiatica</i>
4	Moosli Siyah	<i>Curculigo orchiodes</i>
5	Pippl Kalan (Desi)	<i>Piper longum</i>
6	Pipla Mool	<i>Piper longum</i>
7	Satawar	<i>Asparagus racemosus</i>
8	Zanjabeel (Khushk)	<i>Zingiber officinale</i>
9	Qand Siyah Kohna	<i>Saccharum officinarum</i>

2. MATERIAL AND METHODS

2.1 Chemicals and Reagents

Folin Ciocalteu reagent, 2, 2-diphenyl-1-picryl hydrazyl radical (DPPH), phenazine methosulfate, nicotinamide adenine dinucleotide, sodium nitroprusside (SNP), trichloro acetic acid (TCA), thio barbituric acid (TBA), and L-ascorbic acid were purchased from Lab Trading Laboratory Aurangabad. Catechin and rutin were purchased from Sigma Chemicals, India. All other chemicals and solvents used were of analytical grade available commercially.

2.2 Physico-chemical analysis of Habb-e-asgand

The specification of the Habb-e-Asgand mixture was evaluated by performing physicochemical analyzes of appearance, color, taste, odor, pH value, friability, hardness, weight change, disintegration time, etc.

2.3 Preparation of aqueous extracts

Water-soluble extract: 5 g of the air-dried drug mixture was macerated and coarsely sprayed with 100 ml of water in a closed flask for 24 h with frequent agitation for 6 h and allowed to stand for 18 h. The extract was quickly filtered with precautions due to loss of solvent. A 25 ml aliquot of the filtrate was evaporated to dryness on a flat-bottomed tarred dish at 105 ° C. The percentage of water extraction was calculated concerning the air-dried drug [10, 11].

2.4 Preliminary phytochemical screening of the extract

The extract of Habb-e-Asgand leaves obtained during the extraction process was subjected to preliminary phytochemical examination for the presence of various phytoconstituents using reported methods [11].

2.5 *In vitro* Models

2.5.1 Determination of total antioxidant capacity (TAC)

TAC of samples was determined by the previously described method [12]. This assay is mainly based on the reduction of Mo (VI) to Mo (V) by the drugs/samples and thus the formation of green-colored phosphate/ Mo (V) complex at acidic pH. Samples/standard approx. 0.5 ml at different concentrations of 12.5-150 µg/ml was mixed with 3 ml of a mixture containing 0.6 M sulphuric acid, 28 mM sodium phosphate, and 1% ammonium molybdate. The test tubes containing the above mixtures were incubated at 95°C for 10 min so that the reaction can be completed. After cooling the reaction mixture at room temperature, the absorbance of the sample was measured at 695 nm using a spectrophotometer against a blank solution. Catechin was used as a standard. A blank solution was prepared with 3 ml of the reaction mixture and the same volume of the solvent was added which was used for samples/standard. The blank was also incubated for 10 min at 95°C followed by measurement of absorbance at 695nm. Increased absorbance indicates increased total antioxidant capacity. Standard/samples were used at five different concentrations ranging from 12.5 to 150 µg/mL for each antioxidant assay. Concentrations were selected based on trial and error to fit the concentration that can represent the rational change of antioxidant activity with increasing concentration of samples.

2.5.2 Ferrous reducing antioxidant capacity assay (FRAC)

FRAC of samples was evaluated by the previously described method [12]. The Fe²⁺ can be monitored by measuring the development of Per's Prussian blue at 700 nm. 0.25 ml of standard/sample solutions at different concentrations of 12.5-150 µg/ml, 0.625 ml of potassium buffer (0.2 M), and 0.625 ml of 1% potassium ferricyanide solution were added into the test tubes. The above reaction mixtures were incubated for 20 min at 50°C to complete the reaction. Then 0.625 ml of 10% trichloroacetic acid solution was added to the test tubes. The above mixture was centrifuged at 3000 rpm for 10 min, later which 1.8 ml supernatant was withdrawn from the test tubes and mixed with 1.8 ml of distilled water and 0.36 ml of 0.1% ferric chloride solution. The absorbance was measured at 700 nm using a spectrophotometer against blank. A blank solution contained the same reaction mixture without the sample/standard and was incubated under the same conditions and the absorbance was measured at 700 nm. Increased absorbance of the reaction mixture indicates increased reducing capacity. The experiment was repeated three times at each concentration.

2.6 Free Radical scavenging activity

2.6.1 DPPH radical scavenging assay

The free radical scavenging ability of the drugs was tested by DPPH radical scavenging assay as previously described method [12]. The hydrogen atom donating ability of the drugs was determined by the decolorization of methanol solution of 2, 2-diphenyl-1-picrylhydrazyl (DPPH). DPPH produced purple/violet color in methanol solution and fades in presence of antioxidants to shades of yellow color. A solution of 0.1 mM DPPH in methanol was prepared, and 2.4 ml of this solution was mixed with 1.6 ml of extract in methanol at varied concentrations of 12.5 to 150 µg/ml. The solution mixture was mixed thoroughly and left in dark at room temperature for 30 min. The absorbance was measured spectrophotometrically at 517 nm. Butylated hydroxytoluene (BHT) was used as a reference. Percentage of DPPH radical scavenging activity was calculated by the following equation:

$$\% \text{DPPH radical scavenging activity} = [(A_0 - A_1) / A_0] \times 100 \text{ ----- (1)}$$

Where A_0 is the absorbance of the control, and A_1 is the absorbance of the drugs/standard. Then % of inhibition was plotted against concentration, and from the graph, IC_{50} was calculated. The experiment was repeated three times at each concentration.

2.6.2 Hydroxyl radical scavenging activity

Hydroxyl radical scavenging activity of the drugs was determined by the previously described method [12]. Hydroxyl radical was generated by the Fe^{3+} -ascorbate-EDTA- H_2O_2 system (Fenton reaction). The assay is based on the quantification of the 2-deoxy-D-ribose degradation product, which forms a pink chromogen upon heating with TBA at low pH. The reaction mixture contained 0.8 ml of phosphate buffer solution (50 mmol L^{-1} , pH 7.4), 0.2 ml of drug/standard at different concentrations of 12.5-150 µg/ml, 0.2 ml of EDTA (1.04 mmol L^{-1}), 0.2 ml of $FeCl_3$ (1 mmol L^{-1}) and 0.2 ml of 2-deoxy-D-ribose (28 mmol L^{-1}). The mixtures were kept in a water bath at 37°C and the reaction was started by adding 0.2 ml of ascorbic acid, AA (2 mmol L^{-1}), and 0.2 ml of H_2O_2 (10 mmol L^{-1}). After incubation at 37°C for 1 h, 1.5 ml of HCL (25%). The reaction mixture was heated at 100°C for 15 min and then cooled down with water. The absorbance of the solution was measured at 532 nm with a spectrophotometer. The hydroxyl radical scavenging capacity was evaluated with the inhibition of the percentage of 2-deoxy-D-ribose oxidation on hydroxyl radicals. The percentage of hydroxyl radical scavenging activity was calculated according to the following formula:

$$\% \text{hydroxyl radical scavenging activity} = [A_0 - (A_1 - A_2)] \times 100 / A_0 \text{ ----- (2)}$$

Where A_0 is the absorbance of the control without a sample. A_1 is the absorbance after adding the sample and 2-deoxy-D-ribose. A_2 is the absorbance of the sample without 2-deoxy-D-ribose. Then % of inhibition was plotted against concentration, and from the graph, IC_{50} was calculated. The experiment was repeated three times at each concentration.

2.6.3 Nitric oxide scavenging assay

The method was performed according to the process previously described [13]. 2 ml of 10mM sodium nitroprusside solution which was prepared in phosphate buffer saline (PBS) pH 7.4 was mixed with 0.5 ml of drug solutions at various concentrations ranging from 10 to 50 µg/ml and ascorbic acid at

concentrations ranging from 25 to 200 µg/ml. This mixture was incubated at 25°C. After 150 min, 0.5 ml of incubation solution was collected which was then mixed with 0.5 ml of Griess reagent [1.0 ml sulfanilic acid reagent (0.33% prepared in 20% glacial acetic acid at room temperature for 5 min 1 ml of naphthyl ethylene diamine dihydrochloride (0.1% w/v)]. The mixture was incubated at room temperature for 30 min, followed by the measurement of absorbance at 540 nm using a spectrophotometer (Shimadzu UV-1800).

3. RESULTS AND DISCUSSION

3.1 Physico-chemical analysis of Habb-e-asgand

The Habb-e-Asgand pills were brown and spherical (solid) with a distinctive Asgand taste and smell. The pills were stored in a cool, dark place in tightly closed containers, protected from moisture, light, and temperature. Physico-chemical analysis showed a slightly basic pH of 6.5 as shown in Tables 2 and 3.

Table 2: Organoleptic Character

Sr. No	Parameters	Observation
1	Size	Round
2	color,	Light Brown
4	Taste,	Pungent
5	Appearance	Habb (Pills)

Table 3: Physiochemical Character

Sr. No	Parameters	Habb-e-agand
1	Friability Test	0.219%
2	Hardness Test	12.5 kg/cm ² (Monsanto)
3	Weight Variation	0.0668
4	Disintegration time	55 min
5	the pH of 1% Solution	6.5
6	10% Solution	5.2

3.2 Preliminary phytochemical screening of the extract

The medicinal properties of Habb-e-Asgand pills are perhaps due to the presence of various secondary metabolites such as alkaloids, tannins, flavonoids, proteins, and mucilages. Therefore, preliminary screening tests can be useful in the detection of bioactive ingredients and can subsequently lead to drug discovery and development. Furthermore, these tests facilitate their quantitative estimation and qualitative separation of pharmacologically active chemical compounds [13]. Preliminary phytochemical screening with various qualitative chemical tests revealed the presence of carbohydrates, reducing sugars, protein, amino acid, alkaloids, glycosides, and tannins in the Habb-e-Asgand extract, and flavonoids, saponins, fats & oils, and steroids were absent in the Habb-e-Asgand extract.

3.3 Determination of TAC and FRAC

The TAC and FRAC of HEA are shown in **Table 4**. HEA showed a considerable antioxidant activity compared to standard catechin. At the concentration of 100µg/ml, the absorbance of HEA was in the range of 0.584±0.053 to 1.89±0.076; while at 150µg/ml the range was 0.874±0.043 to 2.394±0.064. Total antioxidant activity was increased with an increase in the dose of the drugs.

HEA showed moderate to high FRAC with increased concentrations of the drugs. At 100µg/ml, the absorbance was in the range of 0.975±0.056 to 3.409±0.065, while at 150µg/ml, the range was 1.345±0.086 to 3.225±0.067. These results demonstrate that HEA possessed the highest TAC and FRAC values, thus closely resembling that of the standards catechin and ascorbic acid.

Table 4: Absorbance HEA at two different concentrations (n=3, X ± SEM)

Drugs	TAC		FRAC	
	At 100 µg/ml	At 150 µg/ml	At 100 µg/ml	At 150 µg/ml
HEA	1.378 ± 0.036 ^a	1.784 ± 0.019	2.235 ± 0.137	2.986 ± 0.107
CA	1.89 ± 0.076	2.394 ± 0.064	-	-
AA	-	-	3.409 ± 0.065	3.225 ± 0.067

Where, CA and AA represent catechin and ascorbic acid; ^a each value is the average of three analyses ± standard deviation

3.4 Determination of DPPH radical scavenging activity of HEA

Table 5 shows the free radical scavenging activity of HEA. At a concentration of 100µg/ml, the scavenging activity of HEA and BHT was 95±0.74 and 59±0.94µg/ml. The extract showed concentration-dependent antioxidant activity by inhibiting the DPPH radical with a good inhibition value. The method is based on the reduction of the DPPH solution in the presence of a hydrogen donor antioxidant, due to the formation of the non-radical form DPPH-H by the reaction. The extract was able to reduce the stable DPPH radical to the yellow diphenyl picrylhydrazine. Ascorbic acid has been found to reduce and discolor 1,1-diphenyl-2-picrylhydrazyl due to its ability to donate hydrogen. [40] HEA extract appears to have the ability to donate hydrogen and act as an antioxidant. The elimination effect increases with the increasing concentration of the extract.

Table 5: Determination of DPPH radical scavenging activity of HEA

Concentration (µg/ml)	BHT	HEA
0	0	0
12.5	80	65
25	93	73
50	95	79
100	95	90
150	95	90

3.5 Determination of hydroxyl radical scavenging activity of HEA

Table 6 shows the hydroxyl radical scavenging activity of HEA and present drugs have shown a dose-dependent activity. The hydroxyl radical is a highly reactive oxygen-centered radical formed by

the reaction of various hydroperoxides with transition metal ions. Attacks proteins, DNA, polyunsaturated fatty acids on membranes, and most biological molecules. At the concentration of 100µg/ml, the scavenging activity of HEA was 55±0.64, whereas at the same concentration catechin was 63±0.84µg/ml.

Table 6: Determination of hydroxyl radical scavenging activity of HEA and CA

Concentration (µg/ml)	CA	HEA
0	0	0
12.5	16	36
25	22	40
50	37	50
100	63	55
150	84	63

3.6 Determination Nitric oxide scavenging assay

The extract showed moderate nitric oxide scavenging activity in a dose-dependent manner (IC₅₀ = 49.60 µg/ml). The plant/plant products may have the property of counteracting the effect of NO formation and, in turn, may be of considerable interest in preventing the harmful effects of excessive NO production in the human body. Furthermore, the elimination activity can also help stop the chain of reactions initiated by the excessive generation of NO harmful to human health. The extract showed moderate nitric oxide scavenging activity. The% inhibition increases with increasing concentration of the extract. NO generation was observed after the incubation of solutions of sodium nitroprusside in PBS at 25°C for 150 min. HEA effectively reduced generation NO. The IC₅₀ was found to be 49.60±1.57 for HEA, and 186.34±3.28µg/ml for ascorbic acid **Table 7**.

Table 7: IC₅₀ value for *in-vitro* nitric acid scavenging activity

Sr.No	Sample	Nitric oxide (µg/ml)
1.	HEA	49.60±1.57
4.	Ascorbic acid	186.34±3.28

4. CONCLUSION

To characterize the antioxidant activity of HEA extracts, it is desirable to subject it to a battery of tests evaluating the range of activities such as hydroxyl radical scavenging activity test, nitric oxide scavenging activity test, free radical scavenger DPPH in formulation by Habb-e-asgand. The *in-vitro* antioxidant activities of the aqueous extract have indicated the effectiveness of the formulation as a source of natural antioxidants that will be used to reduce oxidative stress with consequent health benefits. The activity of Habb-e-asgand could be due to the presence of different polyherbal constituents, which is a known potent antioxidant. Further research is needed for the isolation and characterization of the phytoconstituents responsible for its antioxidant property.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

NOTE:

The study highlights the efficacy of " sigma, Unani " which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

5. REFERENCE

1. George SK, Rajesh R, Kumar S, Sulekha B, Balaram P. A polyherbal ayurvedic drug– Indukantha Ghritha as an adjuvant to cancer chemotherapy via immunomodulation. *Immunobiology*. 2008;213(8):641-9.
2. Singh RS, Dhaliwal R, Puri M. Production of high fructose syrup from Asparagus inulin using immobilized exoinulinase from *Kluyveromyces marxianus* YS-1. *J. Ind. Microbiol. Biotechnol*. 2007;34(10):649-55.
3. Kar P, Asim M, Sarma MP, Patki PS. HD-03/ES: a promising herbal drug for HBV antiviral therapy. *Antivir. Res*.2009;84(3):249-53.
4. Nwaehujor CO, Udeh NE. Screening of ethyl acetate extract of *Bridelia micrantha* for hepatoprotective and anti-oxidant activities on Wistar rats. *Asian Pac. J. Trop. Med*.2011;4(10):796-8.
5. Licata A, Macaluso FS, Craxì A. Herbal hepatotoxicity: a hidden epidemic. *Intern Emerg Med*. 2013;8(1):13-22.
6. Oberbaum M, Samuels N, Ben-Arye E, Amitai Y, Singer SR. Apparent life-threatening events in infants and homeopathy: an alternative explanation. *Hum Exp Toxicol*2012;31(1):3-10.
7. Leone L, Marchitello M, Natilli M, Romano MF. Measuring the effectiveness of homeopathic care through objective and shared indicators. *Homeopathy*. 2011;100(04):212-9.

8. Rutten AL. Comparison of the effectiveness of frequently and infrequently used homeopathic medicines. *Homeopathy*. 2011;100(03):175-82.
9. Krishnan MK. The useful plants of India. Publication and Information Directorate, CSIR, New Delhi. 1992:918.
10. Elsakka M, Pavelescu M, Grigorescu E. *Withania somnifera*, a plant with a great therapeutical future. *Rev Med Chir Soc Med Nat Iasi*.1989;93(2):349-50.
11. Ali M, Khan SA, Chang PS, Haque R, Bhatia K, Ahmad S. Habb-e-Asgand, polyherbal Unani formulation, protects the liver and antioxidative enzymes against paracetamol induced hepatotoxicity. *Pharm. Biol.* 2014;52(4):506-15.
12. Rahman MM, Islam MB, Biswas M, Alam AK. In vitro antioxidant and free radical scavenging activity of different parts of *Tabebuia pallida* growing in Bangladesh. *BMC Res. Notes*.2015;8(1):1-9.
13. Borra SK, Gurumurthy P, Mahendra J, Ch R. Antioxidant and free radical scavenging activity of curcumin determined by using different in vitro and ex vivo models. *J. Med. Plant Res.* 2013;7(36):2680-90.
14. Parmer NS, Prakash S. Evaluation of analgesics, anti-inflammatory, and antipyretic activity. In: *Screening Methods in Pharmacology*. Narosa Publishing House Pvt. Ltd.; New Delhi, India: 200;1(1).211-215.
15. Patil S, Anarthe S, Jadhav R, Surana S. Evaluation of anti-inflammatory and In-vitro Antioxidant Activity of Indian Mistletoe, the Hemiparasite *Dendrophthoe falcata* L. F. (Loranthaceae). *Iranian J Pharm Res.* 2011;10(2):253-259.