

## Review Article

### BRONCHIAL ASTHMA IN THE ERA OF COVID-19

#### Abstract:-

Coronavirus disease also (COVID 19) is a worldwide pandemic contagious infection produced through the Coronavirus SAR.S-CoV2, and aberrant, overactive immunity and "chemokines" have been hypothesized as possible pathogenic reasons for fast COVID-19 development. Because of a poor viral immunogenic reaction with a potential for aggravation evoked by group of similar respiratory virus, asthmatic patients have increased susceptibility to and severe from covid infection. Existing research, however, has not found an anticipated proportion of patients of asthma between COVID patient. Features of the immune responses of type2 nature, such as type2 cytokine such as IL-4, IL-13 and eosinophil buildup, may give protection against Corona virus. Furthermore, traditional asthma treatments like as inhalation of corticosteroid, AIT (allergens immunotherapy), and monoclonal antibodies (anti-IgE) may minimize the chance for disease by the causative agent in asthmatics by reducing inflammation and strengthening antiviral defense. The connections between COVID along with asthma need to be researched upon more. Infection with a respiratory virus causes up to eighty percent of acute asthma attacks in kids and 50% of similar instances in adults, with RV being the most common cause. (1) CoVs are also seen in asthma exacerbations, with an 8.4 percent frequency. Surprisingly, no cases of asthma attacks have been reported as a result of because of COVID-19. As a result, during the SARS and MERS outbreaks, there seem to be minimal reports of asthma attacks. Such specific events still have to be thoroughly documented. The study's main goal was to find out how common asthma was among COVID-19 participants. COVID-19 was used to examine asthmatic and non-asthmatic patients' clinical features and comorbidities. We also looked at the possibility of being admitted to the hospital as a result of asthma and/or the usage of inhaled corticosteroids.

**Keywords:** Covid-19, Bronchial Asthma, Attacks, Immunity, Corona, Allergy

#### Introduction:

Infection with a respiratory virus causes up to eighty percent of acute asthma attacks in kids and 50% of similar instances in adults, with RV being the most common cause (1). (+)CoVs are also seen in asthma exacerbations, with an 8.4 percent frequency. Surprisingly, no cases of asthma attacks have been reported as a result of COVID-19. As a result, during the SARS and MERS outbreaks, there seem to be minimal reports of asthma attacks. Such specific events still have to be thoroughly documented (2). (2)-Asthmatic people have a weakened immune system reaction

**Comment [YT1]:** Use an other synonymous word to avoid redundancy e.g like

**Comment [YT2]:** Do you mean rhino virus (RV)?

**Comment [YT3]:** When you use abbreviation or acronym for the first time in your document, write the full meaning and then the abbreviation in brackets.

**Comment [YT4]:** When you use abbreviation or acronym for the first time in your document, write the full meaning and then the abbreviation in brackets.

**Comment [YT5]:** Do the same

to viral infections, and respiratory viral diseases can trigger or aggravate asthmatic attacks, treating asthma patients becomes more difficult at the time of the COVID19 outbreak. In general, pre-existing asthma appears to have an effect on SARS\_CoV2 vulnerability & disease progression. Previous research, even though, has not found a significant presence of asthma in COVID19 patients. Few studies in published COVID\_19 research included it as a condition, that is inadequate to establish a strong conclusion (3). (3)

**Comment [YT6]:** Rephrase, you can remove the conjunction, even though

A research from Lombardy, Italy, likewise revealed a lesser presence of it. Only data from New York City, United States of America, reveal an asthma prevalence of 9%, however no thorough determination of the actual clinical phenotypes and therapies of included patients is available. According to State of NY mortality data, asthma was not included in the first ten comorbidities. In conclusion, their findings show demonstrate allergic illnesses are not a risk factors or exacerbating factors for COVID\_19 (4). (4)

**Comment [YT7]:** When you use abbreviation or acronym for the first time in your document, write the full meaning and then the abbreviation in brackets.

Symptoms mostly associated with COVID\_19 are dry cough, and shortness of breath which may be associated to an acute asthma worsening. Most primarily fever is common with COVID19, although it can occur with any disease triggered asthma exacerbation. Given the variety of clinical manifestations of COVID\_19, screening techniques for COVID19 should be used to anybody experiencing increasing symptoms of respiration, including those having asthma. People who are commencing screening must wear appropriate personal protective equipment (5). (5)

**Comment [YT8]:** Symptoms of what? Do you mean Asthma?

**Comment [YT9]:** Please put a hyphen between COVID and 19 through the whole of the document

### Can having Asthma Increases the chances of having COVID-19?

Individuals with long term conditions and above 65 years in age were in the CovidCOVID-19 groups at risk, as per the C.D.C-Center for Disease Control and Prevention (CDCCenter for Disease Control and Prevention) (6). Asthmatic patients with a medium to heavy disease were also included in this danger category. Nevertheless, asthma illness was uncommon among individuals of COVID-19, particularly the Chinese observational case series. There were no incidences of asthma or allergic rhinitis recorded in a research of 140 Wuhan patients' clinical features and allergies; nevertheless, only two cases of urticaria were identified (7). The risk factors were examined in a different research involving 549 hospitalised patients in China's Wuhan. Asthma was found to be present in 0.9 percent (n= 5) of the comorbidities (8). The researchers hypothesized that IFN-, IL-12, IL-8, IL-6, IL-1, and TNF-cytokines were produced in response to a Th-1 cellular reaction, whereas a Th-2 immunogenic reaction might be protective opposing the illness. There were 11 different phenotypic and clinical characteristics discovered in 11 separate patients, but no instance of asthma was described. In a separate study of 1590 patients with comorbidities, no occurrence of asthma was found (9). 180 concomitant conditions (12.1 percent) were identified between 1482 hospitalised cases in a monthly update analyzing hospitalisation and case features in the US. These were confirmed cases from laboratory, and the occurrence of the condition was reported to be 17 percent and 27.3 percent in each age categories and in the 18-49 age range, among the 180 cases with comorbidities. However, these are the rates for comorbid conditions only; when all hospitalised cases are taken into account, the incidence drops dramatically. In the areas of New York, asthma cases were reported to be present in 9% of COVID-19 participants. While some meta-analyses found no evidence of a relation between asthmatic patients having an higher danger of Covid 19 mortality in hospitals (11), others found a link between asthma and an increased danger of COVID-19

**Comment [YT10]:** Keep uniformity of subtitle writing, either capitalize each word or sentence case. Review all the titles and subtitles underneath also

mortality in hospitals (12). Patient had not been utilizing inhaler steroids, which is significant. The finding that diffused damage of alveoli and mucous plaques found in patients lung sections were connected to asthmatic.

The presence of lesser report associated to ~~Covid~~ COVID-19 and asthma, according to some experts, could be attributed to three factors (13). To begin with, some patients were misdiagnosed or misidentified, especially in China. It's possible that these patients' asthma was not documented in their hospital records. Another explanation could be that respiratory disorders trigger a distinct immunological reaction.

The third option is that inhaled corticosteroids, which are used to treat chronic respiratory illnesses, may prevent against COVID-19 or improve symptoms.

### **Does Asthma and Allergy or their therapy affecting ACE-2 Receptor Genes Expression in the Airway??**

Formatted: Space After: 0 pt

By attaching to ACE2 receptors, the SARS-CoV-2 penetrates the cell. The appearance of ACE2 gene in asthma sufferers was studied (14,15). Type 2 markers were also measured, including FeNO (fraction exhaled nitric oxides), total Ig E, and blood eosinophil. Allergy sensitivity has also been found to be inversely associated to ACE-2 appearance in the nasal epithelium. Furthermore, a strong negative connection was discovered between Type 2 biomarker (specific IgE, Total Ig E, FeNO) and nasal epithelial IL-13 and ACE-2 expressions. (14). Reduced levels of sensors in the body with which the SARS-CoV-2 attaches might protect allergic asthma patients from illness. The second research contributing in our knowledge of how asthmatic, ICS, and ACE2 expression are linked (15).

### **Does the Sars-CoV-2 Virus Cause Asthma?**

The respiratory tract virus in the past have been linked to attacks of asthma. In previous investigations, respiratory tract viruses like RV (rhinoviruses), RSV (respiratory syncytial viruses), herpes simplex viruses, EnV (enteroviruses), and IfV (influenza) were found to be the majority of the causes of the attacks ~~(16)~~. Non-pandemic coronaviruses were also shown to induce asthma attacks, though less frequently. SARS-CoV and MERS-CoV, two previous pandemic coronaviruses, were not connected to the attacks ~~=(16)~~.

### **COVID-19 and Previously Existing Asthma:**

• **Asthmatic Patients with Inadequate Antiviral Response:** This is widely assumed that antiviral and allergy reactions are two different arms of resistance that are mutually controlled and involve a complex network of interactions. IFNs, a kind of antiviral cytokine, play a critical function in this. It's been suggested that people with asthma are more sensitive to allergic reactions that can override antiviral medication, leaving them more vulnerable to viral infections and having a poor immune response (17). In asthmatic bronchial epithelial cells, IFN production is decreased, which results in altered apoptosis and enhanced rhinovirus (RV) replication. After

Comment [YT11]: Review the spacing after and before titles and subtitles

RV infection, asthma patients have been found to have decreased IFN / expression by bronchial epithelial cells, which has been linked to higher viral load and worse clinical consequences. (18).

- **Eosinophilic inflammation and Covid disease:** Eosinophils are well-known for their function in allergic disorders, such as asthma. The prospective impacts due to eosinophil on Covid are worth noting. Although this behavior is not observed in all conditions, previously conducted research has suggested that eosinophilic reaction must have a part in viral clearance and host antiviral defense. Furthermore, eosinophils with ssRNA can be activated by activating the TLR7MyD88 signaling pathway, which could lead to the clearance of RSV and a reduced dysfunction of lung. The ability in eosinophil in its fight against viral infections may thus account for the decreased presence of asthmatics among COVID19 patients. (19)

### **Asthma patients' susceptibility to COVID-19 infection**

During initial stages of epidemic, research from China and Italy highlighted asthma as one of the key clinical risk factors for SARSCoV2 infection.

Comment [YT12]: Check spacing

According to research conducted in India, Brazil, U.A.E and Russia, decrease of asthma incidence have been reported in COVID19 patients. However, research from the US and UK found that asthma comorbidities percentages in COVID19 patients were comparable to or greater than in the general population. (20)

In summary, the presence of asthma in COVID19 patients varies greatly with different countries; Most countries did not report higher rates of COVID19 infection in asthma patients, but instead comparable or lesser rates of COVID19 disease in comparison to the general public in the relevant area, likely owing to several factors, consisting of a lesser percentage of non-type 2 phenotypes. In fact, a cohort from the nation of Korea appeared that non allergic asthmatic people had increase chances for a positive SARSCoV2 test than patients with allergic asthma. (21).

### **Asthma (severe form) along with COVID-19**

Despite extensive therapy, a small percentage of asthma cases (5-10%) have uncontrolled or inadequately managed asthma. As one might anticipate these patients to be more susceptible to SARSCoV2 infection, but little data is obtainable to support such a theory. Regardless of the lack of solid data, the U.S.A CDC (center for Control and Prevention) issued a warning for patients having moderate forms to severe forms of asthma are more likely of contracting COVID19 and suffering catastrophic consequences as a result of the disease. (22) Although patients with level 5 asthma spent more time in the hospital compared with level 1 asthmatic patients in one variable and multi variable analyses, asthma severity was not independently associated determinant of poor health outcomes from COVID19 in another experiment.

A huge COVID19 hospital community in Italy revealed that patients with an adverse COVID19 outcome (death / requirement for ventilation versus release from home without intrusive treatments) had more severe asthma. (23) At last it is considered everywhere that old age, obese body, having cardiovascular diseases, and diabetic patient are all at risk of having a poor COVID19 score. Chronic respiratory disorders, such as asthma, may or may not be considered

risk factors. Many studies on this topic have devised contradicting plans, suggesting that to types of factors that may affects Covid 19 vulnerability along with the intensity in asthmatic patient. This would include severity of asthma, asthmatic genotypes/phenotype, asthmatic prescriptions, and comorbidities.

The documented occurrences of significant COVID19 incidences in patients with asthma are influenced by the factors other than the patient. Furthermore, local circumstances (screening recommendations or screening suggestions, including the elderly or those with comorbidities such as asthma) and the diagnostic tools used to diagnose asthma and COVID19 can have a crucial influence.(24)

**In the context of COVID-19, do anti-asthmatic medications play a protective or beneficial role??**

---

### **COVID-19 and inhaled antiasthmatic treatments**

Maintaining CSIs is expected to give protection as well, however there is no indication for CSI advantages or downsides in COVID19 . Numerous key questions emerge. Currently, there is really no indication of a higher danger for COVID19 disease among patients with asthma who use ICS on a regular basis. (25)

Asthma patients should continue to use their inhaler corticosteroids and incorporate therapies during the outbreak, according to guidelines and experts (26-29). It is suggested that nebulizers not be used in hospitals because of the danger of illness transmission to healthcare workers and others, and that spirometry need not be performed unless absolutely required. In asthma attacks, a measured dose inhalation through a mouth piece or a closely fitting ventilation mask should be utilized instead of a nebulizer (26). It has been proven that attaching a nebulizer with a filter is useful in catching inhaled aerosolized particles (30-32). The filters help to limit the amount of aerosols that are released into the environment. Mesh nebulizers are a good alternative to jet nebulizers (30). These recommendations should be taken into account when treating COVID19 and asthma patients. There is a recommendation to postpone routine check-up appointments for patients with mild asthma and also those who have controlled asthma and instead tell them by phone calls, e-mail, or text message (26-29). The potential of COVID-19 transmission makes hospital visits for these individuals unfavourable. Patients who are well and have not had an urgent appointment with in past 6-12 months, as well as those who have had one oral steroid in the previous 6 months, may have their face-to-face examinations postponed. Telehealth services may be used to manage these patients remotely-(29). Patients who have had an attack in the last 3-6 months and have a history of an emergency visit or hospitalisation should be given priority. COVID19 screening should be administered to patients admitted with worsening asthma or an acute attack, and the risk of SARS-CoV-2 infection and the need for testing should be questioned- (29).

### **Asthma Attacks and Systemic Corticosteroids**

The usage for systemic steroids in the treatment of COVID-19 associated lung damage is a source of debate. Prednisolone has been linked to the prolongation of viral replication. Short-term administration of systemic steroids at modest doses (0.5-1 mg/kg, 7 days) is effective for COVID-19 associated lung damage, according to the Chinese Thorax Society (31). According to Russel et al., the use of systemic steroids in the SARS-CoV and MERS-CoV infections slowed viral RNA clearance, induced different side effects, and there was no clinical evidence to justify steroid treatment in the COVID-19 illness (32). During the pandemic, the Global Initiative for Asthma (GINA) recommends using oral steroids for moderate attacks since it reduces hospitalizations (26). The Canadian Society recommends the use of systemic steroids in asthma attacks, whether or not they are caused by COVID-19 (28). It's vital to keep patients safe from asthma episodes at this period.. Patients' inhaler steroid doses could be increased to prevent them from having an attack (26,29). During this time, giving patients a written action plan and following up via remote telemedicine are appropriate ways (26,29). During the COVID-19 pandemic, Levin et al. published an acute asthma care protocol for doctors. (35) They believe that high-dose MDI and quick systemic corticosteroid treatment can help prevent asthma attacks from worsening.

Formatted: Pattern: Clear

### **COVID-19 and Biological therapies**

Biologics are used to treat severe asthma in people who have been unable to control their symptoms with existing therapies. These three anti-IL5 / IL5r drugs work primarily to combat eosinophils by lowering or depleting eosinophils in tissues and peripheral circulation. Another 2 therapies (anti-IgE and anti-IL4 / IL13) mostly decrease type 2 immunity. As a result, another central issue is whether eosinophils and/or immunomodulation via the type 2 pathway play a part in changing vulnerability, intensity, resistance, or tolerance to SARSCOV2 infection. Even though the significance of eosinophils during COVID19 sickness is unknown, this has been demonstrated that such a viral disease is linked with significant eosinopenia and that prolonged eosinopenia is related with clinical deterioration and an increased risk of mortality. Various reasons for COVID19-related eosinopenia have been proposed- (36)

According to new recommendations, biological agents should be used throughout the epidemic. Biological drugs do not appear to reduce the immune response to a viral infection, according to the findings. These medicines, when administered in severe cases of asthma, are known to lessen asthma episodes (26-29). In a randomized controlled study of children with allergic asthma aged 9 to 17, it was discovered that omalizumab reduced viral shedding and duration of rhinovirus, the most commonly found virus in asthma attacks (37). During the COVID-19 pandemic, the FDA has approved omalizumab for short-term home administration (38). Biological treatments as home-based administration are encouraged by national recommendations in several nations during pandemics (27,39-41). Clinical trials with omalizumab, mepolizumab, reslizumab, benralizumab, and dupilumab shown that such medications reduced asthma exacerbations while without increasing the risk of viral infection (42-45). During the pandemic, these drugs should be continued in people with severe asthma., based on existing information. The question is whether these medications should be continued as part of COVID-19 treatment. These medicines may be compatible with biologics in severe asthma, although no data is currently available. The decision

to keep a biologic agent or postpone it till a patient recovers should be made on a case-by-case basis with the help of a multidisciplinary council (46).

## Methodology:

To analyze the connection of asthma and Covid 19, utilizing papers of the English language till the June 15, 2020, a systematic research was conducted in the following four digital libraries: PUBMED, MEDLINE, MedRxiv, and Google scholar. "COVID-19" OR "SARS-CoV-2" OR "Coronavirus" AND "Asthma" OR "Allergy" OR "Case series" OR "Comorbidities" OR "Epidemiology" OR "Hospitalization" OR "Risk factors" In addition, the references of the first included research were manually examined to discover other potentially acceptable studies.

**Comment [YT13]:** What do you mean by systematic research? Specify your research methodology. Even the research design is not indicated. Furthermore, the study population is not clarified.

**Comment [YT14]:** Rephrase it to make it more readable.

Following the application of the aforementioned criteria, 45 papers were subjected to a full-text review. Finally, ~~they included~~ 15 papers were included in ~~their~~ the analysis. Within each trial, ~~they utilized~~ the Cloper Pearsons approach was utilized in generating 90 percent confidence intervals ~~C.I (CI confidence intervals)~~ for prevailing cases of asthma. ~~They also used~~ local available information of cases was also used from ~~their~~ hospitals to run a multivariate logistical regression model having numerous inputs to see if having asthma affected intubating chances after managing for gender, age, and Body Mass Index (BMI).

As a result, the proportion of patients with asthma among COVID-19 hospitalised patients is similar to the incidence of asthma in the general community at each research site.

**Comment [YT15]:** Where is the result if it is research? How is possible to discuss with out results presented? If it is a kind of review, you still need to make major amendments on your methodology.

## Result

---

---

## Discussion

According to their data, asthma prevalence among COVID-19 hospitalised patients appeared as were comparable to asthmatic patients and the prevalence were much decreased among influenza hospitalised group. Even after correcting for Body mass index along with age, which is the principal risk factor for severity, asthma is not an independent factor for cause for intubating hospitalised patients having Covid 19.

Primary cause of hospitalization in asthmatic patients with worsening is due to upper respiratory tract infections, coronavirus has no contribution towards this (4). Patient suffering from asthma are not appeared to be largely impacted in instances of SARS-Cov (not the virus that causes Covid 19), despite the lack of data to make this comparison (5, 6).

During the 2019–20 influenza period, 25.2 percent of cases hospitalised having asthma along with flu like symptoms, that was significantly more compared to the four season aggregate of 21% from 2016 till 2020 although, it is significantly larger than the combined presence estimated from the 16 Covid 19 study. (47). In spite of early concerns about asthma patients having unreasonable higher morbidity and death (48), data provided here and elsewhere indicate little indication of a clinically significant link (49,50).

Although after correcting for Body Mass Index and lifespan, that are widely in knowledge risk factors for severe forms and are widely linked with intubation in their design, information through the hospitals may not demonstrate any notable links in asthmatics with increasing intubating odds within COVID-19 cases. The location of the angiotensin Converting enzyme receptor (ACE2) in the respiratory airway epithelium could be one reason why COVID-19 isn't linked to higher hospitalisation rates in people with asthma. Diabetes and hypertension have been linked to increased ACE2 expression, whereas inhaled corticosteroid use has been linked to decreased ACE2 expression, resulting in more difficulty with viral entry (51, 52). Furthermore, patients with asthma, especially those with a primarily allergic phenotype, may have significantly decreased ACE2 expression (52-63). Although the relationship between ACE2 receptor expression and total COVID-19 susceptibility and illness severity is yet unknown, further research is warranted. Unfortunately, we do not have any inhaled-corticosteroid data on patients in our study or at our hospital to further investigate the potential benefits of these drugs, as has been suggested (53). COPD, unlike asthma, increases chances of severe forms of Covid 19 disease in hospitalised cases (54). Higher chances of comorbidity are connected to higher ACE-2 response tissues of the lungs and small airways is linked to this comorbidity (55) Given the varying prevalence of asthma among COVID-19 hospitalised groups, it's probable that comorbidity reporting was uneven across studies, especially since the authors didn't explain how asthma or chronic respiratory illness diagnoses were acquired in any of these investigations. In contrast to the higher-than-expected rates of hypertension and diabetes mellitus among hospitalised patients, which are comorbidities known to be related with severe COVID-19, studies of COPD have revealed prevalence rates that are lower than the population norm. Finally, we realise that our findings could be due to a small sample size, and that more data from asthma and intubation risk investigations would be desirable. Although asthma prevalence varies among COVID-19 publish studies, it appears to be comparable to community prevalence and is certainly lower than what would be expected during seasonal influenza. According to the findings, asthma does not appear to be a substantial risk factor for severe COVID-19 that necessitates hospitalisation or intubation.

### **Conclusions:**

---

According to our analysis of the current data on asthma and COVID19, asthma doesn't really enhance vulnerability to SARS-CoV2 disease or result in a poorer treatment outcome in people with the disease. Asthma doesn't really seem to become a contributing factor for COVID19 in general; nevertheless, its precise role in danger might rely mostly on existence of some of the other behavioral & environmental variables (e.g., smoking, illnesses), as well as the depending upon the severity of asthma (e.g., adherence).

International Asthma Guidelines recommend adhering to basic measures to protect against COVID19 and following asthmatic medications, ICS and biological agents are examples of ICS in asthma patients during the COVID19 pandemic. Some suggested asthmatic patients should manage their existing condition with regulator medicines, especially biologics, because continuous ICS usage doesn't really carry the chance of admission in asthmatic patients with concurrent COVID19 disease. Finally, the choice to prolong or discontinue biological treatment for individuals who have already been infected with SARSCoV2 must be made on an individual basis.

Extensive experimental and clinical research is needed in corroborating such early uncovering of a "dangerous connection" between COVID19 and asthma.

**COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

**References:**

1. Johnston SL, Pattemore PK, Sanderson G, Smith S, Lampe F, Josephs L, Symington P, O'Toole S, Myint SH, Tyrrell DA et al (1995) Community study of role of viral infections in exacerbations of asthma in 9-11 year old children. *BMJ (Clinical research ed)* 310(6989):1225–1229. <https://doi.org/10.1136/bmj.310.6989.1225>
2. Van Bever HP, Chng SY, Goh DY (2004) Childhood severe acute respiratory syndrome, coronavirus infections and asthma. *Pediatr Allergy Immunol* 15(3):206–209. <https://doi.org/10.1111/j.1399-3038.2004.00137.x>
3. The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team (2020) The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) — China, 2020. *China CDC weekly*. February 17 2020
4. Dembic Z (2015) Chapter 6 - cytokines of the immune system: interleukins. In: Dembic Z (ed) *The cytokines of the immune system*. Academic Press, Amsterdam, pp 143–239. <https://doi.org/10.1016/B978-0-12-419998-9.00006-7>
5. Shaker MS, Oppenheimer J, Grayson M, et al. COVID-19: pandemic contingency planning for the allergy and immunology clinic. *J Allergy Clin Immunol Pract* 2020 Mar 26 [Epub ahead of print]. pii: S2213-2198(20)30253-1. doi: 10.1016/j.jaip.2020.03.012
6. Centers for Disease Control and Prevention. Coronavirus disease 2019 (COVID-19) situation summary Available from: <https://www.cdc.gov/coronavirus/2019-ncov/index.html>, Accessed 15<sup>th</sup> Mar 2020

7. Zhang JJ, Dong X, Cao YY, Yuan YD, Yang YB, Yan YQ, et al. Clinical characteristics of 140 patients infected with SARSCoV-2 in Wuhan, China. *Allergy* 2020. Doi: 10.1111/all.14238.
8. Li X, Xu S, Yu M, Wang K, Tao Y, Zhou Y, et al. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *J Allergy Clin Immunol* 2020 pii: S00916749(20)304954.
9. Guan WJ, Liang WH, Zhao Y, Liang HR, Chen ZS, Li YM, et al. Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J* 2020;55(5). Pii: 2000547. Doi: 10.1183/13993003.00547-2020
10. Garg S, Kim L, Whitaker M, O'Halloran A, Cummings C, Holstein R, et al. Hospitalization rates and characteristics of patients hospitalized with laboratory-confirmed coronavirus disease 2019 – COVID-NET, 14 States, March 1-30, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69(15):458-64. Doi: 10.15585/mmwr.mm6915e3.
11. 1001/ jama.2020.6775 10. Emami A, Javanmardi F, Pirbonyeh N, Akbari A. Prevalence of underlying diseases in hospitalized patients with COVID-19: a systematic review and meta-analysis. *Arch Acad Emerg Med* 2020;8(1):e35. eCollection 2020.
12. Williamson E, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, et al. OpenSAFELY: factors associated with COVID-19-related hospital death in the linked electronic health records of 17 million adult NHS patients. COVID19 SARS-CoV-2 preprints from medRxiv and bioRxiv 2020. Doi: 10.1101/2020.05.06.20092999
13. Halpin DMG, Faner R, Sibila O, Badia JR, Agusti A. Do chronic respiratory diseases or their treatment affect the risk of SARS-CoV-2 infection? *Lancet Respir Med* 2020;8(5):436-8.
14. Jackson DJ, Busse WW, Bacharier LB, Kattan M, O'Connor GT, Wood RA, et al. Association of respiratory allergy, asthma, and expression of the SARS-CoV-2 receptor ACE2. *J Allergy Clin Immunol* 2020 pii: S0091-6749(20)30551-0. Doi: 10.1016/j.jaci.2020.04.009
15. Peters MC, Sajuthi S, Deford P, et al. SARS-COV-2 related genes in sputum cells in asthma: relationship to demographic features and corticosteroids. *Am J Respir Crit Care Med* 2020. Doi: 10.1164/rccm.202003-0821OC.
16. Zheng XY, Xu YJ, Guan WJ, Lin LF. Regional, age and respiratory-secretion-specific prevalence of respiratory viruses associated with asthma exacerbation: a literature review. *Arch Virol* 2018;163(4):845-53.
17. Gonzales-van Horn SR, Farrar JD (2015) Interferon at the crossroads of allergy and viral infections. *J Leukoc Biol* 98(2):185–194. <https://doi.org/10.1189/jlb.3RU0315-099R>

18. Zhu J, Message SD, Mallia P, Keadze T, Contoli M, Ward CK, Barnathan ES, Mascelli MA, Kon OM, Papi A, Stanciu LA, Edwards MR, Jeffery PK, Johnston SL (2019) Bronchial mucosal IFN-alpha/beta and pattern recognition receptor expression in patients with experimental rhinovirus-induced asthma exacerbations. *J Allergy Clin Immunol* 143(1):114–125.e114. <https://doi.org/10.1016/j.jaci.2018.04.003>
19. Rosenberg HF, Dyer KD, Domachowske JB (2009) Eosinophils and their interactions with respiratory virus pathogens. *Immunol Res* 43(1–3):128–137. <https://doi.org/10.1007/s12026-008-8058-5>
20. Avdeev S, Moiseev S, Brovko M, Yavorovskiy A, Umbetova K, Akulkina L, et al. Low prevalence of bronchial asthma and chronic obstructive lung disease among intensive care unit patients with COVID-19. *Allergy*. 2020;75(10):2703–4. <https://doi.org/10.1111/all.14420>.
21. Yang JM, Koh HY, Moon SY, Yoo IK, Ha EK, You S, et al. Allergic disorders and susceptibility to and severity of COVID-19: a nationwide cohort study. *J Allergy Clin Immunol*. 2020;146(4):790–8. <https://doi.org/10.1016/j.jaci.2020.08.008>.
22. Coronavirus 19. Center for Disease Control and Prevention. <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/asthma.html>. Updated Jan. 20, 2021.
23. Caminati M, Vultaggio A, Matucci A, Senna G, Almerigogna F, Bagnasco D, et al. Asthma in a large COVID-19 cohort: prevalence, features, and determinants of COVID-19 disease severity. *Respir Med*. 2021;176:106261. <https://doi.org/10.1016/j.rmed.2020.106261>.
24. Eger K, Bel EH. Asthma and COVID-19: do we finally have answers? *Eur Respir J*. 2020 Dec;30:2004451.
25. Assaf SM, Tarasevych SP, Diamant Z, Hanania NA. Asthma and severe acute respiratory syndrome coronavirus 2019: current evidence and knowledge gaps. *Curr Opin Pulm Med*. 2021;27:45–53.
26. Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention. Available from: [www. Ginaasthma.org](http://www.ginaasthma.org)
27. COVID-19 rapid guideline: severe asthma. NICE guideline [NG166]Published date: 03 April 2020.
28. Licskaia C, Yang CL, Ducharme FM, Radhakrishnan D, Podgerse D, Ramsey C, et al. Addressing therapeutic questions to help Canadian physicians optimize asthma management for their patients during the COVID-19 pandemic. *Canadian J Respir Crit Sleep Med* Available from: <https://doi.org/10.1080/24745332.2020.1754027>

29. Shaker MS, Oppenheimer J, Grayson M, Stukus D, Hartog N, Hsieh EWY, et al. COVID-19: pandemic contingency planning for the allergy and immunology clinic. *J Allergy Clin Immunol Pract* 2020;8(5):1477-88.e5.
30. Ari A. Practical strategies for a safe and effective delivery of aerosolized medications to patients with COVID-19. *Respir Med* 2020:105987. Doi: 10.1016/j.rmed.2020.105987
31. McGrath J, O'Toole C, Joyce GBM, Byrne M, MacLoughlin R. Investigation of fugitive aerosols released into the environment during high-flow therapy. *Pharmaceutics* 2019;11:254.
32. Wittgen BP, Kunst PW, Perkins WR, Lee JK, Postmus PE. Assessing a system to capture stray aerosol during inhalation of nebulized liposomal cisplatin. *J Aerosol Med* 2006;19:385-91.
33. Shang L, Zhao J, Hu Y, Du R, Cao B. On the use of corticosteroids for 2019-nCoV pneumonia. *Lancet* 2020;395(10225):683-4.
34. Russell CD, Millar JE, Baillie JK. Clinical evidence does not support corticosteroid treatment for 2019-nCoV lung injury. *Lancet* 2020 Feb 15;395(10223):473-5.
35. Levin M, Morais-Almeida M, Ansotegui IJ, Bernstein J, Chang YS, Chikhladze M, et al. Acute asthma management during SARS-CoV2-pandemic 2020. *World Allergy Organ J* 2020 May 14:100125. Doi: 10.1016/j.waojou.2020.100125.
36. Roca E, Ventura L, Zattra CM, C Lombardi. EOSINOPENIA: an early, effective, and relevant COVID-19 biomarker? *QJM*. 2021;114(1):68-9. <https://doi.org/10.1093/qjmed/hcaa259>.
37. Esquivel A, Busse WW, Calatroni A, Togias AG, Grindle KG, Bochkov YA, et al. Effects of omalizumab on rhinovirus infections, illnesses, and exacerbations of asthma. *Am J Respir Crit Care Med* 2017;196(8):985-92.
38. Self-Administration of Xolair Pre-Filled Syringe (PFS) for Asthma during the COVID-19 Pandemic. <https://www.gene.com/medical-professionals/medicines/xolair>. Accessed 23-Apr-2020
39. <http://www.siaaic.org/wp-content/uploads/2020/04/SIAAIC.pdf>. Accessed 8-Apr-2020. Documento-di-Indirizzo-
40. Malipiero G, Paoletti G, Puggioni F, Racca F, Ferri S, Marsala A, et al. An academic allergy unit during COVID-19 pandemic in Italy. *J Allergy Clin Immunol* 2020 Apr 16:S0091-6749(20)30489-9. Doi: 10.1016/j.jaci.2020.04.003
41. Mitchell P, Leigh R. A drug safety review of treating eosinophilic asthma with monoclonal antibodies. *Expert Opin Drug Saf* 2019;18(12):1161-70.

42. Agache I, Rocha C, Beltran J, et al. Efficacy and safety of treatment with biologicals (benralizumab, dupilumab and omalizumab) for severe allergic asthma. *Allergy* 2020 Feb 16. Doi: 10.1111/all.14235.
43. Pavord ID, Korn S, Howarth P, et al. Mepolizumab for severe eosinophilic asthma (DREAM): a multicentre, double-blind, placebo-controlled trial. *Lancet* 2012;380(9842):651-9.
44. FitzGerald JM, Bleecker ER, Nair P, et al; CALIMA study investigators. Benralizumab, an anti-interleukin-5 receptor  $\alpha$  monoclonal antibody, as add-on treatment for patients with severe, uncontrolled, eosinophilic asthma (CALIMA): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet* 2016;388(10056):2128-41.
45. Castro M, Corren J, Pavord ID, et al. Dupilumab efficacy and safety in moderate-to-severe uncontrolled asthma. *N Engl J Med* 2018;378(26):2486-96.
46. Morais-Almeida M, Aguiar R, Martin B, Ansotegui IJ, Ebisawa M, Arruda LK, et al. COVID-19, asthma, and biologic therapies: What we need to know. *World Allergy Organ J* 2020 May 16:100126. Doi: 10.1016/j.waojou.2020.100126.
47. U.S. Centers for Disease Control and Prevention. FluView interactive: laboratory-confirmed influenza hospitalizations. Atlanta, GA: U.S. centers for Disease Control and Prevention; 2020 [accessed 2020 May 7]. Available from: <https://gis.cdc.gov/grasp/fluview/FluHospChars.html>.
48. Johnston SL. Asthma and COVID-19: is asthma a risk factor for severe outcomes? *Allergy* 2020;75:1543–1545.
49. Razzaghi H, Wang Y, Lu H, Marshall KE, Dowling NF, Paz-Bailey G, Et al. Estimated county-level prevalence of selected underlying medical conditions associated with increased risk for severe COVID-19 illness: United States, 2018. *MMWR Morb Mortal Wkly Rep* 2020; 69:945–950.
50. Zhu Z, Hasegawa K, Ma B, Fujiogi M, Camargo CA, Liang L. Association of asthma and its genetic predisposition with the risk of severe COVID-19. *J Allergy Clin Immunol* 2020;146:327–329, e4.
51. Peters MC, Sajuthi S, Deford P, Christenson S, Rios CL, Montgomery MT, et al. COVID-19-related genes in sputum cells in asthma: relationship to demographic features and corticosteroids. *Am J Respir Crit Care Med* 2020;202:83–90.
52. Jackson DJ, Busse WW, Bacharier LB, Kattan M, O'Connor GT, Wood RA, et al. Association of respiratory allergy, asthma, and expression of the SARS-CoV-2 receptor ACE2. *J Allergy Clin Immunol* 2020;146: 203–206, e3.

53. Halpin DMG, Faner R, Sibila O, Badia JR, Agusti A. Do chronic respiratory diseases or their treatment affect the risk of SARS-CoV-2 infection? *Lancet Respir Med* 2020;8:436–438.
54. Guan W-J, Liang W-H, Zhao Y, Liang HR, Chen ZS, Li YM, et al.; China medical Treatment Expert Group for COVID-19. Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J* 2020;55:2000547.
55. Leung JM, Yang CX, Tam A, Shaipanich T, Hackett TL, Singhera GK, Et al. ACE-2 expression in the small airway epithelia of smokers and COPD patients: implications for COVID-19. *Eur Respir J* 2020;55: 2000688.
56. Acharya, Sourya, Samarth Shukla, and Neema Acharya. “Gospels of a Pandemic- A Metaphysical Commentary on the Current COVID-19 Crisis.” *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH* 14, no. 6 (June 2020): OA01–2. <https://doi.org/10.7860/JCDR/2020/44627.13774>.
57. Arora, Devamsh, Muskan Sharma, Sourya Acharya, Samarth Shukla, and Neema Acharya. “India in ‘Flattening the Curve’ of COVID-19 Pandemic - Triumphs and Challenges Thereof.” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 43 (October 26, 2020): 3252–55. <https://doi.org/10.14260/jemds/2020/713>.
58. Bawiskar, Nipun, Amol Andhale, Vidyashree Hulkoti, Sourya Acharya, and Samarth Shukla. “Haematological Manifestations of Covid-19 and Emerging Immunohaematological Therapeutic Strategies.” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 46 (November 16, 2020): 3489–94. <https://doi.org/10.14260/jemds/2020/763>.
59. Burhani, Tasneem Sajjad, and Waqar M. Naqvi. “Telehealth - A Boon in the Time of COVID 19 Outbreak.” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 29 (July 20, 2020): 2081–84. <https://doi.org/10.14260/jemds/2020/454>.
60. Butola, Lata Kanyal, Ranjit Ambad, Prakash Kesharao Kute, Roshan Kumar Jha, and Amol Dattarao Shinde. “The Pandemic of 21st Century - COVID-19.” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 39 (September 28, 2020): 2913–18. <https://doi.org/10.14260/jemds/2020/637>.
61. Dasari, Venkatesh, and Kiran Dasari. “Nutraceuticals to Support Immunity: COVID-19 Pandemic- A Wake-up Call.” *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH* 14, no. 7 (July 2020): OE05–9. <https://doi.org/10.7860/JCDR/2020/44898.13843>.
62. Dhok, Archana, Lata Kanyal Butola, Ashish Anjankar, Amol Datta Rao Shinde, Prakash Kesharao Kute, and Roshan Kumar Jha. “Role of Vitamins and Minerals in Improving Immunity during Covid-19 Pandemic - A Review.” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 32 (August 10, 2020): 2296–2300. <https://doi.org/10.14260/jemds/2020/497>.
63. Gawai, Jaya Pranoykumar, Seema Singh, Vaishali Deoraaji Taksande, Tessy Sebastian, Pooja Kasturkar, and Ruchira Shrikant Ankar. “Critical Review on Impact of COVID 19 and Mental Health.” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL*

SCIENCES-JEMDS 9, no. 30 (July 27, 2020): 2158–63.  
<https://doi.org/10.14260/jemds/2020/470>.

UNDER PEER REVIEW

