

Review Article

Blood Coagulation: A Potentially Lethal Manifestation in Covid-19 Patients

ABSTRACT:

Blood clotting is a well-known phenomenon which is intended to provide protection in case of any external cellular/ tissue injury. However, there lies a concern due to this phenomenon in Covid-19 patients that is leading to unusual thrombotic presentations. Since the emergence of novel ~~coronavirusecorona~~ coronavirus, World has come across various benign as well as lethal manifestations in COVID-19 patients and one such life-threatening manifestation which needs rigorous attention is the genesis of strange blood clots that can travel and get lodged into several parts of the COVID-19 patients leading to various clinical presentations. COVID-19 infection is caused due to interaction of spike glycoproteins of ~~coronavirusecorona~~ coronavirus with ACE 2 receptors ~~on the hostof host~~ cell surface. This interaction in the arteries, veins or capillaries could lead to injury ~~in the vascularin vascular~~ wall of blood vessels that can directly/Indirectly lead to coagulation and clotting cascades activation and subsequent formation of internal blood clots. However, it is undesirable to have the presence of these clots as they could lead to certain fatal complications that need emergency medical intervention or else, they may lead ~~to the deathto death of a patientof patient~~. Due to the severity of this manifestation, early detection of these blood clots in covid-19 patients become very important intervention which could be done by observing certain specific signs and symptoms and/or with the help of various laboratory biomarkers like D-dimer, platelet count, erythrocyte sedimentation rate (ESR), ferritin, fibrinogen, etc. Once the early diagnosis is made, ~~the patientpatient~~ can be treated appropriately with the help of anticoagulant therapy, which includes use of oral and parenteral anticoagulant drugs. This way the complications of blood coagulation like thrombo-embolism, could be prevented.

KEYWORDS:

- Covid-19
- Blood Clotting

Comment [1]: A lot works have left to be done:
The author of this manuscript have to restructure the review section of Abstract. The objective, the reason why this review is required, Review methods, conclusions sections are missed. What types of documents were included and excluded are not mentioned enough.

Comment [2]: This statement is not clear, please rewrite

- ACE 2 receptors
- Anticoagulants.

Comment [3]: No need of bullet, put alphabetical order and remove bulet

BACKGROUND:

Severe acute respiratory syndrome coronavirus-2, designated as SARS-CoV-2 first recognised in the middle of an outbreak in Wuhan City, Hubei Province, China, ~~is;~~ ~~is~~ the causative organism of Coronavirus disease 2019 (COVID-19). On 31th December ~~in;~~ 2019, it was first conveyed to the World Health Organization (WHO).

Comment [4]: Reference is required here

The World Health Organisation announced this ~~coronaviruseorona-virus~~ outbreak as a global health crisis on 30th of January 2020. Then in the same year on 11th of March, the World Health Organisation declared COVID-19 a global pandemic. Since then, over the course of time with the advent of new mutated ~~coronaviruseorona-virus~~ strains, dynamics of COVID-19 have kept on evolving.

Comment [5]: What this paragraph talk about?? Not clear. Please write it and insert the references

Comment [6]: Reference?????

Comment [7]: Reference

SARS-CoV-2 is a virus composed of a single strand of Ribonucleic acid (RNA), glycoprotein, spike protein and an ~~envelope~~envelop. It is a virus of the pulmonary system and so when it enters in the human body, it primarily infects the predominant organ of pulmonary system which is the lung.[1] This access is facilitated by the linking of the receptor-binding domain (RBD) of S1 subunit present on the spike protein of virus with the angiotensin-converting enzyme-2 (ACE-2) receptor of the host which are principally articulated in type II pneumocytes, serving as a pool for the virus.[2][3]

Comment [8]: Move this paragraph to next to the first paragraph of background

A major proportion of individuals infected with COVID-19 remain asymptomatic and recover without requiring any medical intervention.]

Comment [9]: Discuss about current status of COVID 19 at least in your country.

However, common clinical presentations in symptomatic patients include dry cough, sore throat, fever and breathlessness.[4] Although a variety of other symptoms have been seen in patients infected with different mutated strains of ~~coronaviruse~~corona virus over the course of ~~the pandemic~~of pandemic.

Blood clots have been discovered in the minute blood vessels of various organs like lungs, heart, liver, and kidney, on ~~post-mortem~~post-mortem examinations of a significant percentage of COVID-19 patients. These blood clots can cause ischemia of the tissues or organs in which they are logged ~~-[5].~~ Thus, this aspect of covid-19 disease has become an area of research and requires adequate concern.

Comment [10]: This is not enough, more justification why this review required should be needed here.

Comment [11]: Who attacked more by COVID 19?? State 1 - 2 paragraph about severity of COVID 19 by sex, age, human health status and others.

OBJECTIVES:

The main objective of this review article is to collect, summarize and present all the important data and information in regard to blood coagulation which has come out to be a most important hematological appearance in patients suffering from covid-19 disease. ~~This article comprises the~~comprises of the complete story of blood coagulation in covid-19 patients ~~under the following~~under following heads:

- Introduction
- Pathophysiology
- Clinical picture
- Laboratory findings
- Treatment

Comment [12]: What is this??? Is this objective of the current review?? Move to its appropriate place.

INTRODUCTION:

The pandemic of Coronavirus disease 2019 (COVID-19) has been producing substantial mortality and morbidity since its outbreak in December 2019. COVID-19 has been linked up with various coagulative disorders and hypercoagulable states in a growing number of studies. Coronavirus infection could trigger numerous systemic coagulative and inflammatory reactions, as per the hypothesized mechanism. Increased production of proinflammatory cytokine is a result of host inflammatory responses, which triggers coagulation cascade activation and consumptive coagulopathy.[6] When compared to historical data, numerous observational studies have found a greater prevalence of venous thrombotic events amid the corona virus infected individuals under intensive care unit (ICU) admission.[7] However, some studies have also testified thrombotic events in arteries of individuals infected with corona virus.[8] Experts urge anticoagulant prophylaxis for all ~~coronaviruseorona virus~~ infected patients who are seriously unwell.

Comment [13]: Move this to appropriate places in background before justification of review

METHODOLOGY:

This is a Narrative review article which has been prepared after going through various literature works from sources such as PubMed and Google scholar.

Comment [14]: What were/was your inclusion and exclusion criterias, by time, place and types?? Please add all thes information here.

PATHOPHYSIOLOGY:

After primary infection, the virus moves into the systemic circulation and attaches itself to the Angiotensin-converting enzyme-2 (ACE-2) expressing endothelial cells that ~~formforms~~ the lining of the blood vessels. This linkage aids the virus, resulting in injury to endothelial cells and harm to the surface of blood vasculature. The injury to the vasculature can also happen because of inflammation in endothelial cells that can lead to cytokine release and subsequent blood vessel injury. Following injury to endothelial cells in blood vessels, recruitment and aggregation of platelets together with collagen takes place at the place of injury. This aggregation is enhanced ~~by the release~~ ~~by-release~~ and action of the Von Willebrand factor. ~~CoagulationSoon-coagulation~~ is initiated by activation ~~of the~~ ~~coagulationof-coagulation~~ cascade following the intrinsic pathway. The genesis of clots in blood vasculature can be initiated following sequential activation of all the twelve clotting factors, followed by conversion of prothrombin to form thrombin and then conversion of fibrinogen to form insoluble and elongated fibrins. These fibrins ~~cross-linker~~ ~~cross-link~~ with platelets by covalent bonds which results in formation of a firm interlocking network of fibrin

~~clot~~ ~~at the location~~ ~~at location~~ of injury. This leads to formation of haemostatic plugs. This haemostatic plug can lead to various complications, of which thrombosis is one of the most commonly perceived ~~complication~~ ~~complication~~ among Covid-19 individuals.

Virchow described three primary events which predispose to thrombus formation and are collectively known as Virchow's triad. It includes: -

❖ **Endothelium injury:**

It triggers platelet activation and genesis of thrombus by the action of von Willebrand factor (vWF) and tissue factor in the heart and in the arterial circulation, where coagulation is impeded by high rates of blood flow. Activation and adherence of platelets is an important ~~prerequisite~~ ~~pre-requisite~~ for formation of thrombus and therefore clots in arteries and heart are exclusively rich in platelets. Hence, in cases of coronary artery disease and acute myocardial infarction, platelet-inhibitors like aspirin, etc are brought in use. There are several prothrombotic events that can lead to what is called endothelial activation or dysfunction. These events include varied exposures to infectious agents, physical injury, altered blood flow, mediators of inflammation, abnormalities of metabolism such as hypercholesterolemia or ~~homocysteinemia~~ ~~homocystinemia~~, and toxins assimilated from cigarette smoke.

Following are some important prothrombotic alterations: -

- **Procoagulant changes** - On activation by cytokines, endothelial cells reduce the thrombomodulin expression, which is a main regulator of thrombin activity. This can cause continuous thrombin activation, causing stimulation of platelets and augmentation of inflammation through Protease-activated receptors (PARs) expressed over platelets and other inflammatory cells. Along with this, the inflamed endothelium also reduces the activity of anticoagulants like protein C and tissue factor protein inhibitors.

- **Antifibrinolytic effects** - Cells of endothelium on activation release plasminogen activator inhibitors (PAIs) ~~that~~ ~~that~~ limit lysis of fibrins.

❖ **Altered blood flow (Stasis or turbulent):**

Normally, blood flow in vessels is laminar in ~~such~~ ~~aa-such~~ way that the platelets and other cellular elements flow in the centre in the lumen of ~~vessels~~ ~~vessel~~, around this central flow of cellular elements is the layer of plasma which moves slowly and separates it from the endothelium. Endothelial injury is caused by turbulent flow

Comment [15]: Be consistency throughout the documents, no need of putting bullet

which leads to formation of counter-currents that contribute to cardiac and arterial thrombosis. Whereas, ~~the major~~ cause behind formation of venous thrombi is stasis.

- Abnormal flow of blood promotes endothelial activation, which enhances procoagulant activity and leukocyte adhesion.
- It disturbs laminar flow and platelets come in contact with endothelium.
- It also safeguards the removal of active clotting factors by and inhibits the inflow of inhibitors of clotting factors.

Altered blood flow aids thrombosis in several clinical syndromes such as aneurysms, acute myocardial infarctions, rheumatic mitral valve stenosis, polycythaemia vera, sickle cell anaemia, etc.

❖ ***Hypercoagulability of blood:***

~~Hypercoagulability, also~~ ~~Hypercoagulability~~ also known as ~~thrombophilia, is defined~~ ~~thrombophilia is defined~~ as any haematological disorder predisposing to thrombosis. Hypercoagulability particularly contributes ~~to thrombosis~~ ~~in thrombosis~~ of veins. It can be classified as primary disorders that are genetic and secondary disorders that are acquired. The most common events leading to thrombophilia are point mutations in the prothrombin gene and factor V gene.

- One of the mutations leading to hypercoagulability is mutation of a single nucleotide in factor V, ~~called factor~~ ~~referred as factor~~ V Leiden. This mutation leads ~~to substitution~~ ~~in substitution~~ of glutamine by arginine at 506 amino acid residue that results in factor V becoming resistant to splitting and deactivation by protein C, resulting in loss of an antithrombotic counterregulatory pathway.
- Another single nucleotide mutation in the prothrombin gene ~~at the 3'-at 3'~~ untranslated region can be linked ~~to up with~~ hypercoagulability. It leads to raised levels of prothrombin and a threefold increase in venous thrombosis.
- Apart from above mentioned genetic factors, raised homocysteine levels also aid in thrombosis of arteries and veins. Increased levels of homocysteine could be due to an inherited deficiency of ~~cystathionine~~ ~~cystathione~~ β -~~synthase-~~ ~~synthetase~~. [9][10]

CLINICAL PICTURE:

In general, the location of blood clot in the body determines the clinical features a patient suffers:

- A blood clot in heart or lung could cause-
 - i. Chest pain
 - ii. Shortness of breath
 - iii. Upper body discomfort in the arms, back, neck, or jaw
- A blood clot ~~in the brain~~ in brain could cause-
 - i. Headache
 - ii. Paralysis
 - iii. Dizziness
 - iv. Trouble in speaking or understanding speech
- A blood clot in deep veins of the leg could cause-
 - i. Pain
 - ii. Redness
 - iii. Warmth
 - iv. Swelling

Corona positive individuals can come up with a variety of abnormalities of coagulation as well as complications of thrombosis, some of which are as follows:

- Pulmonary embolism (PE)
- Venous thromboembolism (VTE)
- Sepsis-induced coagulopathy (SIC)
- Disseminated intravascular coagulation (DIC)
- Thrombotic ~~microangiopathy~~ micro-angiopathy
- Microvascular thrombosis
- Micro-thrombosis

LABORATORY FINDINGS:

Following are some of the potentially useful Laboratory findings in COVID-19: -

- **Tests in which levels are likely to increase:**
 - C-reactive protein (CRP)

- D-dimer
- Activated partial thromboplastin (aPTT)
- Aspartate aminotransferase (AST)
- Alanine aminotransferase (ALT)
- Lactate dehydrogenase (LDH)
- Fibrinogen
- Prothrombin time (PT)
- CBC (platelets and lymphocytes)

The levels of above-mentioned markers are usually expected to increase in the acute ~~in~~ acute stage of the disease.

Comment [16]: Who said all this?References are missed???

Formatted: Indent: Left: 0"

□ **Tests in which levels are likely to decrease:**

- Albumin
- Activated partial thromboplastin (aPTT)
- CBC (platelets and lymphocytes)
- Fibrinogen
- Prothrombin time (PT)

The levels of above-mentioned markers are usually expected to decrease in the late ~~in~~ late stage of disease.

There are specific laboratory markers which are helpful for diagnosis, treatment and prognosis of blood coagulation and its related disorders in patients with COVID-19. These include: -

- **D-dimer:** D-dimer is an outcome of denaturation of fibrin. It is one of the fragments of protein formed on termination of a blood clot in the body ~~in~~ body. In normal circumstances, it is either undetectable or detectable at a very low level. It is detected only when there is formation and breakdown of blood clots in the body. It's presence in blood indicates fibrinolysis.[11] In patients with COVID-19, raised D-dimer readings are observed to be associated with bad prognosis and increased mortality. In patients with COVID-19, Systemic Inflammatory Response Syndrome (SIRS)

~~leads~~lead to coagulation cascade activation which ultimately is responsible for such high D-Dimer levels.[12]

- **Platelet count:** Platelets are also designated as thrombocytes. These are colourless bits of blood cells that prevent or stop bleeding by forming clots. Thrombocytopenia is detected in ~~corona-positive~~~~corona-positive~~ patients but the incidence varies as per the severity of disease. Severely diseased patients have a platelet count ranging between $23 \times 10^9/L$ to $31 \times 10^9/L$ on average, which is lower as compared to those with non-severe disease. Platelets are involved in inflammatory signals as well as the immune response. Platelets may assist target haemostasis and immunological responses against possible infectious agents to limit microbial invasion by combining thrombotic and immune recruiting roles.
- **C-reactive protein (CRP):** It is considered to be an "acute stage protein"~~that,~~~~that~~ is an early marker of any infectious or inflammatory conditions. Normally, the concentration of CRP in blood is less than 10 mg/L. Although within 6 to 8 hours, it rapidly increases and peaks within 48 hours from the disease onset.[13] Half- life of C-reactive protein is about 19 hours and when the inflammatory stage ends, its concentration decreases.
- **Erythrocyte sedimentation rate (ESR):** It is a procedure or method to evaluate blood by measuring the speed of sinking of erythrocytes (red blood cells) at the lowermost part of a test tube. Raised ESR can lead to various joint abnormalities like osteoarthritis, it can also be seen as a precursor of renal and hepatic ~~dysfunction;~~~~therefore~~~~dysfunction~~ ~~therefore~~ raised ESR in individuals with COVID-19 could be a sign of bad prognosis.
- **Fibrinogen:** Fibrinogen is a complex of glycoprotein that is synthesized ~~in the liver~~~~in~~ ~~liver~~. It is enzymatically converted by thrombin into fibrin as a consequence of any tissue or vascular injury. [14] The fibrin clots so formed primarily ~~act~~~~acts~~ to occlude blood vessels in order to stop bleeding. Several studies have found remarkably higher fibrinogen levels in critically diseased patients.
- **Ferritin:** It is an intracellular iron-storage protein. Various studies have suggested elevated levels of serum ferritin in individuals with severe COVID-19 disease.[15] In autopsies of 12 patients with COVID-19 as a cause of death, raised levels of serum ferritin were found.[16]

- **Procalcitonin:** Procalcitonin is the precursor of calcitonin, which regulates calcium~~regulates the calcium~~ homeostasis. preprocalcitonin is cleaved by endopeptidase to form procalcitonin. The level of procalcitonin can reflect the severity of disease in corona positive patients.

Formatted: Font color: Auto

Formatted: Space After: 0 pt

TREATMENT:

The treatment protocols for Covid-19 disease have kept on changing with the course of the pandemic~~of pandemic~~. In general, the treatment for covid-19 is definite and is based on the severity of disease and presence or absence of some risk factors. Currently, a wide range of therapeutic modalities are available for management of COVID-19 under Emergency Use Authorization (EUA) issued by the Food~~by Food~~ and Drug Administration (FDA). These include: -

- Anti-inflammatory drugs
- Anti-SARS-CoV-2 monoclonal antibodies
- Antiviral drugs
- Immunomodulators agents [17]

Anticoagulant drugs are usually used to treat blood clots. These drugs~~These group of drugs~~ not only prevent existing clots from getting bigger but they also keep a check on new clot formation. One observational study examined the effects of anticoagulants in people who were hospitalized with COVID-19. It found that people who were treated with anticoagulants in the hospital had a more positive outcome than those who were not.[18] Therefore, WHO recommends the use of low dose anticoagulants in hospitalized patients for preventing the formation of blood~~the blood clots in~~ clots formation in blood vessels.

Chart 1. Parenteral anticoagulants: -

Comment [17]: Say it table, not chart (not appropriate)

INDIRECT THROMBIN INHIBITORS	DIRECT THROMBIN INHIBITORS
Heparin (unfractionated)	Bivalirudin
Low molecular weight heparins (Enoxaparin, Reviparin, Nadroparin,	Argatroban

Ardeparin, Parnaparin)	
Fondaparinux	
Danaparoid	

Chart 2. Oral anticoagulants: -

VIT K ANTAGONISTS	DIRECT FACTOR Xa INHIBITOR	ORAL DIRECT THROMBIN INHIBITOR
Bishydroxycoumarin (Dicumarol)	Rivaroxaban	Dabigatran-etexilate
Warfarin sod.	Apixaban	
Acenocoumarol (Nicoumalone)		
Ethyl-biscoum-acetate		

[19-25]

There are some measures which are certainly helpful in preventing blood coagulation and associated disorders. These are as follows:

- Stay active- A sedentary lifestyle can increase the risk of developing blood clots therefore **being been** fit and active is very important for which regular exercises are very important.
- Lose weight- Shedding excess weight can help lower your risk for developing blood clots.
- Quit smoking- Smoking can harm the blood vessel lining and lead to formation of blood clots.

● **Impacts of COVID 19**

Prevention and Control Methods

CONCLUSION:

Comment [18]: Discuss about economic and social impacts of COVID 19 to make the document more better

Formatted: Font: Bold, Font color: Auto

Formatted: Font: Bold, Font color: Auto

Comment [19]: What prevention and Control methods can be used?? You have to Mention

Formatted: Space Before: 0 pt, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

Formatted: Font: (Default) Arial, 11 pt, No underline

Since the time ~~of the emergence of~~ ~~emergence~~ of ~~novel noble~~ ~~coronaviruses~~ ~~corona virus~~, scientists as well as medical professionals around the world have made a lot of progress in unfolding facts related to various aspects of the virus. Based on this study, we can state that blood coagulation is indeed a very lethal manifestation of COVID-19 disease. It occurs as a consequence of endothelium injury which can be due to binding of spike proteins with ACE 2 receptors or due to release of Inflammatory cytokines. This leads to formation of blood coagulants that can ultimately cause various abnormalities of coagulation and complications associated with thrombosis. Although timely diagnosis based on various signs and symptoms, and with the help of various laboratory markers; followed by timely medical intervention using anticoagulant therapy can prevent the patient from entering the state of critical condition or even death. |

Comment [20]: Add more and What do you recommend standing from your review finding in addition to what you already mentioned

———— There ~~is a lot is lot~~ of scope and requirement of research in areas related to COVID-19 as the corona virus is there to stay with us for a long time and we have to find ways to combat its lethal effects on human beings.

Prevention and Control Methods |

Comment [21]: What prevention and Control methods can be used?? You have to Mention

REFERENCES:

1. Jain A. COVID-19 and lung pathology. Indian J Pathol Microbiol. 2020;63(2):171.
2. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. Cell. 2020;181(2):271-280.e8.
3. Lin L, Lu L, Cao W, Li T. Hypothesis for potential pathogenesis of SARS-CoV-2 infection—a review of immune changes in patients with viral pneumonia. Emerg Microbes Infect. 2020;9(1):727–32.
4. Kakodkar P, Kaka N, Baig MN. A comprehensive literature review on the clinical presentation, and management of the pandemic Coronavirus disease 2019 (COVID-19). Cureus [Internet]. 2020; Available from: <http://dx.doi.org/10.7759/cureus.7560>
5. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. J Thromb Haemost. 2020;18(4):844–7.

6. Connors JM, Levy JH. COVID-19 and its implications for thrombosis and anticoagulation. *Blood*. 2020;135(23):2033–40.
7. Helms J, Tacquard C, Severac F, Leonard-Lorant I, Ohana M, Delabranche X, et al. High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicentre prospective cohort study. *Intensive Care Med*. 2020;46(6):1089–98.
8. Mao L, Jin H, Wang M, Hu Y, Chen S, He Q, et al. Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol*. 2020; 10:10.
9. Kumar V, Abbas AK, Fausto N, Aster JC. Robbins and Cotran pathologic basis of disease, professional edition ~~Ebook~~~~E-book~~. 8th ed. Saunders; 2009.
10. Mohan H. Textbook of pathology. 8th ed. New Delhi, India: Jaypee Brothers Medical; 2018.
11. Demelo-Rodríguez P, Cervilla-Muñoz E, Ordieres-Ortega L, Parra-Virto A, Toledano-Macías M, Toledo-Samaniego N, et al. Incidence of asymptomatic deep vein thrombosis in patients with COVID-19 pneumonia and elevated D-dimer levels. *Thromb Res*. 2020;192:23–6.
12. Léonard-Lorant I, Delabranche X, Séverac F, Helms J, Puzet C, Collange O, Schneider F, Labani A, Bilbault P, Molière S, Leyendecker P. Acute pulmonary embolism in patients with COVID-19 at CT angiography and relationship to D-dimer levels. *Radiology*. 2020 Sep;296(3):E189-91.
13. Young B, Gleeson M, Cripps AW. C-reactive protein: a critical review. *Pathology*. 1991;23(2):118–24.
14. Demelo-Rodríguez P, Cervilla-Muñoz E, Ordieres-Ortega L, Parra-Virto A, Toledano-Macías M, Toledo-Samaniego N, García-García A, García-Fernández-Bravo I, Ji Z, de-Miguel-Diez J, Álvarez-Sala-Walther LA. Incidence of asymptomatic deep vein thrombosis in patients with COVID-19 pneumonia and elevated D-dimer levels. *Thrombosis research*. 2020 Aug 1; 192:23-6.
15. Zhou B, She J, Wang Y, Ma X. Utility of ferritin, procalcitonin, and C-reactive protein in severe patients with 2019 Novel Coronavirus disease [Internet]. Research Square. 2020. Available from: <http://dx.doi.org/10.21203/rs.3.rs-18079/v1>
16. Fox SE, Akmatbekov A, Harbert JL, Li G, Brown JQ, Heide RSV. Pulmonary and cardiac pathology in covid-19: The first autopsy series from New Orleans [Internet]. 2020. Available from: <http://dx.doi.org/10.1101/2020.04.06.20050575>

17. Coopersmith CM, Antonelli M, Bauer SR, Deutschman CS, Evans LE, Ferrer R, et al. The surviving sepsis campaign: Research priorities for Coronavirus disease 2019 in critical illness: Research priorities for Coronavirus disease 2019 in critical illness. *Crit Care Med*. 2021;49(4):598–622.
18. Bs IP, V F, A L, Md AR, Md Z, Gn N. Association of Treatment Dose Anticoagulation with In-Hospital Survival Among Hospitalized Patients with COVID-19.
19. Tripathi KD. *Essentials of medical pharmacology*. 8th ed. New Delhi, India: Jaypee Brothers Medical; 2018.
20. Acharya, Sourya, Samarth Shukla, and Neema Acharya. “Gospels of a Pandemic- A Metaphysical Commentary on the Current COVID-19 Crisis.” *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH* 14, no. 6 (June 2020): OA01–2. <https://doi.org/10.7860/JCDR/2020/44627.13774>.
21. Arora, Devamsh, Muskan Sharma, Sourya Acharya, Samarth Shukla, and Neema Acharya. “India in ‘Flattening the Curve’ of COVID-19 Pandemic - Triumphs and Challenges Thereof.” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 43 (October 26, 2020): 3252–55. <https://doi.org/10.14260/jemds/2020/713>.
22. Bawiskar, Nipun, Amol Andhale, Vidyashree Hulkoti, Sourya Acharya, and Samarth Shukla. “Haematological Manifestations of Covid-19 and Emerging Immunohaematological Therapeutic Strategies.” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 46 (November 16, 2020): 3489–94. <https://doi.org/10.14260/jemds/2020/763>.
23. Burhani, Tasneem Sajjad, and Waqar M. Naqvi. “Telehealth - A Boon in the Time of COVID 19 Outbreak.” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 29 (July 20, 2020): 2081–84. <https://doi.org/10.14260/jemds/2020/454>.
24. Butola, Lata Kanyal, Ranjit Ambad, Prakash Kesharao Kute, Roshan Kumar Jha, and Amol Dattaroa Shinde. “The Pandemic ~~of the 21st~~ of the 21st Century - COVID-19.” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 39 (September 28, 2020): 2913–18. <https://doi.org/10.14260/jemds/2020/637>.
25. Dasari, Venkatesh, and Kiran Dasari. “Nutraceuticals to Support Immunity: COVID-19 Pandemic- A Wake-up Call.” *JOURNAL OF CLINICAL AND DIAGNOSTIC*

RESEARCH 14, no. 7 (July 2020): OE05–9.
<https://doi.org/10.7860/JCDR/2020/44898.13843>.