

## ANÆSTHETIC MANAGEMENT OF LEFT CEREBELLOPONTINE ANGLE ACOUSTIC NERVE SCHWANNOMA IN SITTING POSITION WITH INTRAOPERATIVE ELECTROMYOGRAPHIC FACIAL NERVE MONITORING

### Abstract:

~~A~~~~We managed a~~ case of left ~~cerebellopontine~~CP angle acoustic nerve schwannoma in sitting position with ~~EMG~~ monitoring for facial nerve ~~was managed~~. ~~The~~A 40 year old, male, ASA II, presented with complaints of difficulty in walking, left sided hearing loss, diminution of vision since 3 years ~~and~~ MRI ~~suggestive~~~~suggestive~~ of left CPA acoustic nerve schwannoma. Patient was operated under general ~~anesthesia~~~~anesthesia~~ in sitting position with Mayfield clamp on head without using muscle relaxant as ~~the~~ plan was to use EMG monitoring during tumor resection. Anesthesia was maintained with sevoflurane and propofol and no muscle relaxant was used, ~~still yet~~ patient was immobile. Throughout the procedure ~~ha~~emodynamic stability was maintained with ~~injection~~ noradrenaline ~~injection~~. Procedure was uneventful with complete excision of tumor and without damaging the facial nerve. Excision of this deep-seated tumor in sitting position without using muscle relaxant ~~is~~are very difficult to manage. This surgery was very challenging from ~~anesthesia~~~~anaesthetic~~ point of view as sitting position itself increases the risk of complications and avoiding movement of patient with Mayfield clamp without using muscle relaxant was a tough job.

**Comment [U1]:** State the full meaning since it is the first time it is being mentioned.

**Comment [U2]:** State full meaning

**Keywords:** Sitting position, EMG monitoring, ~~n~~No muscle relaxant

**Comment [U3]:** Add one more keyword possibly CPA

### Introduction

The sitting position was introduced in clinical practice by De Marte in 1973(1).~~However, But~~ it has been declining in popularity since ~~the~~ 1980's due to high incidence of complications. Sitting position ~~has been~~ used for resection of tumors in cerebellopontine angle, pineal region, other technically difficult tumors in infratentorial region as well as surgeries of cervical spine. ~~The a~~Advantages of this position is ~~that it~~ provides drainage of CSF and blood, provides easy access to airway and promotes favorable changes in ventilatory mechanics. This position has several life threatening complications such as venous air embolism, postural cardiovascular effects because of general anesthesia, macroglossia, pneumocephalous, quadriplegia~~and~~ peripheral nerve injury(2).

**Formatted:** Not Superscript/ Subscript

**Formatted:** Not Superscript/ Subscript

Intraoperative electromyographic (EMG) monitoring of the facial nerve during acoustic neuroma resection allows the early detection of surgical stimulation and reduces nerve dysfunction postoperatively(3). Acoustic nerve shwannoma are the most common tumors of the cerebellopontine angle (CPA)(4), accounting for more than 90% of all such tumors with the risk of injuring facial nerve during surgeries. Usage of intraoperative EMG monitoring will definitely reduce the risk of damaging facial nerve, but becomes more challenging for an anaesthetist to maintain motionless patient in sitting position with Mayfield clamp on head of the patient. ~~we managed~~A case of left CP angle tumor posted for excision in sitting position with EMG monitoring ~~was managed~~in our institute.

**Formatted:** Not Superscript/ Subscript

**Formatted:** Not Superscript/ Subscript

### Case report

A 40 year old (45kg, 155 cm) male brought to the hospital with complaints of difficulty in walking, left sided hearing loss and diminution of vision for the past 3 years. MRI was done which revealed a well marginated heterogeneously enhancing extra axial mass lesion in left CP angle cistern s/o acoustic nerve schwannoma. He was post referred for excision of the tumour. Thorough preoperative checkup was done. preop HRCT was S/O a solitary pulmonary nodule in left upper lobe s/o infective etiology. All routine workups were done including 2D ECHO which was normal.

**Comment [U4]:** State the full meaning since it is the first time it is being used.

**Comment [U5]:** Rephrase for better understanding





Fig.1 Preoperative MRI Scan

**Comment [U6]:** Source

**Comment [U7]:**



(Figure 2

Comment [U8]: Title and source

]

Patient was transferred to OT. All ASA monitors were attached, 18G Intracath was secured intravenous I.V. fluid NS started. Patient was preoxygenated with 100% oxygen at 5lit/min of oxygen and premedicated with inj. lidocaine 2% 1.5mg/kg. Induction done with Inj. thiopentone 5 mg/kg and Inj. fentanyl 2mcg/kg after that ventilation was confirmed with bag and mask and muscle relaxant Inj. vecuronium 0.1mg/kg was given. Patient was ventilated for 3 minutes and intubated with ET tube size 8 mm. After confirming the air entry is equal on both sides, ET tube was fixed. Left radial artery was cannulated for direct arterial pressure monitoring and arterial blood gas analysis intraoperatively. Triple lumen central line was cannulated in right subclavian vein. Anesthesia Anaesthesia was maintained with sevoflurane (MAC 0.8 -0.9) and oxygen and air (40%-60%). The bladder was catheterised

Comment [U9]: Full meaning

Comment [U10]: Full meaning

for urine output monitoring. After the continuous observation of CVP, arterial pressure and peak pressures, patient was slowly kept in sitting position without disturbing the hemodynamic parameters. Knees were kept at the level of the heart while the shoulders were strapped to the operating table from behind to avoid drooping. Over flexion of neck was avoided; pillows were placed below thighs and legs on both sides and pressure points were padded properly. A Mayfield frame was used, which was clamped to the side rails of the operating table to make the head and trunk immobile with respect to each other. Before fixing the frame to the scalp inj. fentanyl 1 mcg/kg was administered. After final positioning, Arterial line transducer was zeroed at the level of the middle ear. Intraoperatively analgesic was maintained with fentanyl 1mcg/kg/hr.

**Comment [U11]:** Full meaning



Fig 23: Intra operative vitals

**Comment [U12]:** State source even if it is your health facility




Fig 3: Intraoperative EtCO2 monitoring



Fig 4: intraoperative noradrenaline and propofol infusions

As the surgeon planned for facial nerve monitoring, the patient was maintained on inj. propofol infusion at 0.1-0.2 mg/kg/hr and Sevoflurane MAC was kept at 0.5 to 0.6; inj. noradrenaline infusion was started at 4 ml/hr to maintain MAP between 70mmHg to 90 mmHg. All vital parameters were maintained at a stable rate throughout the procedure. Serial arterial blood gases monitoring was done. There was no episodes of air embolism through out the surgery which is the most devastating complication in sitting position.

**Comment [U13]:** Be specific 0.1 or 0.2??

**Comment [U14]:** State the unit

The surgery was completed in 11 hours without any complications; patient was movshifted to the ICU with ET tube insitu. He was put on inj. midaz 5mg/hr with inj. fentanyl 1mcg/kg boluses every 2 hoursly for the night and extubated on the next day after stopping sedation. Patient was fully conscious and oriented, obeying commands, GCS was E4V5M6, and no facial palsy was present.

**Comment [U15]:** Full meaning and acronym in bracket

## Discussion

CPA tumors are mostly benign, slow-growing tumors with low potential for malignancy (~1%) (1). Between 5 and 10% of all intracranial tumors are located in the cerebellopontine angle (5). The most common tumors at the CPA are vestibular schwannoma, meningioma, and epidermoid tumors (6). Vestibular schwannoma accounts for 75 to 85% of all CPA tumors; Schwannomas are the primary lesion of cranial nerves involving trigeminal, facial, glossopharyngeal, vagus, and sometimes even accessory cranial nerve (7). The diagnosis of CPA tumors is made based on history, physical examination, audiometric and radiological evaluation. Magnetic Resonance Image (MRI) is the gold standard for the diagnosis of CPA tumors. Treatment options for CPA tumors include observation, radiation therapy, or microsurgery (8). In patients undergoing acoustic neuroma resection, intraoperative EMG monitoring of the facial nerve has resulted in improved preservation of facial nerve function postoperatively (1). The anaesthetic management of these patients should not only provide an immobile patient but also provide an optimal environment for facial EMG monitoring. The advantages of sitting position includes improved surgical access to the posterior fossa by facilitating gravity assisted drainage of blood and CSF (9) and decreasing ICP; It also improves surgical orientation, access to the midline structures and decreases the amount of surgical retraction needed to gain access to deeper structures; but this position however has many disadvantages such as the venous air embolism accounting for 25%-75%; cardiovascular instability, macroglossia, quadriplegia and pneumocephalus (10). Patients in the sitting position must be returned to the supine position rapidly for resuscitative measures in case of an acute cardiovascular collapse. From the surgical aspect, the sitting position gives good surgical access to the operative site (11), improves venous drainage, gives a better view of facial area for monitoring evoked responses from cranial nerve stimulation and allows for better ventilation. Conversely, the sitting position can present complications such as air emboli, postural hypotension and serious cardiac arrhythmias due to surgical stimulation of cranial nerves and brainstem. Precordial Doppler or transesophageal echocardiography monitoring improves the detection of small venous air embolism enabling its early treatment and diminishing its consequences (12).

**Formatted:** Not Superscript/ Subscript

**Formatted:** Not Superscript/ Subscript

**Formatted:** Not Superscript/ Subscript

**Formatted:** Not Superscript/ Subscript

**Formatted:** Not Superscript/ Subscript

**Formatted:** Not Superscript/ Subscript

**Comment [U16]:** Full meaning

**Formatted:** Not Superscript/ Subscript

**Formatted:** Not Superscript/ Subscript

**Formatted:** Not Superscript/ Subscript

## CONCLUSION

We have successfully managed a case of acoustic nerve schwannoma excision in sitting position with propofol infusion with facial nerve EMG monitoring.

**Comment [U17]:** Rephrase and add more contents to the conclusion

## REFERENCES

1. Porter JM, Pidgeon C, Cunningham AJ. The sitting position in neurosurgery: a critical appraisal. *British journal of anaesthesia*. 1999 Jan 1;82(1):117-28.
2. Albin. MD MS, Babinski M, Maroon JC, Jannetta PJ. Anesthetic management of posterior fossa surgery in the sitting position. *Acta Anaesthesiologica Scandinavica*. 1976 Apr;20(2):117-28.
3. Amano M, Kohno M, Nagata O, Taniguchi M, Sora S, Sato H. Intraoperative continuous monitoring of evoked facial nerve electromyograms in acoustic neuroma surgery. *Acta neurochirurgica*. 2011 May;153(5):1059-67.
4. Kankane VK, Warade AC, Misra BK. Nonvestibular schwannoma tumors in the cerebellopontine angle: A single-surgeon experience. *Asian journal of neurosurgery*. 2019 Jan;14(1):154.
- 5) Zúccaro G, Sosa F. Cerebellopontine angle lesions in children. *Child's Nervous System*. 2007 Feb 1;23(2):177-83.
- 6) Farid N. Imaging of vestibular schwannoma and other cerebellopontine angle tumors. *Operative Techniques in Otolaryngology-Head and Neck Surgery*. 2014 Mar 1;25(1):87-95.
- 7) Lak AM, Khan YS. Cerebellopontine Angle Cancer. *StatPearls [Internet]*. 2021 May 4.
- 8) Bennett M, Haynes DS. Surgical approaches and complications in the removal of vestibular schwannomas. *Otolaryngologic Clinics of North America*. 2007 Jun 1;40(3):589-609.
- 9) Jagannathan S, Krovvidi H. Anaesthetic considerations for posterior fossa surgery. *Continuing Education in Anaesthesia, Critical Care and Pain*. 2014 Oct 1;14(5):202-6.
- 10) Lalenoh DC. ANESTHESIA OF THE POSTERIOR FOSSA SURGERY. *NEUROANESTHESIA AND CRITICAL CARE (NACC) COURSE SEPTEMBER 19TH-20ST 2014*.
- 11) Howell VL, Collins MM, Rochlen LR. ANESTHESIA FOR POSTERIOR FOSSA MASS. *Neuroanesthesia: A Problem-Based Learning Approach*. 2018 Sep 15:9.
- 12) Gracia I, Fabregas N. Craniotomy in sitting position: anesthesiology management. *Current Opinion in Anesthesiology*. 2014 Oct 1;27(5):474-83.

Formatted: Font: Bold

Formatted: Numbered + Level: 1 +  
Numbering Style: 1, 2, 3, ... + Start at: 1 +  
Alignment: Left + Aligned at: 0" + Tab after:  
0.3" + Indent at: 0.3"