

Study Protocol

Assessment of Compliance Level of Healthcare Personnel at AVBRH to Appropriate Hand Hygiene Practices: A Study Protocol

Abstract-

Background: WHO recommends washing hands with water and soap or alcohol -based hand rubs for cleaning hands. Hand hygiene has been considered as central component in every programs related to infection control. Its applicable to healthcare personnel handling critical patients as well. Poor hand hygiene during intrapartum and newborn care is associated with increased death rates among newborns due to sepsis. This study aims to assess the knowledge, attitude and practices of hand washing by healthcare staff in hospital.

Methodology: will be an observational study conducted among the healthcare personnel at AVBRH hospital, Wardha. Tool consisting of self-administered observation record form on knowledge and practices of hand washing will be used for data collection from 60 healthcare workers. Data will be analyzed using appropriate statistical tests.

Expected Results- Appropriate levels of knowledge and practices about proper hand hygiene is expected within healthcare workers of AVBRH. Availability of alcohol -based hand rubs, sanitizers, soap and water is expected at all times.

Conclusion- Considering the pandemic situation, availability and use of hand sanitizers can be effective to limit the spread of infections COVID -19.

Keywords- WHO, Infection Control, Hand hygiene, hand washing, healthcare workers, compliance, knowledge, practices.

Introduction:

The WHO has recommended streamline or a piece of advice for hand hygiene and its bottom line is to wash hands by using soap and water while it seems dirty or involving blood sanguineous or other body fluids or after the lavatory. Since, the guidelines of WHO suggested washing hands with water and soap or one can be used alcohol -based hand rubs for hand washing. Hand hygiene has been considered as a foundation or central component or main ingredient in every programs related to infection control which is held by any of the healthcare personnel who are the supporters of a cause of infection in critical patients through their hands. WHO introduced "Five moments for

Comment [آذر1]: Explain the reason: COVID-19

Comment [آذر2]: It is better to use past tense for the whole of the manuscript

Comment [آذر3]: handwashing

Comment [آذر4]: program

Comment [آذر5]: Insert a valid reference

Hand Hygiene" to reduce the problems concerned with hand washing. Acc, to the recommendation of WHO, the time duration required for the proper cleaning and removal of transient flora should be extended upto 15 - 20 seconds. Due to poor hand hygiene among the HCWs, their is major cause of spreading of Nosocomial infection which leads to the increase in morbidity and mortality within the hospitalized patients. Our developing countries leads a high prevalence of infection by 19% which is a challenge for healthcare personnel. Washing hands cleanly and neatly is effectively simplest method to prevent 50% of infection.

Hand hygiene has been considered as the treating interference that will cause the cross- transmission of pathogens in a healthy healthcare environment and hence, Hand hygiene according, to WHO has been proven to decrease the occurrence of Nosocomial infections. Relevant hand hygiene practices has shown the reduction rates of all infections such as gastrointestinal, respiratory tract and skin infections. In today's Modern period, WHO has included the implementation of alcohol--based lotions or rubs. -The highest rate of infection is found in the ICU, which is the place where very critically ill patients are placed, in such a critical care areas chances of infections is majorly by touching the non-movable objects which may directly leads to object contamination. The WHO has put forward an application of evidence - based practice of hand hygiene which is referred as " five - moments pf-of hand hygiene". This five moments of hand hygiene is regarding to-the washing of hands before performing any aseptic procedure, after the condition of being laid open to undesirable body fluids, or touching a patient or its surrounding. But from the another point of view, regular hand hygiene may be challenging to the HCWs lead hand dermatitis because of water, detergents, and disinfectants.

Now when we perform this hand washing method so it includes after doing some daily activities such as after arrival at work places it could be anything, after removal of gloves or socks, etc, before taking meals, after excretion of any sort of body wastes urination and defecation, after coming in contact with the contaminated equipment(1). Most of the healthcare associated infections are transmitted via healthcare worker's. This is because most of the pathogenic organisms stays-on the body for maximum of 60 minutes(1). The WHO (World Health Organization) has recommended the five steps moments of hand hygiene which should be performed by all healthcare workers before coming in contact with the patient or performing aseptic procedures, after dealing with the high risk body fluid procedure, etc (1). Sometimes in many hospitals, many of the healthcare workers had bad practices of hand hygiene, as they do not wash their hands cleanly and sometimes not at all_(2). Washing hands daily for effective measure so as to prevent the infection. In hospital's major transmission or spread of infection is mainly among the front line survivor's that is healthcare worker's(4). To avoid or prevent the spreading of infection among HCWs they have to apply alcohol-based hand sanitizers, or other antiseptic agents (4). The whole world is fighting with novel coronavirus pandemic since 2019, so as the duty of healthcare worker's they can't avoid coming in contact with CoV 19 positive patients. Hence, hand hygiene is very much important for HCW(4).-The whole world is fighting with

Comment [آذر16]: what is the meaning of Acc?

Comment [آذر17]: Up to

Comment [آذر18]: Please describe the abbreviations in the first use

Comment [آذر19]: There is a major

Comment [آذر10]: nosocomial

Formatted: Highlight

Comment [آذر11]: lead

Comment [آذر12]: Insert a valid reference

Comment [آذر13]: Why do you use a capit...

Comment [آذر14]: have

Comment [آذر15]: , and

Comment [آذر16]: modern

Comment [آذر17]: alcohol-based

Comment [آذر18]: delete a

Comment [آذر19]: delete the

Comment [آذر20]: lead

Comment [آذر21]: insert to: to as

Comment [آذر22]: these

Comment [آذر23]: it's

Comment [آذر24]: HCWs'

Comment [آذر25]: Insert a valid reference

Comment [آذر26]: handwashing

Comment [آذر27]: workplaces

Comment [آذر28]: healthcare-associated

Comment [آذر29]: insert of

Comment [آذر30]: It should be used in the (...)

Comment [آذر31]: high-risk

Comment [آذر32]: What does it mean?

Comment [آذر33]: front line survivors that (...)

Comment [آذر34]: Where is reference no 3?

Comment [آذر35]: Covid 19 or COVID-19 (...)

Comment [آذر36]: HCWs or HCW?

~~the novel coronavirus pandemic since 2019~~. Novel coronavirus is the type of SARS-CoV-2 hence also known as COVID-19 or CoV-19. The emergence of the novel coronavirus CoV-19 has ~~an~~-major challenges for the public all around the world(5).

Comment [آذر37]: It is repetitive

For reducing the transmission of this pandemic COVID-19 is only by the simplest method i.e., hand hygiene and this is very effective among the public(5). ~~The origin of the novel corona virus which is the main cause of the current pandemic named COVID-19 is in Wuhan, China(7). This infection has started in December 2019, and now the spread has reached ~~to~~-113 countries(7). SARS-CoV-2 or COVID -19 is mostly presents in Human's this does not infect any animals. This disease shows some symptoms such as fever, cough, and dyspnea_(7). Just like Severe acute respiratory syndrome coronavirus (SARS-CoV) and Middle ~~east~~ respiratory syndrome coronavirus (MERS -CoV), these SARS -CoV -2/ COVID-19 also transmitted from person to person through respiratory droplets during sneezing and also by close contact_(7). In healthcare sector the infection caused due to insufficient hand hygiene which is directly linked with the mortality rate, economical burden also the patients suffers a lot(10). Emergency Medical Service (EMS) is ~~a highly~~ rated contaminated department where any type of patient arrives from different surrounding and are thus a high-risk way of spreading microbial transmission. In EMS, hand hygiene practices are rarely ~~been studies~~ since, as we all know that this is a proven infection control intervention_(10). Healthcare associated infections ~~has many~~ reasons lacking for the compliance of ~~hand-washing~~ practices_(11). This may includes lacking of required proper equipment, ratios of staff with patients which is very low, also some of them may have allergic infection from hand washing products, incomplete knowledge of some staff regarding the risks behind the infection and procedures, regarding bio-safety some of them shows a very casual attitudes towards infection control and doesn't even value time_(11). In some sector's, ~~the~~ main focus was given on the compliance or bad attitude of ~~the~~ healthcare worker's towards hygiene, so many of studies also done for the barriers but instead, they are lacking with ~~the~~-knowledge and attention related with the pediatric settings regarding hand hygiene compliance_(12).~~

Comment [آذر38]: Wearing a mask is more important than hand wash bec the virus is airborne

Comment [آذر39]:

Comment [آذر40]: presenting

Comment [آذر41]: East

Comment [آذر42]: These sentences must be entered in the first paragraph of the introduction

Comment [آذر43]: In the healthcare sector the infection is caused due to insufficient hand hygiene which is directly linked with the mortality rate, the economical burden also the patients suffer a lot

Comment [آذر44]: a highly

Comment [آذر45]: high-risk

Comment [آذر46]: been studied

Comment [آذر47]: Healthcare-associated

Comment [آذر48]: have

Comment [آذر49]: handwashing

Comment [آذر50]: attention-related

Comment [آذر51]: children

Comment [آذر52]: infections

Comment [آذر53]: cause

As we all know small ~~children's~~ are generally affected by ~~the infection's~~ and caused harmful diseases therefore, it is very much essential to give focus on such settings in hospitals. In ~~the~~ case of nursing education, compliance ~~of-with~~ hand hygiene is one of the most difficult tasks of outcomes_(13). It is somewhat difficult to ensure nursing students ~~to~~-recognize the use of hand hygiene when, where, what, and how in increasing rates of healthcare-associated infections_(13). Patient safety is the first most priority for any healthcare system, and ~~the~~ most effective measure regarding hand hygiene_(14). And for this, it is very much important to know that the healthcare staff has proper ~~has proper~~-knowledge about the technique and awareness about infection_(14). Otherwise, if the health staff itself doesn't have proper information then this can lead to ~~the~~- a major increase in the compliance related to the nosocomial infection(14). The only aim here was to analyze the effect of different methods, training related to hand hygiene, and education among nursing staff to ensure whether they ~~maintainig~~ proper awareness over the time_(14). Chances of having

transmission of microbial ~~vector's~~ vectors such as *Staphylococcus aureus* and other common respiratory infection's ~~is~~ occurs mainly through face-touching ~~behaviour~~ behavior (15). Most of the people had bad habits of not covering ~~the~~ their face by using clothes before going outside. They may reach in hospital after coming from surrounding's with any type of microbial vector stuck on their face and these personnel's may come directly in contact with patient's or vice versa (15). ~~The~~ hand hygiene is globally a critical infection prevention or infection control practice across all over the healthcare sector's (16). There are many approaches ~~that~~ come's forward to monitor hand hygiene compliance vary from ~~a~~ simple methods such as observational methods or using different products to taking examples of advanced methods such as the automated electronic monitoring system method. ~~The~~ Current approach supports the multimodal supplemented by educating towards enhancement of hand hygiene performance (16). ~~A~~ The practice of having hand hygiene at right time is a very much effective major for the prevention of healthcare-associated infections. This can ~~be~~ easily undertaken by any of the healthcare professionals (17). But in case of changing ~~behaviour~~ behavior of staff requires a multifaceted approach to ensure the appropriate practice of hand hygiene. This article gives main highlights on two important aspects i.e., the need ~~of~~ for hand hygiene, when it is to be needed and how it is to be done (17). To convince the user's we only need to educate them and pay their attention ~~towards~~ to the importance of hand hygiene, give the evidence in reducing bacterial flora which mostly present on the palm's of both the hands (18). The present study was done in a rural hospital ~~so as~~ to demonstrate the presence of bacterial flora. In order to reduce the outcome of floral bacteria present on the hands. It is very important to give the emphasis on teaching the healthcare workers the technique of using alcohol-based hand rub in different ways (18). Professionals were lacking with the congruence in hand hygiene training for infection prevention and control i.e., IPC. Therefore, we need to give ~~emphasize~~ emphasis on the approach to train the professionals using the concept of: "Train-the-Trainers"(TTT) especially especially for the Infection- prevention and control personnel and observe the impact on hand hygiene knowledge (19). As we know, ~~H~~ hand ~~H~~ hygiene is an essential factor for the safety of patients, for the quality care, this creates major challenges for the healthcare personnel (20). To ~~overcomes~~ with this major challenge, especially nursing staff has the proper knowledge about hand hygiene as they stay more in contact with the patients (20). It is very effective to provide the healthcare worker's with feedback related to their daily practices of hand hygiene, to monitor the compliance of hand hygiene to improve multidisciplinary program (21). Product using or consumption is less time consuming and is very much helpful in monitoring and collecting information about the frequency of hand hygiene and this information can be used as feedback for the healthcare personnel (21). To decrease the level of pathogen transmission and nosocomial infections the hand hygiene plays a very important role in every healthcare ~~facilities~~ facility. However, most of the interventions have not been sustainable (22). The given review gives an overview in the community regarding hand hygiene which includes some aspects such as hand drying, and some issues related to hand hygiene behavior (23). Hand hygiene play's a role of a barrier

between the transmission of microbial infection both in the community and healthcare sector's. Alcohol-based hand rub's is ~~a~~the simplest method to stay away from bacterial infections_(23). Most of the healthcare workers may recommends the problems with hand hygiene there are many factors ~~which~~that influence the level of hand hygiene_(24). Lacking ~~of~~products, as well as facilities, lack with the inappropriate location which may create the barriers. We aimed at making ~~the~~ hand hygiene practices very feasible while healthcare delivery. To ensure that the needed infrastructure is in the proper place so that it doesn't cause any barrier ~~for~~to the healthcare worker's_(24). Introducing alcohol hand rubs comes with the essential barrier's for the good hand hygiene practices. This is related ~~with the~~to the increasing compliance especially when it is integrated within multi-modal strategies_(24). Infection can be spread in any way such as if any person coughs or sneezes then microorganisms can easily ~~contaminated~~be transmitted via droplets for example airborne microorganisms easily infects the person via direct contact_(25). Therefore hand hygiene is essential. In particular situations such as in pandemic outbreaks, it is very much necessary to interrupt the transmission chain of any virus and ~~this can only be done by practicing proper hand hygiene~~_(25). As ~~of now~~ currently ~~we all~~, the whole world is dealing with the novel corona virus, but now it is very much in control of spreading and this is only due to proper hand sanitization and covering our face with masks. The transmission of infection is under control now. ~~Pandemic outbreak~~ Control in a pandemic can be achieved by self-isolation, maintaining social distancing, good hand hygiene, and strictly follow up for infection control tools both in hospital premises and in the public area_(25). To achieve ~~the~~ success on ~~P~~pandemic outbreak, one needs to use formulated effective hand sanitizers agents such as water, soaps, sanitizer liquids, water-based or alcohol-based hand ~~sanitize~~ sanitizer, antimicrobial soaps such that, and now at present each and every hospital are widely using ~~this~~these effective hand sanitizers_(25). Related ~~evidences~~evidence are reflected from GBD studies_(26-30).

Comment [آذر54]: Not all organisms are transmitted through hand contamination. Airborne viruses are transmitted by air and through the breathing

Methods-

Site of study - Acharya Vinoba Bhave Rural Hospital, Sawangi, Wardha.

Study Population - Healthcare professionals including nursing staff's at Acharya Vinoba Bhave Rural Hospital, Wardha.

Study Duration - 2 months study.

Study design. - Observational study.

Participants - Healthcare worker's especially main focus was given on nursing staff.

Sample size - 60 HCWs.

Research tool - Tool consisting of self - administered observation record form contains observations including knowledge of hand washing and practice of hand washing among nurses.

Source of data collection - Data ~~will be~~were collected ~~by~~ using observation recorded form.

Expected outcomes-

This observational study ~~will be~~ was done at *Acharya Vinoba Bhave Rural Hospital, Wardha* to assess the level of compliance of healthcare personnel to appropriate hand hygiene practices. To analyses the drift or direction of hand hygiene practices for 2 months. The study ~~will be~~ was done on 60 healthcare worker-s and the main focus will be given ~~on-~~ to nursing staff as they are in close contact with the patients. We ~~had done-~~ did this study so as to get good results about proper hand hygiene within HCW ~~as-~~ to reduce the rate of transmission. We ~~were~~ expected at least 50% of the participants to have a good knowledge of hand hygiene practices.

Discussion-

WHO recommended five moment alcohol-based hand rubbing formula which has much better antimicrobial property. The commonly used cleaning agents among the HCW~~s~~ was soap and water. Most of the HCWs had the training on hand washing, from multiple sources. Many of the doctors who ~~were directly comes-who directly~~ came in contact with the patients had trainings on hand-washing than other HCWs. Having knowledge of hand sanitization is important because basic hand hygiene is very much important in preventing transmission of infection within patients and health workers. Some constraints may be observed among them such as lack of water and soap and knowledge of how to wash hands. For drying hands after washing most ~~preferable-~~ preferable method is drying in free air. Hand washing can help ~~the~~ people from the bacterial infection or diseases that can cause by viruses, bacteria's~~2~~ etc. This only protect people from the transmission with ~~directly~~ physical contact. But this does not mean that all the germs all over the body ~~has~~ have been killed~~;-A~~ and the illness caused by ~~the~~ bacteria or viruses or any type of germs can be very serious. Diseases like Diarrhea or respiratory infection and many more can be preventable by simply washing our hands daily. Unwashed hands can transfer many ~~infection~~ infections from one person to another person by touching their hands to objects such as handrails, surfaces of tables, toys of small children. Most of ~~the~~ skin and eye infections can be prevented only by hand washing. Few of the related studies were reported. Khanam et. al. assessed knowledge, attitude and practice on use~~s~~ of plastic products, their disposal (31). Khandelwal et. al. assessed the knowledge of ~~H~~ hepatitis B virus infection and it's control practices among dental students (32). A ~~S~~ similar study about Dengue fever was reported by Meshram et. al. (33).- Pasari et. al. assessed knowledge about covid-19 and practices among-in hemodialysis technicians (34). Swapnil P. et. al. reported on knowledge, attitude~~2~~ and practices of allied health care professional students towards universal precaution (35-48).

Conclusion- Considering the pandemic situation, availability and use of hand sanitizers can be effective to limit the spread of infections COVID -19.

References -

Comment [آذ55]: How respiratory infection can be prevented by hand washing???

1. Jemal S. Knowledge and practices of hand washing among health professionals in Dubti Referrals Hospital, Dubti, Afar, Northeast Ethiopia. *advances in preventive medicine*. 2018 ~~Nov~~; 22;2018.
2. Gwarzo GD. Hand Hygiene practices among healthcare workers in a public hospital in North-western Nigeria. *Nigerian Journal of Basic and Clinical Sciences*. 2018 ~~Jul~~; 1;15(2):109.
3. Tyagi M, Hanson C, Schellenberg J, Chamarty S, Singh S. Hand hygiene in hospitals : an observational study in hospitals from two southern states of India. *BMC Public Health*. 2018 ~~Nov~~; 27; 18(1):1299.
4. Araghi F, Tabary M, Gheisari M, Abdollahimajd F, Dadkhahfar S. Hand Hygiene Among HealthCare workers During COVID -19 Pandemic :Challenges and Recommendations. *Dermatitis*. 2020 ~~Jul/Aug~~; 31 (4): 233-237.
5. Golin AP, Choi D, Ghahary A. Hand Sanitizers: A review of ingredients, mechanisms of action, modes of delivery, and efficacy against coronavirud. *Am J Infect Control*. 2020 ~~Sep~~; 48(9) : 1062-1067.
6. Shobowale EO, Adegunle B, Onyedibe K. An assessment of hand hygiene practices of healthcare workers of a semi-urban teaching hospital using the five moments of hand hygiene. *Niger Med J*. 2016 ~~May-Jun~~; 57(3):150-4.
7. Lotfinejad N, Peters A, Pittet D. Hand hygiene and the novel coronavirus pandemic: the role of healthcare workers. *J Hosp Infect*. 2020 ~~Aug~~; 105(4):776-777.
8. von Lengerke T, Lutze B, Krauth C, Lange K, Stahmeyer JT, Chaberny IF. Promoting Hand Hygiene Compliance. *Dtsch Arztebl Int*. 2017 Jan 20;114(3):29-36.
9. Le CD, Lehman EB, Nguyen TH, Craig TJ. Hand Hygiene Compliance Study at a large Central Hospital in Vietnam. *Int J Environ Res Public Health* 2019 ~~Feb~~; 19;16(4):607.
10. Vikke HS, Vittinghus S, Giebner M, Kolmos HJ, Smith K, Castren M, Lindstrom V. Compliance with hand hygiene in emergency medical services: an international observational study. *Emerg Med J*. 2019 Mar; 36(3):171-175.
11. Shanu S.J., Trivandrum, A study to assess the hand hygiene practices among Health care workers in CSICU. Code no :6197 ~~Nov~~ 2011.
12. Monteiro Abigail, "" Evaluating Hand Hygiene Compliance Among Healthcare workers in a Specialized Paediatric Hospital." Thesis, Geogia State University, 2018. <https://scholarworks.gsu.edu/iph-theses/592>.
13. Labrague LJ, Mc Enroe- Petite DM, van de Mortel T, Nasirudeen AMA. A systematic review on hand hygiene knowledge and compliance in student nurses. *Int Nurs Rev*. 2018 ~~Sep~~; 65(3): 336-348.
14. Martos - Cabrera MB, Mota-Romero E, Martos - Garcia R, Gomez-Urguiza JL, Suleiman - Martos N, Albendin - Garcia L, Canadas- De la Fuente GA. Hand Hygiene Teaching Strategies among Nursing Staff: A systematic Review. *Int J Environ Res Public Health*. 2019 ~~Aug~~; 22;16(17):3039.
15. Kwok YL, Gralton J, Mclaws ML. Face touching: a frequent habit that has

- implications for hand hygiene. *Am J Infect Control*.2015 ~~Feb~~;43(2):112-4.
16. Masroor N, Doll M, Stevens M, Bearman G. Approches to hand hygiene monitoring: From low to high technology approches. *Int J Infect Dis*.2017 ~~Dec~~; 65:101-104.
17. Kilpatrick C, Hosie L, Storr J. Hand hygiene--when and how should it be done* *Nurs Times*. 2013 ~~Sep 25~~ ~~Oct 1~~;109(38):16-8PMID:24313110.
18. Kapil R, Bhavsar HK, Madan M. Hand hygiene in reducing transient flora on the hands of healthcare workers: an educational intervention. *Indian J Med Microbiol*. 2015 ~~Jan~~ ~~Mar~~;33(1):125-8.
19. Tartari E, Fankhauser C, Masson - Roy S, Marguez - VillarrealH, Feenandez Moreno I, Rodriguez Navaz ML, Sarabia O, Bellissimo- Rodrigues F, Hernandez -de Mezerville M, Lee YF, Aelami MH, Mehtar S, Agostinho A, Camilleri L, Allegranzi B, Pires D, Pittet D. Train-the-Trainers in hand hygiene: a standardized approach to guide education in infection prevention and Control.2019 ~~Dec 30~~; 8(1):206.
20. Konicki T, Miller E. Use of a Simulation Intervention to examine differences in nursing students hand hygiene knowledge , beliefs, and behaviours. *Nurse Educ Today*. 2016 ~~Oct~~;45:96-101.
21. Boyce JM. Measuring healthcare worker hand hygiene activity: current practices and emerging technologies. *Infect Control Hosp Epidemiol*. 2011 ~~Oct~~; 32(10):101628.
22. Neo JR, Sagha - Zadeh R, Vielemeyer O, Franklin E. Evidence - based practices to increase hand hygiene compliance in healthcare facilities: An integrated review: *Am J Infect Control*.2016 Jun 1; 44(6)"691-704.
23. Jumaa PA. Hand hygiene: simple and complex. *Int J Infect Dis*. 2005 ~~Jan~~; 9(1):3-14.
24. Allegranzi B, Sax H, Pittet D. Hand hygiene and healthcare system change within multi-modal promotion: a narrative review. *J Hosp Infect*. 2013 ~~Feb~~; 83 suppl 1:S3 -10.
25. Jing JLJ, Pei YiT, Bose RJC, McCarthy JR, Tharmalingam N, Madheswaran T. Hand Sanitizers: A Review on Formulation Aspects, Adverse Effects, and Regulations, *Int J Environ Res Public Health*. 2020 May 11;17(9):3326.
26. Murray, Christopher J L, Aleksandr Y Aravkin, Peng Zheng, Cristiana Abbafati, Kaja M Abbas, Mohsen Abbasi-Kangevari, Foad Abd-Allah, et al. "Global Burden of 87 Risk Factors in 204 Countries and Territories, 1990–2019: A Systematic Analysis for the Global Burden of Disease Study 2019." *The Lancet* 396, no. 10258 (October 2020): 1223–49. [https://doi.org/10.1016/S0140-6736\(20\)30752-2](https://doi.org/10.1016/S0140-6736(20)30752-2).
27. Vos, Theo, Stephen S Lim, Cristiana Abbafati, Kaja M Abbas, Mohammad Abbasi, Mitra Abbasifard, Mohsen Abbasi-Kangevari, et al. "Global Burden of 369 Diseases and Injuries in 204 Countries and Territories, 1990–2019: A Systematic Analysis for the Global Burden of Disease Study 2019." *The Lancet* 396, no. 10258 (October 2020): 1204–22. [https://doi.org/10.1016/S0140-6736\(20\)30925-9](https://doi.org/10.1016/S0140-6736(20)30925-9).

28. Wang, Haidong, Kaja M Abbas, Mitra Abbasifard, Mohsen Abbasi-Kangevari, Hedayat Abbastabar, Foad Abd-Allah, Ahmed Abdelalim, et al. "Global Age-Sex-Specific Fertility, Mortality, Healthy Life Expectancy (HALE), and Population Estimates in 204 Countries and Territories, 1950–2019: A Comprehensive Demographic Analysis for the Global Burden of Disease Study 2019." *The Lancet* 396, no. 10258 (October 2020): 1160–1203. [https://doi.org/10.1016/S0140-6736\(20\)30977-6](https://doi.org/10.1016/S0140-6736(20)30977-6).
29. Kinyoki DK, Ross JM, Lazzar-Atwood A, Munro SB, Schaeffer LE, Abbasalizad-Farhangi M, et al. Mapping local patterns of childhood overweight and wasting in low- and middle-income countries between 2000 and 2017. *Nat Med* 2020;26(5):750-759.
30. Lozano R, Fullman N, Mumford JE, Knight M, Barthelemy CM, Abbafati C, et al. Measuring universal health coverage based on an index of effective coverage of health services in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020.
31. Khanam, N., V. Wagh, A.M. Gaidhane, and S.Z. Quazi. "Knowledge, Attitude and Practice on Uses of Plastic Products, Their Disposal and Environmental Pollution: A Study among School-Going Adolescents." *Journal of Datta Meghe Institute of Medical Sciences University* 14, no. 2 (2019): 57–60. https://doi.org/10.4103/jdmimsu.jdmimsu_27_19.
32. Khandelwal, V., S. Khandelwal, N. Gupta, U.A. Nayak, N. Kulshreshtha, and S. Baliga. "Knowledge of Hepatitis B Virus Infection and Its Control Practices among Dental Students in an Indian City." *International Journal of Adolescent Medicine and Health* 30, no. 5 (2018). <https://doi.org/10.1515/ijamh-2016-0103>.
33. Meshram, K., A. Meshram, S. Chopra, and U. Gajbe. "Impact of Community Based Awareness Education about Dengue by Assessing Knowledge and Preventive Practices in Rural Population." *International Journal of Research in Pharmaceutical Sciences* 11, no. 3 (2020): 3729–35. <https://doi.org/10.26452/ijrps.v11i3.2539>.
34. Pasari, A.S., A. Bhawane, M.R. Balwani, P. Tolani, V. Ramteke, and N. Deshpande. "Knowledge about Covid-19 and Practices among Hemodialysis Technicians in the Covid-19 Pandemic Era." *International Journal of Nephrology* 2020 (2020). <https://doi.org/10.1155/2020/6710503>.
35. Swapnil, P., D. Sarika, R. Alka, and P. Varsha. "Assessment of Knowledge, Attitude and Practices of Allied Health Care Professional Students towards Universal Precaution." *Indian Journal of Forensic Medicine and Toxicology* 14, no. 4 (2020): 280–85. <https://doi.org/10.37506/ijfmt.v14i4.11485>.
36. Rai A, Datarkar A, Borle RM. Are maxillomandibular fixation screws a better option than Erich arch bars in achieving maxillomandibular fixation? A randomized clinical study. *Journal of oral and maxillofacial surgery*. 2011 Dec 1;69(12):3015-8.
37. Agrawal A, Timothy J, Cincu R, Agarwal T, Waghmare LB. Bradycardia in neurosurgery. *Clinical neurology and neurosurgery*. 2008 Apr 1;110(4):321-7.

38. Bourne R, Steinmetz JD, Flaxman S, Briant PS, Taylor HR, Resnikoff S, Casson RJ, Abdoli A, Abu-Gharbieh E, Afshin A, Ahmadi H. Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the Global Burden of Disease Study. *The Lancet Global Health*. 2021 Feb 1;9(2):e130-43.
39. Borle RM, Nimonkar PV, Rajan R. Extended nasolabial flaps in the management of oral submucous fibrosis. *British Journal of Oral and Maxillofacial Surgery*. 2009 Jul 1;47(5):382-5.
40. Agrawal A, Cincu R, Goel A. Current concepts and controversies in the management of non-functioning giant pituitary macroadenomas. *Clinical neurology and neurosurgery*. 2007 Oct 1;109(8):645-50.
41. Franklin RC, Peden AE, Hamilton EB, Bisignano C, Castle CD, Dingels ZV, Hay SI, Liu Z, Mokdad AH, Roberts NL, Sylte DO. The burden of unintentional drowning: global, regional and national estimates of mortality from the Global Burden of Disease 2017 Study. *Injury prevention*. 2020 Oct 1;26(Suppl 1):i83-95.
42. Chole RH, Gondivkar SM, Gadbaile AR, Balsaraf S, Chaudhary S, Dhore SV, Ghonmode S, Balwani S, Mankar M, Tiwari M, Parikh RV. Review of drug treatment of oral submucous fibrosis. *Oral oncology*. 2012 May 1;48(5):393-8.
43. Korde SD, Basak A, Chaudhary M, Goyal M, Vagga A. Enhanced nitrosative and oxidative stress with decreased total antioxidant capacity in patients with oral precancer and oral squamous cell carcinoma. *Oncology*. 2011;80(5-6):382-9.
44. Gondivkar SM, Gadbaile AR. Gorham-Stout syndrome: a rare clinical entity and review of literature. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*. 2010 Feb 1;109(2):e41-8.
45. Gadbaile AR, Chaudhary M, Gawande M, Hande A, Sarode S, Tekade SA, Korde S, Zade P, Bhowate R, Borle R, Patil S. Oral squamous cell carcinoma in the background of oral submucous fibrosis is a distinct clinicopathological entity with better prognosis. *Journal of Oral Pathology & Medicine*. 2017 Jul;46(6):448-53.
46. Gadre PK, Ramanojam S, Patankar A, Gadre KS. Nonvascularized bone grafting for mandibular reconstruction: myth or reality?. *Journal of Craniofacial Surgery*. 2011 Sep 1;22(5):1727-35.
47. Sorte K, Sune P, Bhake A, Shivkumar VB, Gangane N, Basak A. Quantitative assessment of DNA damage directly in lens epithelial cells from senile cataract patients. *Molecular vision*. 2011;17:1.
48. Basak S, Rajurkar MN, Mallick SK. Detection of *Blastocystis hominis*: a controversial human pathogen. *Parasitology research*. 2014 Jan;113(1):261-5.