

## Original Research Article

### PRELIMINARY PHYTOCHEMICAL, ANTIOXIDANT AND HYPOLIPIDEMIC POTENTIAL OF AQUEOUS EXTRACT OF *FERULA ASAFOETIDA*-AN *IN VITRO* STUDY

Running title: Phytochemical and antioxidant activity of aqueous extract of *Ferula asafoetida*

#### ABSTRACT

**Aim:** To analyse the preliminary phytochemical, antioxidant, anti-cholesterol potential of aqueous extract of *Ferula asafoetida*.

**Background:** Hyperlipidemia is considered as one of the leading causes behind the occurrence of deadly disorders like diabetes, cardiovascular diseases, atherosclerosis etc. It is characterised by elevated levels of plasma lipids, mainly total cholesterol. Antioxidants are compounds which can inhibit oxidative damage. is the herbaceous plant belonging to the family Umbelliferae. It is used as spice in food and also used as digestive aid. It is used in the treatment of asthma and whooping cough and it also helps to reduce blood pressure.

**Methods:** Aqueous extract of *Ferula asafoetida* was prepared by hot percolation method. The screening of phytochemical constituents, assessment of in vitro antioxidant activity and anticholesterol activity were done using standard procedures and the data were analysed statistically using one-way analysis of variance (ONE-WAY ANOVA) and the significance was considered at the levels of  $p < 0.05$ .

**Results:** *Ferula asafoetida* extract was rich in phytochemicals and possessed potent in vitro antioxidant activity. Anti-cholesterol activity of *Ferula asafoetida* extract was examined and it was observed that the plant extract exhibited significant anti cholesterol potential in a dose dependent manner with an IC 50 value of 400  $\mu\text{g/ml}$ .

**Conclusion:** The study established the *in vitro* antioxidant and hypolipidemic potential of aqueous extract of *Ferula asafoetida*. It is concluded that the extract of *Ferula asafoetida* possesses potent antioxidant and anticholesterol activity.

**Keywords:** *Ferula asafoetida*, anti- cholesterol, antioxidant, phytochemical screening, Innovative technology, Novel method

## INTRODUCTION

*Ferula asafoetida* is a herbaceous plant belonging to the Umbelliferae family. It has been used as a spice in food for more than thousands of years. It is also used as a digestive aid. It can change the dull food into an appetizing meal by their colour, flavor and pungency (1). The resin gum obtained from *Ferula asafoetida* is used to treat whooping cough and ulcer. It is used in the treatment of cancer, women's allergies, blood pressure, and chemoprotive therapy (2). It is used in modern treatments in whooping cough , asthma, and also it reduces cholesterol (must cite) . The oil of *Ferula asafoetida* is reported to have antifungal activity (must cite). It has cardioprotective effects in low doses and cardiotoxic effects in higher doses (must cite). It is also used in organic farming to kill several insect pests (must cite). It helps to treat gastrointestinal, neurological and respiratory disorders (3).

Phytochemicals are the compounds produced in the plants. Secondary metabolites are present only in plants possessing various biological activities(4). They help in the normal processing and are capable of prevention or treatment of various disorders(5). Phytochemicals generally are regarded as an important compounds of research interest because of their possible health effects(6). Antioxidant is the material which terminates the chain reaction by removing the free radical(7). Antioxidants reduce the risk of cancer, cardiovascular diseases, diabetes and other diseases which are caused by oxidative stress(8).

Hyperlipidemia is a secondary metabolic disorder. It is characterised by elevated levels of serum triglycerides, cholesterol and low density lipoproteins and decreased serum levels of high density lipoproteins(9). Hyperlipidemia is regarded as a major risk factor for the premature development of cardiovascular disease like atherosclerosis, hypertension, coronary heart disease

etc. (10). It is also associated with other metabolic disorders like obesity, diabetes mellitus etc (11). Statin drugs are widely used for the treatment of hypercholesterolemia as it is an inhibitor of the key enzyme HMG CoA reductase, which is involved in the cholesterol biosynthesis (12). Our team has extensive knowledge and research experience that has translate into high quality publications(13),(14),(15),(16),(17),(18),(19),(20),(21),(22),(23),(24),(25),(26),(27),(28), (29),(30), (31),(32)Hence the study aimed to assess the anticholesterol and antioxidant activity of *Ferula asafoetida* extract by *in vitro* methods.

## **MATERIALS AND METHODS**

### **1. Preparation of Aqueous extract of *Ferula asafoetida*:**

*Ferula asafoetida* was purchased from a herbal health care centre. Crushed and made into powder. 80% aqueous extract was obtained. The extract was prepared by a hot percolation method. Phytochemical analysis, assessment of *in vitro* antioxidant and anti cholesterol potential was evaluated (33).

### **2. Phytochemical Screening test:**

#### **Test for phlobatannin:**

1ml of the extract was treated with 1ml of 1% HCl and boiled for 10 mins. The formation of red color precipitate indicates the presence of phlobatannin.

#### **Test for Carbohydrates**

Three to five drops of Molisch reagent was added with 1 mL of the extract and then 1 mL of concentrated sulphuric acid was added carefully through the side of the test tube. The mixture was then allowed to stand for two minutes and diluted with 5 mL of distilled water. The development of a red or dull violet ring at the junction of the liquids showed the presence of carbohydrates.

### **Test for Flavonoids**

Few drops of 1% liquid ammonia were taken in a test tube and along with it 1ml of the extract was added resulting in the formation of yellow color thereby indicating the presence of flavonoids.

### **Test for Alkaloids**

2ml of sample was mixed with 2ml of HCl. Then 6 drops of HCN was added and further 2 drops of picric acid was added that resulted in a creamish pale yellow ppt indicating the presence of alkaloids.

### **Test for Terpenoids**

2 ml of sample along with 2ml of chloroform and 3 ml of con.H<sub>2</sub>SO<sub>4</sub> was added. Red color ppt obtained indicates the presence of terpenoids.

### **Test for proteins**

One milliliter of ninhydrin was dissolved in 1 mL of acetone and then a small amount of extract was added with ninhydrin. The formation of purple colour revealed the presence of protein.

### **Detection of saponins**

Foam test: A fraction of the extract was vigorously shaken with water and observed for persistent foam (34).

### **Test for steroids**

One ml of chloroform was mixed with 1 mL of extract and then ten drops of acetic anhydride and five drops of concentrated sulphuric acid were added and mixed. The formation of dark red colour or dark pink colour indicates the presence of steroids.

### **3. *In vitro* antioxidant activity (DPPH free radical scavenging activity)**

Scavenging of 2, 2-Diphenyl-1-picrylhydrazyl (DPPH) radicals was assessed by the method of Hatano et al, (1989). DPPH solution (1.0 ml) was added to 1.0 ml of extract at different concentrations (0.1 to 0.5mg/ml). The mixture was kept at room temperature for 50 minutes and the activity was measured at 517 nm. Ascorbic acid at the same concentrations was used as

standard. The capability to scavenge the DPPH radical was calculated and expressed in percentage (%) using following formula:

$$\text{DPPH radical scavenging (\%)} = \frac{\text{Control OD} - \text{Sample OD}}{\text{control OD}} \times 100$$

#### 4. *In vitro anti-cholesterol activity:*

The anti-cholesterol assay was carried out as described as per the kit method (Spinreact, S.A.U-Ctra Santa Coloma, Girona, Spain). Cholesterol was dissolved in chloroform at a concentration of 2.5 mg/mL/ml. Ten microliter of the extract was pipetted into a microtiter plate followed by the addition of 2000  $\mu\text{L}$  of R1 reagent and 10  $\mu\text{L}$  of cholesterol as sample. Twenty microliter of distilled water and 2000  $\mu\text{L}$  of R1 reagent were used as blank. Negative control consisted of 20  $\mu\text{L}$  cholesterol and 2ml R1; standard consisted of 20  $\mu\text{L}$  simvastatin and 2mL R1 reagent. The contents were incubated between 0-30 min at room temperature and the absorbance was read at 500 nm in a UV-Vis spectrophotometer against reagent blank. Anti-cholesterol assay of the extract was calculated using the following equation:

$$\text{Inhibition (\%)} = \frac{\text{Negative control} - \text{Sample}}{\text{Negative OD}} \times 100$$

#### 5. Statistical analysis:

The data were subjected to statistical analysis using one-way analysis of variance (ANOVA) and Duncan's multiple range test to assess the significance of individual variations between the groups. In Duncan's test, significance was considered at the level of  $p < 0.05$ .

## RESULTS

### Phytochemical screening of *Ferula asafoetida*

The results of phytochemical analysis showed the presence of proteins, flavonoids, alkaloids, terpenoids, saponins and steroids in *Ferula asafoetida*. The carbohydrates and the amino acids are absent in *ferula asafoetida* (Table 1). The presence of phytochemicals in the extract might be the underlying reason for the therapeutic properties of *Ferula asafoetida*.

Table 1: Phytochemical screening of *Ferula asafoetida* aqueous extract.

Phytochemicals	Presences
----------------	-----------

Proteins	+
Flavonoids	+
Alkaloids	+
Terpenoids	+
Saponins	+
Steroids	+
Carbohydrates	-
Amino Acids	-

+ = It indicates the presence of the phytochemical in the *ferula asafoetida*.

- = It indicates the absence of the phytochemical in the *ferula asafoetida*.

#### **Antioxidant activity of aqueous extract of ferula asafoetida:**

Antioxidant activity of aqueous seed extracts of *Ferula asafoetida* was determined by performing DPPH free radical scavenging assay (Figure 1). The extract showed a dose-dependent increase in the antioxidant activity although its activity is less compared to the standard vitamin C. The P value is  $p < 0.05$ .




Figure 1: The figure represents the antioxidant potential of aqueous extract of *Ferula asafoetida* by DPPH assay. The X axis represents the concentration in ug/ml and the Y axis represents the % of inhibition. Green bar represents ascorbic acid, orange bar represents aqueous extract of *Ferula asafoetida* . Each bar represents the Mean±SEM of 3 independent observations. Significance at  $p < 0.05$ .

#### **AntiCholesterol activity of aqueous extract of ferula asafoetida:**

Anticholesterol activity of *Ferula asafoetida* aqueous extract revealed a dose dependent increase in the percentage of inhibitory activity. The extract showed a potent anti cholesterol activity with a  $IC_{50}$  value of 450 µg/ml (Figure 2). The activity of the extract is less compared to the standard drug simvastatin in all the tested concentrations. Simvastatin is a standard anti-cholesterol drug which acts as HMG CoA reductase inhibitor. It showed that the extract is showing less activity compared to simvastatin. The p value is  $p < 0.05$ .

## Anticholesterol potential of aqueous extract of *Ferula asafoetida*




Figure 2: The figure represents the hypolipidemic potential of aqueous extract of *Ferula asafoetida*. The X axis represents the concentration in µg/ml and the Y axis represents the % of inhibition. Blue bar represents the standard Simvastatin, orange bar represents aqueous extract of *Ferula asafoetida*. Each bar represents Mean  $\pm$  SEM of 3 independent observations. Significance at  $p < 0.05$ .

## DISCUSSION

Plants are an excellent source of biologically active compounds known as phytochemicals. It is well established that free radicals are the major cause of various chronic and degenerative diseases, like coronary heart disease, inflammation, stroke, diabetes mellitus and cancer (35). The medicinal value of plants is related to the content of phytochemicals including phenolic compounds, flavonoids, alkaloids, tannins etc. (36). The results of phytochemical analysis showed the presence of proteins, flavonoids, alkaloids, terpenoids,

saponins and steroids in *Ferula asafoetida* (Table 1). The presence of phytochemicals in the extract might be the underlying reason for the therapeutic properties of *Ferula asafoetida*.

Antioxidant activity of aqueous seed extracts of *Ferula asafoetida* was determined by performing DPPH free radical scavenging assay (Figure 1). The extract showed a dose-dependent increase in the antioxidant activity although its activity is less compared to the standard vitamin C. Free radicals are molecules possessing unpaired electrons. Antioxidants are the most effective ingredients compounds to eliminate free radicals which can create oxidative stress and are possible protective tools that protect the cells from reactive oxygen species (37). They retard the progress of many diseases as well as lipid peroxidation. Additionally, they also possess anti-inflammatory, anti-viral and anti-cancer properties ((37,38). *Ferula asafoetida* extracts which are rich in phytochemicals which have great importance in free radical scavenging. The effect of antioxidants on the DPPH free radical scavenging was considered due to their hydrogen donating ability. The extract of *Ferula asafoetida* exhibited a significant antioxidant potential with standard IC<sub>50</sub> value of 380 µg/ml. Further studies may be needed to find out the potential health benefits of the extracts in prevention and scavenging of free radicals.

Investigation of anticholesterol activity of *Ferula asafoetida* aqueous extract revealed a dose dependent increase in the percentage of inhibitory activity. The extract showed a potent anti cholesterol activity with a IC<sub>50</sub> value of 450 µg/ml (Figure 2). The activity of the extract is less compared to the standard drug simvastatin in all the tested concentrations. Simvastatin is a standard anti-cholesterol drug which acts as HMG CoA reductase inhibitor (39). Although statin groups of drugs do well in many people, the use of it is associated with potential adverse effects which include cognitive loss, neuropathy, pancreatic and hepatic dysfunction, and sexual dysfunction (40). Hence our study showed that even though the extract is showing less activity compared to simvastatin, since the extract is natural in origin it can avoid the adverse side effects created by the synthetic drugs.

## CONCLUSION

The present study can be concluded that the extract of *Ferula asafoetida* possesses potent antioxidant and anticholesterol activity. Further detailed studies on in vitro cell lines and *in vivo*

experimental animal models need to be done on the plant extract for the formulation of the anti-cholesterol drug towards the clinical utility.

## REFERENCES

1. Amalraj A, Gopi S. Biological activities and medicinal properties of Asafoetida: A review. *Afr J Tradit Complement Altern Med*. 2017 Jul;7(3):347–59.
2. Mahendra P, Bisht S. *Ferula asafoetida* : Traditional uses and pharmacological activity [Internet]. Vol. 6, *Pharmacognosy Reviews*. 2012. p. 141. Available from: <http://dx.doi.org/10.4103/0973-7847.99948>
3. Mahapatra KD, Kumar B. A REVIEW ON THERAPEUTIC USES OF OCIMUM SANCTUM LINN (TULSI) WITH ITS PHARMACOLOGICAL ACTIONS [Internet]. Vol. 3, *International Journal of Research in Ayurveda and Pharmacy*. 2012. p. 645–7. Available from: <http://dx.doi.org/10.7897/2277-4343.03512>
4. Zengin G, Uysal A, Diuzheva A, Gunes E, Jekő J, Cziáky Z, et al. Characterization of phytochemical components of *Ferula halophila* extracts using HPLC-MS/MS and their pharmacological potentials: a multi-functional insight. *J Pharm Biomed Anal*. 2018 Oct 25;160:374–82.
5. Tajbakht P, Talebi M, Rahimmalek M. Genetic diversity among and within *Ferula asafoetida* H. Karst. populations using molecular and phytochemical markers [Internet]. Vol. 155, *Phytochemistry*. 2018. p. 19–29. Available from: <http://dx.doi.org/10.1016/j.phytochem.2018.07.004>
6. Sayyar HT. Evaluation of Phytochemical Screening, Antimicrobial and Antioxidant Activities of Ethanol Extracts of *Cucumis flexuosus* and *Cucumis reticulatus* Seeds [Internet]. Vol. 41, *The Pakistan Veterinary Journal*. 2021. p. 142–6. Available from: <http://dx.doi.org/10.29261/pakvetj/2020.089>
7. Mohammad NS, Dehpour AA, Ebrahimzadeh MA, Fazel NS. Antioxidant activity of the methanol extract of *Ferula assafoetida* and its essential oil composition [Internet]. Vol. 60, *Grasas y Aceites*. 2009. p. 405–12. Available from: <http://dx.doi.org/10.3989/gya.010109>
8. Karimian V, Sepehry A, Barani H, Ebrahimi SN, Mirjalili MH. Productivity, essential oil variability and antioxidant activity of *Ferula assa-foetida* L. oleo-gum-resin during the plant exploitation period [Internet]. Vol. 32, *Journal of Essential Oil Research*. 2020. p. 545–55. Available from: <http://dx.doi.org/10.1080/10412905.2020.1794988>
9. Djamil R, Rahmat D, Zaidan S, Latifah MN. Anticholesterol Activity of Okra Fruit Extract (*Abelmoschus esculentus* (L) Moench) and Its Nanoemulsion in vivo [Internet]. Vol. 12,

Pharmacognosy Journal. 2020. p. 316–20. Available from:  
<http://dx.doi.org/10.5530/pj.2020.12.50>

10. Ramachandran AV, Ansarullah, Jadeja RN, Thounaojam MC, Patel V, Devkar RV. Antihyperlipidemic potential of a polyherbal preparation on triton WR 1339 (Tyloxapol) induced hyperlipidemia: A comparison with lovastatin [Internet]. Vol. 3, International Journal of Green Pharmacy. 2009. p. 119. Available from: <http://dx.doi.org/10.4103/0973-8258.54900>
11. Sikarwar MS, Patil MB. Antihyperlipidemic activity of *Salacia chinensis* root extracts in triton-induced and atherogenic diet-induced hyperlipidemic rats. *Indian J Pharmacol.* 2012 Jan;44(1):88–92.
12. Buhaescu I, Izzedine H. Mevalonate pathway: a review of clinical and therapeutical implications. *Clin Biochem.* 2007 Jun;40(9-10):575–84.
13. Wu F, Zhu J, Li G, Wang J, Veeraraghavan VP, Krishna Mohan S, et al. Biologically synthesized green gold nanoparticles from Siberian ginseng induce growth-inhibitory effect on melanoma cells (B16). *Artif Cells Nanomed Biotechnol.* 2019 Dec;47(1):3297–305.
14. Chen F, Tang Y, Sun Y, Veeraraghavan VP, Mohan SK, Cui C. 6-shogaol, a active constituents of ginger prevents UVB radiation mediated inflammation and oxidative stress through modulating Nrf2 signaling in human epidermal keratinocytes (HaCaT cells). *J Photochem Photobiol B.* 2019 Aug;197:111518.
15. Li Z, Veeraraghavan VP, Mohan SK, Bolla SR, Lakshmanan H, Kumaran S, et al. Apoptotic induction and anti-metastatic activity of eugenol encapsulated chitosan nanopolymer on rat glioma C6 cells via alleviating the MMP signaling pathway [Internet]. Vol. 203, *Journal of Photochemistry and Photobiology B: Biology.* 2020. p. 111773. Available from: <http://dx.doi.org/10.1016/j.jphotobiol.2019.111773>
16. Babu S, Jayaraman S. An update on  $\beta$ -sitosterol: A potential herbal nutraceutical for diabetic management. *Biomed Pharmacother.* 2020 Nov;131:110702.
17. Malaikolundhan H, Mookkan G, Krishnamoorthi G, Matheswaran N, Alsawalha M, Veeraraghavan VP, et al. Anticarcinogenic effect of gold nanoparticles synthesized from *Albizia lebbek* on HCT-116 colon cancer cell lines. *Artif Cells Nanomed Biotechnol.* 2020 Dec;48(1):1206–13.
18. Han X, Jiang X, Guo L, Wang Y, Veeraraghavan VP, Krishna Mohan S, et al. Anticarcinogenic potential of gold nanoparticles synthesized from *Trichosanthes kirilowii* in colon cancer cells through the induction of apoptotic pathway. *Artif Cells Nanomed Biotechnol.* 2019 Dec;47(1):3577–84.
19. Gothai S, Muniandy K, Gnanaraj C, Ibrahim IAA, Shahzad N, Al-Ghamdi SS, et al. Pharmacological insights into antioxidants against colorectal cancer: A detailed review of the possible mechanisms. *Biomed Pharmacother.* 2018 Nov;107:1514–22.

20. Veeraraghavan VP, Hussain S, Balakrishna JP, Dhawale L, Kullappan M, Ambrose JM, et al. A Comprehensive and Critical Review on Ethnopharmacological Importance of Desert Truffles: *Terfezia clavaryi*, *Terfezia boudieri*, and *Tirmania nivea* [Internet]. *Food Reviews International*. 2021. p. 1–20. Available from: <http://dx.doi.org/10.1080/87559129.2021.1889581>
21. Sathya S, Ragul V, Veeraraghavan VP, Singh L, Niyas Ahamed MI. An in vitro study on hexavalent chromium [Cr(VI)] remediation using iron oxide nanoparticles based beads. *Environmental Nanotechnology, Monitoring & Management*. 2020 Dec 1;14:100333.
22. Yang Z, Pu M, Dong X, Ji F, Priya Veeraraghavan V, Yang H. Piperine loaded zinc oxide nanocomposite inhibits the PI3K/AKT/mTOR signaling pathway via attenuating the development of gastric carcinoma: In vitro and in vivo studies. *Arabian Journal of Chemistry*. 2020 May 1;13(5):5501–16.
23. Rajendran P, Alzahrani AM, Rengarajan T, Veeraraghavan VP, Krishna Mohan S. Consumption of reused vegetable oil intensifies BRCA1 mutations. *Crit Rev Food Sci Nutr*. 2020 Oct 27;1–8.
24. Barma MD, Muthupandiyani I, Samuel SR, Amaechi BT. Inhibition of *Streptococcus mutans*, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish. *Arch Oral Biol*. 2021 Jun;126:105132.
25. Samuel SR. Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life? *Int J Paediatr Dent*. 2021 Mar;31(2):285–6.
26. Samuel SR, Kuduruthullah S, Khair AMB, Shayeb MA, Elkaseh A, Varma SR. Dental pain, parental SARS-CoV-2 fear and distress on quality of life of 2 to 6 year-old children during COVID-19. *Int J Paediatr Dent*. 2021 May;31(3):436–41.
27. Tang Y, Rajendran P, Veeraraghavan VP, Hussain S, Balakrishna JP, Chinnathambi A, et al. Osteogenic differentiation and mineralization potential of zinc oxide nanoparticles from *Scutellaria baicalensis* on human osteoblast-like MG-63 cells [Internet]. Vol. 119, *Materials Science and Engineering: C*. 2021. p. 111656. Available from: <http://dx.doi.org/10.1016/j.msec.2020.111656>
28. Yin Z, Yang Y, Guo T, Veeraraghavan VP, Wang X. Potential chemotherapeutic effect of betalain against human non-small cell lung cancer through PI3K/Akt/mTOR signaling pathway. *Environ Toxicol*. 2021 Jun;36(6):1011–20.
29. Veeraraghavan VP, Periadurai ND, Karunakaran T, Hussain S, Surapaneni KM, Jiao X. Green synthesis of silver nanoparticles from aqueous extract of *Scutellaria barbata* and coating on the cotton fabric for antimicrobial applications and wound healing activity in fibroblast cells (L929). *Saudi J Biol Sci*. 2021 Jul;28(7):3633–40.
30. Mickymaray S, Alfaiz FA, Paramasivam A, Veeraraghavan VP, Periadurai ND, Surapaneni KM, et al. Rhaponticin suppresses osteosarcoma through the inhibition of PI3K-Akt-mTOR pathway. *Saudi J Biol Sci*. 2021 Jul;28(7):3641–9.

31. Teja KV, Ramesh S. Is a filled lateral canal – A sign of superiority? [Internet]. Vol. 15, Journal of Dental Sciences. 2020. p. 562–3. Available from: <http://dx.doi.org/10.1016/j.jds.2020.02.009>
32. Kadanakuppe S, Hiremath S. Social and Behavioural Factors Associated with Dental Caries Experience among Adolescent School Children in Bengaluru City, India [Internet]. Vol. 14, British Journal of Medicine and Medical Research. 2016. p. 1–10. Available from: <http://dx.doi.org/10.9734/bjmmr/2016/24021>
33. Sood R. Asafoetida (*Ferula asafoetida*): A high-value crop suitable for the cold desert of Himachal Pradesh, India [Internet]. Vol. 12, Journal of Applied and Natural Science. 2020. p. 607–17. Available from: <http://dx.doi.org/10.31018/jans.v12i4.2418>
34. Iranshahy M, Iranshahi M. Traditional uses, phytochemistry and pharmacology of asafoetida (*Ferula assa-foetida* oleo-gum-resin)-a review. J Ethnopharmacol. 2011 Mar 8;134(1):1–10.
35. Mondal M, Hossain MS, Das N, Khalipha ABR, Sarkar AP, Islam MT, et al. Phytochemical screening and evaluation of pharmacological activity of leaf Methanolic extract of *Colocasia affinis* Schott [Internet]. Vol. 5, Clinical Phytoscience. 2019. Available from: <http://dx.doi.org/10.1186/s40816-019-0100-8>
36. Jamuna, Jamuna. Screening of in vitro antioxidant activity of methanolic leaf and root extracts of *Hypochoeris radicata* L. (Asteraceae) [Internet]. Journal of Applied Pharmaceutical Science. 2012. Available from: <http://dx.doi.org/10.7324/japs.2012.2722>
37. Kaur C, Kapoor HC. Anti-oxidant activity and total phenolic content of some Asian vegetables [Internet]. Vol. 37, International Journal of Food Science and Technology. 2002. p. 153–61. Available from: <http://dx.doi.org/10.1046/j.1365-2621.2002.00552.x>
38. Tutour BL, Le Tutour B. Antioxidative activities of algal extracts, synergistic effect with vitamin E [Internet]. Vol. 29, Phytochemistry. 1990. p. 3759–65. Available from: [http://dx.doi.org/10.1016/0031-9422\(90\)85327-c](http://dx.doi.org/10.1016/0031-9422(90)85327-c)
39. Rana MG. ANTIHYPERLIPIDEMIC AND ANTIOXIDANT ACTIVITY OF BIOFRACTIONS OF FLOWERS OF *HIBISCUS ROSA-SINENSIS* IN POLOXAMER 407 INDUCED HYPERLIPIDEMIC RATS [Internet]. World Journal of Pharmacy and Pharmaceutical Sciences. 2017. p. 1665–82. Available from: <http://dx.doi.org/10.20959/wjpps20177-9618>
40. Prati D, Berzuini A. Faculty Opinions recommendation of Mitochondrial damage-associated molecular patterns as potential proinflammatory mediators in post-platelet transfusion adverse effects [Internet]. Faculty Opinions – Post-Publication Peer Review of the Biomedical Literature. 2016. Available from: <http://dx.doi.org/10.3410/f.726177867.793516093>

UNDER PEER REVIEW