

**PHARMACOLOGICAL ACTIVITY REVIEW ON SELECTED INDIAN
TRADITIONAL MEDICINAL PLANTS**

ABSTRACT

The medicinal plants are the rich source of therapeutic phytoconstituents. The wide range of plants possesses vast range of medicinal uses in humanity. In the present study a review was conducted for selected plants for the medicinal activity they possess which are scientifically published in reputed journals. The activity researched is antioxidant, anti-inflammatory, antibacterial, antiulcer, hepatoprotective and anticancer activity etc. The present review shown that Pongamia pinnata, Nerium oleander methanol extract and Andrographis paniculata, Macaranga peltata alcoholic extracts possess wide range of therapeutic activities.

Comment [a1]: Please add applied conclusion at the end of abstract!

Key words: Antibacterial, Antifungal, Pongamia pinnata, Nerium oleander, Andrographis paniculata, Macaranga peltata

INTRODUCTION

Medicinal plants are used for curing innumerable diseases ([Mohammadabadi et al., 2009](#); [Heidarpour et al. 2011](#)). In terms of medicinal uses mixture of constituents found in extracts of plants are more effective than isolated compounds ([Mohammadabadi and Mozafari, 2018](#)). Many herbs in nature possess tissue regenerating property as they possess pharmacologically active compound in minute quantity along with energy boosting molecules such as carbohydrates, lipids and proteins ([Mortazavi et al., 2005](#); [Zarrabi et al., 2020](#)).

Comment [a2]: Please add reference for your used sentences. For this, you can apply added references

The plants possess various therapeutic activities, which should be brought to the notice of the scientific field for the systematic evaluation ([Heidarpour et al. 2011](#); [Mohammadabadi and Mozafari, 2019](#)). Hence, an attempt was made to review the select plants to know the possessing therapeutic activity. The plants selected are Pongamia pinnata, Nerium oleander, Andrographis paniculata and Mekaranga peltata

Pongamia pinnata

Singh RK et al. [1] reported, the anti-inflammatory activity of seed extracts of *Pongamia pinnata* in rat using carrageenan, bradykinin, PGE induced models and inflammation intensity measured by production of inflammatory molecules histamine and 5-HT. The result indicates that all extracts of seed (ethanol, petroleum ether, chloroform and acetone) shown anti-inflammatory activity when administered intra-peritoneally. (i.p)

Kumar P et al. [2] reported, the *Pongamia pinnata* flower and flower buds were having antibacterial and antifungal activity. Flower extracts shown higher antibacterial activity against *Staphylococcus aureus* and *Klebsiella pneumoniae* as compared to pod extract. Flower extract shown higher antifungal activity than pod extract.

Raut RW et al. [3] reported, extracellular synthesis of silver nanoparticles using dried leaves of *Pongamia pinnata* (L) pierre. The method represents an example of clean, nontoxic and eco-friendly method for obtaining silver nanoparticles. The capping around each particle furnish uniform chemical environment formed by the bioorganic compound present in the leaf broth, which may be mainly responsible for the stabilization of the particles. The antibacterial effect of nanosized silver colloidal solution against *Staphylococcus aureus* (ATCC 6538), *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* (ATCC9027) and *Escherichia coli* (ATCC 8739) reveals high efficacy of silver nanoparticles as a strong antibacterial agent

Sikarwar MSet al. [4]reported, antidiabetic activity of *Pongamia pinnata* leaf extracts in alloxan-induced diabetic rats. The *Pongamia pinnata* aqueous extract and ethanol extract showed potent antidiabetic effect in alloxan-induced diabetic rats and reduced the mortality rate significantly.

Rani MSet al. [5]reported, antibacterial activity of *Pongamia pinnata* on pathogens of clinical isolates. The seed extract of *Pongamia pinnata* with methanol and ethanol solvent at 100µg/ml concentration showed significant

antibacterial activity on selected (*Pseudomonas aeruginosa*, *Serratiamarcescens*, *Proteus vulgaris*, *Micrococcusluteus*, *Klebsiella pneumonia*, *Staphylococcus aureus*) in clinical isolates

Bhandirge SK et al. [6] reported,*Pongamia pinnata* bark ethanol extract having wound healing activity (excision and incision wound model). The study showed that 10% ethanolic extract ointment shown significant activity then 5% ointment, povidone iodine ointment used as standard.

Ratnam KV et al. [7] reported, that the *Pongamiapinnata* bark having anthelmintic & antiinflammatory activity. He performed anthelmintic activity in earth worms. The ethanol extract paralysed worms in 30min as compared to standard albendazole 26min and death time 35min & 30min respectively. The ethanol extracts 50mg/ml shown significant anti-inflammatory activity.

Kage DN et al. [8]reported,isolation of karajachromene from the seeds of *Pongamia pinnata*. Karajachromene at doses 25mg/kg & 50mg/kg shown 40.48% & 59.6% inhibition of paw oedema respectively as compared to standard diclofenac sodium (63.01%) at 10mg/kg body weight. He concluded that karajachromene exhibits anti-inflammatory reaction. The seed oil extracted using n-hexane for 20hr and after storage for 15days in cold condition the karajachromene crystals sedimented at the bottom test tubes. .

Dwivedi D et al. [9] reported, wound healing, anti-microbial and antioxidant potential of *Pongamia pinnata* in wistar rats. The results confirm that *Pongamia pinnata* having potent significant wound healing activity. The results confirmed by changes in wound contraction, increased tensile strength, increased hydroxyproline and hexosamine content, modulation of pro inflammatory and anti-inflammatory cytokine, moderate antimicrobial activity and in vivo antioxidant activity.

Shukla R et al. [10]reported, wound healing of ointment prepared by using *Pongamiaglabra*, *Piper nigrum*, *Momordicacharantia*. The study showed that

Momordicacharattia & *Pongamiaglabra* which were administered orally and topically with *Piper nigrum* shown significant activity and in ointment formulation can be used as market product and the effect can be enhanced by oral dose in anaemic burn wound model

Balasooriya Det al.[11] reported, wound healing potential of bark paste of *Pongamiapinnata* along with hirudotherapy: A single case study. The bark paste of *Pongamia pinnata* and paste of leaves of *Flueggealeucopyrus* was very effective and shown excellent healing effect in chronic and non-healing wound.

Nerium oleander

Kumar Set al. [12] reported, anti-inflammatory activity and antipyretic activity of *Nerium oleander* in wistar rats. The result shows that chloroform and alcoholic extract having significant anti-inflammatory activity and antipyretic activity, but petroleum ether & aqueous extract having less activity.

Singhal KG et al.[13] reported, hepatoprotective and antioxidant activity of methanolic extract of flowers of *Nerium oleander* against CCl₄-induced liver injury in rats. The result reflected the 'in vitro' effectiveness of the extract in 'in vivo' in terms of lipid peroxidation inhibitory capacity and further confirmed the significant hepatoprotective activity of the methanol extract of flowers of *Nerium oleander* L. along with its antioxidant mechanisms of action.

Benson KF et al.[14]reported, the antioxidant, anti-inflammatory, anti-apoptotic, and skin regenerative properties of an aloe vera-based extract of *Nerium oleander* leaves (NAE-8®). The result shows that NAE-8® extract has multiple beneficial effects to a cellular antioxidant protection system, and reduces cellular free radical production, both in the absence and in the presence of an inflammatory molecule

Akgun SG et al. [15]reported, wound healing activity of *Nerium oleander* in burn wound model. The study showed that burn wound can be analysed by

inflammatory molecules (Malonaldehyde, Glutathione, Myeloperoxidase, TNF-alpha, IL-beta.) activity in all stages of inflammatory reaction. Silver sulphadiazine used as standard. Author concluded that *Nerium oleander* having anti-inflammatory activity & reduced inflammatory molecules & decreased %DNA which is high in burn wound

Hase GJet al. [16] reported, phytochemical studies of *Nerium oleander* L. using GC-MS. The GCMS analysis of the solvent free extract of *Nerium Oleander* L. root having potent anticancer, anti-inflammatory, antimicrobial, 5- α -reductase inhibitor, nematocide, pesticide, analgesic, antioxidant activity due to presence of several phytoconstituents. Gas chromatography-mass spectrometry (GC-MS) analysis of *Nerium oleander* revealed the existence of the p-cresol, l-arginine, 2- methoxy phenol, mequinol, 2- methoxy phenyl ester, 2,4,6-decatrienoic acid etc. (58 compounds)

Ko YSet al. [17] reported, oleandrin and its derivative odoroside A, both cardiac glycosides exhibit anticancer effects by inhibiting invasion via suppressing the STAT-3 Signaling Pathway. The results suggest that oleandrin and odoroside A have anticancer effects by inhibiting invasion/metastasis in both MDA-MB-231 and RT-R-MDA-MB-231 cancer cells. The anticancer effects of oleandrin and odoroside A might be due to the suppression of phospho-STAT-3-mediated pathways that are involved in the regulation of invasion-related molecules, such as cancer stem cell markers and EMT-related proteins.

Kgosana KG[18]reported, the effects of extraction techniques and quantitative determination of oxalates in *Nerium oleander* as feed and other feeds. He concluded that aqueous extraction proved to be more valuable, safe, reliable and effective compared to the organic extraction methods.

Dey P et al.[19] reported, that oleander stem and root standardized extracts mitigate acute hyperglycaemia by limiting systemic oxidative stress response in

diabetic mice. The oleander stem and root extract significantly lowers hyperglycaemia and insulin resistance without affecting insulin sensitivity.

Newman RA et al. [20] reported,antiviral effects of oleandrin. The findings are *Nerium oleander* having effective antiviral activity on enveloped viruses such as HIV-I, HTLV- I, influenza and venezuelan equine encephalitis. olenadrinactive against HIV-I, HTLV- I, venezuelan equine encephalitis, chikungunya virus and SARS-CoV-2.

Andrographis paniculata

Mishra PK et al.[21] reported, the antibacterial activity of *Andrographis paniculata* leaves against clinical pathogens namely *S.aureus*, *E.faecalis*, *P.aeruginosa*, *P.vulgaris* and *E.coli*. The result showed that 75% methanol extract is having antibacterial principles in high percentage & active against methicillin resistant *S.aureus*, *E.faecalis*&*M.tuberculosis* also.

Deepak S et al.[22] reported,antioxidant & antimicrobial activity of *Andrographis paniculata* in DPPH radical scavenging assay & total reducing capacity. The leaf extracts shown considerable antioxidant & antimicrobial activity.

Pongtuluran OBet al. [23]reported, antiviral &immunostimulant activity of *Andrographis paniculata*. The study showed that *Andrographis paniculata* ethanol extract inhibited SRV(Simian Retro Virus) replication similar to positive control Lamivudine & having immunity enhancing property by stimulating lymphocyte cell proliferation at low concentration(1-16microgram) and extracted andrographolide was estimated by HPLC method.

Saini S et al.[24]reported, the traditional medicinal plants having wound healing activity along with anti-inflammatory, antioxidant, antimicrobial and analgesic activities. The study showed that petroleum ether & ethanolic extracts

of *Andrographis paniculata* having effective wound healing activity and other supporting pharmacological activities.

Zou W et al.[25] reported, anti-inflammatory effect of *Andrographis paniculata* on pelvic inflammatory disease in rats through down regulation of the NF-kB pathway. In study oral administration showed significant anti-inflammatory activity in pathogen induced PID rats with a potential mechanism of inhibiting the NF-kB pathway. In US clinics PID is treated by antibiotics as recommended by CDC (centre for disease control) only.

Polash SA et al. [26] reported, Investigation of the phytochemicals, antioxidant, and antimicrobial activity of the *Andrographis paniculata* leaf and stem extracts. *A. paniculata* leaf and stem extracts were prepared using different solvents such as water and ethanol as the polar solvents and hexane as the nonpolar solvent. The ethanolic stem extract showed the highest total phenol content and the aqueous stem extract showed the maximum free radical scavenging activity.

Gan L et al.[27] reported, isolation of diterpenoid lactones with anti-inflammatory effects from the aerial parts of *Andrographis paniculata*. In present work isolated and identified 17 diterpenoid lactones from *A. paniculata* including 2 new compounds, most of the diterpenoid lactones showed potent anti-inflammatory activity by reducing the secretion of IL-6 and TNF- α in LPS-stimulated RAW 264.7 macrophages. Author also discovered the key pharmacophores of diterpenoid lactones and developed a QSAR model to illustrate the significant relationship between the chemical structures and anti-inflammatory effects.

Mussard E et al.[28] reported, that *Andrographis paniculata* and its bioactive diterpenoids protect dermal fibroblasts against inflammation and oxidative stress. In the study, methanol extract with the bioactive compound ANDRO (comprising 0.87% of ANDRO) showed a high antioxidant effect in HDFa

cells. The methanol extract had the highest levels of bioactive compounds, followed by ethanol, water and chloroform. A large quantity of alkaloids, saponins, flavonoids, tannins, terpenoids and steroids has been found, including the highest amount of andrographolide.

Liang Yet al. [29] reported, chromosome level genome assembly of *Andrographis paniculata*. In the present study, analyzed the Cyp450 gene family in *A. paniculata*, and identified 205 putative CYP450 genes with conserved motifs. The results showed that all major classes of Cyp450 reported by the Nutzmann and Osbourn (2014) could be found in the *A. paniculata* genome. The number of Cyp450 genes with high expressions is larger in roots than in leaf. In the *A. paniculata* genome, author identified a total of 53 putative terpene synthase genes, most of which belong to the TPS-a and TPS-b subfamilies.

Macaranga peltata

Subrahmanyam VMet al.[30]reported, antibacterial and antifungal potentials of *macarangapeltata*. The results showed that leaf yields 47% methanolic extract and 30% methanolic extract obtained from stem bark. In the study also concluded that leaf extract having better antibacterial activity against gram positive microbes than gram negative microbes.

Bijesh K et al. [31] reported, isolation and characterization of antibacterial compounds from *Macaranga peltata* against clinical isolates of *Staphylococcus aureus*. The antimicrobial effect of methanol extract of *M. peltata* leaves may be due to the individual activity or synergistic activity of these identified phytochemical compounds. The following compounds identified by LCMS techniques, compounds are shikmic acid, Musennin, Rhamnetin, Lupeol acetate, Corilagin and Quercetrin.

VermaM et al.[32] reported, antibacterial and antifungal potentials of *macarangapeltata*. The leaf and stem bark samples collected, shade dried and

powdered. The methanol extracts of these samples were obtained by soxhlet extraction method. The yield obtained from leaf was 47% and from stem bark was 30%. Both the extracts proved moderate anti-bacterial activity, among them leaf extracts showed better anti-bacterial activity than the stem bark extract against both gram-positive and gram-negative bacteria.

Magadula JJ[33] reported, a review of phytochemical and pharmacological activity of the genus *Macaranga*. The 300 known species of *Macaranga* plants, less than 30 plant species have been investigated phytochemically, this calls for further work to be done on other known species of this genus. The major classes of compounds reported in the literature include flavonoids, stilbenes, terpenes, tannins, coumarins and others. The pharmacological review of the genus indicated many flavonoids and stilbenes to be isolated from the leaves and exhibited strong, moderate to weak anticancer properties, also showed significant antioxidant, anti-inflammatory and antimicrobial activity

Badarudheen R et al.[34] reported, antibacterial activity of *Macaranga peltata*. The comparison of zone of inhibition in study indicates that methanolic extract of stem and leaves has better antibacterial activity than the acetone and petroleum ether fruit extract.

Nehete M et al. [35] reported, the antioxidant, antimicrobial and wound healing potential of *macarangapeltata* bark extracts. The study confirmed that *Macaranga peltata* bark methanol extract having antimicrobial, antioxidant and wound healing activity

Palakka S et al.[36] reported, antioxidant property of extracts of *Mecaranga peltata* by DPPH free radical scavenging activity. In the result the extracts demonstrated promising antioxidant activity, antioxidant activity correlated with phenolic content. The ethanol extract proved to be having highest antioxidant activity.

Palakka S et al. [37] reported,*In vitro* anticancer screening of ethanolic extracts of *Macaranga peltata* leaves. The ethanolic extract of *Macrangapeltata* leaves were subjected to MTT assay using HeLa and SK-Mel-28 cell lines. The extract gave promising result for the anticancer activity on both the cell lines with better effectiveness against the cervical cell cancer.

Thrinitha B et al. [38]reported,reportedhepatoprotective activity of various extracts of *Macarangapeltata* (Roxb.) on paracetamol-induced hepatotoxicity rats. The administration of methanolic concentrates of *Macaranga peltata* attenuated the hepatic marker enzymes and silymarin restored enzyme activities to normal values.

NOTE:

The study highlights the efficacy of "Traditional Medicine" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

References:

- [Heidarpour F, Mohammadabadi MR, ISM Zaidul, B Maherani, N Saari, AA Hamid, F Abas, MYA Manap, MR Mozafari 2011. Use of prebiotics in oral delivery of bioactive compounds: a nanotechnology perspective. Pharmazie 66 \(5\), 319-324](#)
- [Mortazavi SM, Mohammadabadi MR, MR Mozafari 2005. Applications and in vivo behaviour of lipid vesicles. NanoliposomesFrom Fundamentals to Recent Developments, 67-76.](#)
- [Mohammadabadi MR, El-Tamimy M, R Gianello, MR Mozafari 2009. Supramolecular assemblies of zwitterionicnanoliposome-polynucleotide complexes as gene transfer vectors: Nanolipoplex formulation and in vitro characterisation Journal of liposome research 19 \(2\), 105-115.](#)
- [ZarrabiA, AlipoorAmroAbadiM, KhorasaniS, MohammadabadiM, A Jamshidi, S Torkaman, E Taghavi, MR Mozafari, B Rasti 2020. Nanoliposomes and Tocosomes as Multifunctional Nanocarriers for the Encapsulation of Nutraceutical and Dietary Molecules. Molecules 25 \(3\), 638.](#)

Mohammadabadi MR, Mozafari MR 2018. Enhanced efficacy and bioavailability of thymoquinone using nanoliposomal dosage form. Journal of Drug Delivery Science and Technology 47 (1), 445-453.

1. Mohammadabadi MR, Mozafari MR 2019. Development of nanoliposome-encapsulated thymoquinone: evaluation of loading efficiency and particle characterization. Journal of Biopharmaceuticals 11 (4), 39-46

1-2. Singh RK, Pandey BL. Anti-inflammatory activity of seed extracts of *Pongamia pinnata* in rat. Indian J. of Physio. Pharmacol. 1996 Oct 1;40:355-8.

2-3. Kumar P, Semalty A, Semalty M. Antibacterial and Antifungal activity of *Pongamia pinnata*. Indian drugs.2010;47(3):48- 50

3-4. Raut RW, Kolekar NS, Lakkakula JR, Mendhulkar VD, Kashid SB. Extracellular synthesis of silver nanoparticles using dried leaves of *Pongamiapinnata* (L) pierre. Nano-Micro Lett. 2010 Jun 1; 2(2):106-13.

4-5. Sikarwar MS, Patil MB. Antidiabetic activity of *Pongamia pinnata* leaf extracts in alloxan-induced diabetic rats. Int. J. Ayur. Res. 2010 Oct; 1(4):199.

5-6. Rani MS, Dayanand CD, Shetty J, Vegi PK, Kutty AM. Evaluation of antibacterial activity of *Pongamiapinnatalinn* on pathogens of clinical isolates. Am. J. Phytomed. Clin. Thera.. 2013; 1(8):645-51.

6-7. Bhandirge SK, Tripathi AS, Bhandirge RK, Chinchmalatpure TP, Desai HG, Chandewar AV. Evaluation of wound healing activity of ethanolic extract of *Pongamia pinnata* bark. Drug Res. 2015 Jan 21; 65(6):296-9.

7-8. Ratnam KV, Aishwarya KV. In vitro evaluation of anthelmintic and anti-inflammatory activities of *Pongamia pinnata* bark. J Glob. Trends Pharm Sci 2016; 7(2): 3228 - 3232

8-9. Kage DN, Tabassum N, Malashetty VB, Deshpande R, Seetharam YN. Isolation and pharmacological studies of karanjachromene from the seeds of *Pongamia pinnata* (L. Pierre). Int. J Cur Res Rev.2016; 8(17):35.

Formatted: English (United States),
Pattern: Clear

- 9.10. Dwivedi D, Dwivedi M, Malviya S, Singh V. Evaluation of wound healing, anti-microbial and antioxidant potential of *Pongamia pinnata* in wistar rats. J. Tr. Compl. Medi. 2017; 7(1):79-85.
- 10.11. Shukla R, Kashaw V. Wound healing prospective of *Pongamiaglabra*, *Piper nigrum* and *Momordicacharantia* on albino rats using anemic burn wound model. J. Dr. Deli. Thera. 2018; 8(6):146-54.
- 11.12. Balasooriya D, Karunarathna C, Uluwaduge I. Wound healing potential of bark paste of *Pongamiapinnata* along with hirudotherapy: A single case study. J. Ayur. Inte. Med. 2021 Apr 2.
- 12.13. Kumar S, Anand GR. Evaluation of anti-inflammatory activity of *Nerium oleander*. Pharmacia. 2010;1:33-6.
- 13.14. Singhal KG, Gupta GD. Hepatoprotective and antioxidant activity of methanolic extract of flowers of *Nerium oleander* against CCl₄-induced liver injury in rats. Asian Pacific J. Trop. Med.. 2012 Sep 1;5(9):677-85.
- 14.15. Benson KF, Newman RA, Jensen GS. Antioxidant, anti-inflammatory, anti-apoptotic, and skin regenerative properties of an *Aloe vera*-based extract of *Nerium oleander* leaves (NAE-8®). Clini. Cosmetic. Inves. Dermatol. 2015;8:239.
- 15.16. Akgun SG, Aydemir S, Ozkan N, Yuksel M, Sardas S. Evaluation of the wound healing potential of *Aloe vera*-based extract of *Nerium oleander*. Northern clini. Istanbul. 2017;4(3):205.
- 16.17. Hase GJ, Deshmukh KK, Pokharkar RD, Gaje TR, Phatanagre ND. Phytochemical Studies on *Nerium oleander* L. using GC-MS. Int. J. Pharm.Phyto. Res. 2017;9(6):885-91.
- 17.18. Ko YS, Rugira T, Jin H, Park SW, Kim HJ. Oleandrin and its derivative odoroside A, both cardiac glycosides, exhibit anticancer effects by inhibiting invasion via suppressing the STAT-3 signaling pathway. Int. J. Mol. Sci.. 2018;19(11):3350.

- ~~18~~.~~19~~.Kgosana KG. The effects of extraction techniques and quantitative determination of oxalates in *Nerium oleander* and feeds. J. ~~of~~-Veterinary Res.- 2019;86(1):1-9.
- ~~19~~.~~20~~.Dey P, Saha MR, Roy Choudhuri S, Sarkar I, Halder B, Poddar-Sarkar M, Sen A, Chaudhuri TK. *Oleander* stem and root standardized extracts mitigate acute hyperglycaemia by limiting systemic oxidative stress response in diabetic mice. Adv. ~~in~~-Pharmacol. Sci. 2019; 8
- ~~20~~.~~21~~.Newman RA, Sastry KJ, Arav-Boger R, Cai H, Matos R, Harrod R. Antiviral effects of oleandrin. J. Exp. Pharmacol. 2020;12:503.
- ~~21~~.~~22~~.Mishra PK, Singh RK, Gupta A, Chaturvedi A, Pandey R, Tiwari SP, Mohapatra TM. Antibacterial activity of *Andrographispaniculata* (Burm. f.) Wall ex Nees leaves against clinical pathogens. J. Pharm Res. 2013; 7(5): 459-62.
- ~~22~~.~~23~~.Deepak S, Pawar A, Shinde P. Study of antioxidant and antimicrobial activities of *Andrographis paniculata*. Asian J. Plant Sci. Res. 2014; 4(2): 31-41.
- ~~23~~.~~24~~.Pongtuluran OB, Rofaani E. Antiviral and immunostimulant activities of *Andrographis paniculata*. Hayati J. Biosci. 2015; 22(2): 67-72.
- ~~24~~.~~25~~.Saini S, Dhiman A, Nanda S. Traditional Indian medicinal plants with potential wound healing activity: a review. Int. J. Pharm. Sci. Res. 2016 May 1;7(5):1809.
- ~~25~~.~~26~~.Zou W, Xiao Z, Wen X, Luo J, Chen S, Cheng Z, Xiang D, Hu J, He J. The anti-inflammatory effect of *Andrographispaniculata* (Burm. f.) Nees on pelvic inflammatory disease in rats through down-regulation of the NF- κ B pathway. BMC Comp. Alter. Medi. 2016; 16(1): 1-7.
- ~~26~~.~~27~~.Polash SA, Saha T, Hossain MS, Sarker SR. Investigation of the phytochemicals, antioxidant, and antimicrobial activity of the *Andrographis paniculata* leaf and stem extracts. Adv. Biosci. Biotech. 2017; 8(05): 149.

- ~~27~~28.Gan L, Zheng Y, Deng L, Sun P, Ye J, Wei X, Liu F, Yu L, Ye W, Fan C, Liu J. Diterpenoid lactones with anti-inflammatory effects from the aerial parts of *Andrographis paniculata*. Mol. 2019;24(15):2726.
- ~~28~~29.Mussard E, Jousselin S, Cesaro A, Legrain B, Lespessailles E, Esteve E, Berteina-Raboin S, Toumi H. *Andrographis paniculata* and its bioactive diterpenoids protect dermal fibroblasts against inflammation and oxidative stress. Antioxidants. 2020; 9(5): 432.
- ~~29~~30.Liang Y, Chen S, Wei K, Yang Z, Duan S, Du Y, Qu P, Miao J, Chen W, Dong Y. Chromosome level genome assembly of *Andrographis paniculata*. Frontier. Genetic. 2020; 30(11): 701.
- ~~30~~31.Subrahmanyam VM, Rao V, Dhanaraj SA, Vasanth Raj P. Investigation of antibacterial and antifungal potentials of *Macaranga peltata*. Int J Cur Res Rev. 2013 Apr; 5(07).
- ~~31~~32.Bijesh K, Sebastian D. Isolation and characterization of antibacterial compounds from *Macaranga peltata* against clinical isolates of *Staphylococcus aureus*. Int. J. Biol. Pharm. Res. 2013;4:1196-203.
- ~~32~~33.Verma M. NK, Thakar M, Subrahmanyam VM, Rao V, Dhanaraj SA. Investigation of antibacterial and antifungal potentials of *macarangapeltata*. Int. J. Curr. Res. Rev. 2013; 5(7): 17-24
- ~~33~~34.Magadula JJ. Phytochemistry and pharmacology of the genus *Macaranga*: a review. J. Med. Plant. Res. 2014;8(12):489-503.
- ~~34~~35.Badarudheen R, Lubna N, Dufaida KM, Mashhoor K. Antibacterial Activity of *Macaranga peltata*.Int. J. Adv. Res. Emerging Discipli.2015; 3(2): 77-83
- ~~35~~36.Nehete M,Tatke P. Antioxidant, antimicrobial and wound healing potential of *macarangapeltata* bark extracts. Eur. J. Pharm. Med. Res. 2016; 3(10): 313-320
- ~~36~~37.Palakkal S, Ganesan V. Evaluation of Antioxidant Property of Extracts of *Macaranga peltata* by DPPH Free Radical Scavenging Activity. Asian J. Pharm. Health Sci. 2018;8(4).

37,38.Palakka S, Ganesan V. In vitro anticancer screening of ethanolic extracts of *Macaranga peltata* leaves. Pharm. innovation Int. J. 2019; 8(4): 787-8

38,39.Thrinitha B, Murali R, Manichandrika P. Studies on Hepatoprotective Activity of Various Extracts of *Macarangapeltata* (Roxb.) on Paracetamol-Induced Hepatotoxicity Rats. J. Evol. Medi. Dent. Sci. 2020;9(22):1692-7.

UNDER PEER REVIEW