

Knowledge and Awareness about the importance of stem cells in dentistry - A Survey

ABSTRACT

BACKGROUND:

Stem cells, the hero of regenerative medicine, are self renewable and has the potential to regenerate. Dental stem cells have proven to be an excellent source for mesenchymal cells. Dental mesenchymal cells can be applied for the replacement of dental and oral tissues against various tissue-damaging conditions including dental caries, periodontitis, and oral cancers, as well as for systemic regulation of excessive inflammation in immune disorders, such as autoimmune diseases and hypersensitivity. In this study, a survey was conducted among the dental students of a private dental hospital regarding their awareness about stem cell therapy used in dentistry.

MATERIALS AND METHODS: The study was conducted with the help of a questionnaire which was circulated with the help of social media and the internet. The target population were the dental practitioners and students of Saveetha dental college. The questionnaire consisted of 15 questions that tested their knowledge and awareness regarding stem cell therapy in dentistry.

RESULTS: A majority of the undergraduate dental students were aware of basics in stem cell therapy in dentistry. The majority were willing to practice regenerative therapy in future or invest for the same in one way or the other. The overall response of and the percent analysis were calculated for each question.

CONCLUSION: It can be concluded that the dental students of Saveetha dental college have sufficient knowledge and awareness regarding dental stem cells. Practice of dental stem cell therapy may be a big step now but is definitely going to be an advantage in the future.

KEYWORDS

Mesenchymal stem cells, regenerative dentistry, tissue engineering, innovative technique, novel method

RUNNING TITLE: Survey on the knowledge of importance of stem cells in dentistry

INTRODUCTION

Stem cells play an important role in dentistry as they are undifferentiated cells with the capability to produce different types of cells and treat diseases.(1) They are self renewable and can regenerate missing tissues. Teeth are said to be the most natural and noninvasive source of stem cells.(2) Various research on dental stem cells have opened the gate for new treatment protocols in the field of endodontics, periodontology and various oral and maxillofacial procedures(3). There are two types of stem cells, namely, adult stem cells and embryonic stem cells. Dental stem cells are a type of adult stem cells. Owing to the regenerative properties of dental stem cells and their plasticity, they can be a good source of mesenchymal cells.(3,4) These mesenchymal cells play a significant role in organ development and postnatal repair. Though mesenchymal cells are regenerative they show immune-regulatory capacity and tropic activity. These features are important for the establishment of a regenerative micro-environment at the sites of tissue injury. (5) Mesenchymal cells secrete various bio active factors that inhibit apoptosis and scarring.

Dental pulp stem cells are stem cells which are present in the pulp. They were first derived from permanent third molars of adult humans. Recently they have gained attention due to the number of research and their accessibility and the associated relatively low cost of integration into regenerative therapy. In India, the application and awareness of dental stem cells are still in the budding stage and hence more research needs to be done.(6)

In dentistry, these dental pulp stem cells have already been introduced for alveolar bone augmentation which is basically increasing the alveolar bone volume.(7)

Adult stem cells have great potential for the regeneration of craniofacial structures and this can hence contribute to tissue engineering.(8) Craniofacial structures like mandibular condyle, calvarial bone, cranial suture, and subcutaneous adipose tissue have all been engineered from mesenchymal cells.(9) Long term preservation of these dental stem cells are being looked into. Preservation of dental stem cells can be used for future personalised medication and treatment. This mirrors the idea of banking umbilical cord blood.(10)

It is often said that the oral cavity mirrors the rest of the body. Owing to the strong association between the oral health and overall health of an individual, advancements are being looked into in order to maintain teeth and oro facial bone to combat general decline in oral health. Though the use of dental implant and prosthodontics procedures are common and effective, they are expensive and can result in rejection or failure.(10,11) Recent research is shifting focus to regenerative medicine in dentistry. Use of the regenerative medicine in dentistry requires a proper knowledge about the different types of stem cells and their applications. In this study, a survey was conducted among the dental students of a private dental hospital regarding their awareness about stem cell therapy used in dentistry. Our team has extensive knowledge and research experience that has translate into high quality publications (12), (13),(14),(15),(16),(17),(18),(19),(20),(21),(22),(23),(24),(25),(26),(27),(28),(29),(30),(31),(32)

MATERIALS AND METHODOLOGY

This cross-sectional study was conducted among the undergraduate and postgraduate students of Saveetha Dental College and Hospital, Chennai. To evaluate the knowledge of the participants about stem cells and their importance in dentistry , a self-administered questionnaire containing 9 questions was prepared.The questions revolved around the awareness of the dental students regarding the types of stem cells, stem cell banks,sources of dental stem cells, and their willingness to inculcate the practice of dental stem cell therapy. The questionnaire was validated and later distributed to the participants. A web-based questionnaire was also developed using Google forms and was circulated. The participation of the subjects was kept voluntary and

nobody was not obligated to fill the form. Questions were answered with “yes” or “no” or by marking the correct responses. Frequency analysis and percentage analysis were done with the obtained results. Data was recorded in Microsoft Excel and exported to the statistical package of social science for windows (SPSS) and subjected to statistical analysis. Descriptive statistics were done and Chi square tests are used for comparison.

RESULTS

In this study around 150 responses were collected. The overall response and the percent analysis were calculated for each question. 4% of first year students, 8% of second year students, 18% of third year and 22% of final year students were aware about the different types of stem cells. Pearson chi square test shows p value is 0.639 (p value > 0.05) (Fig 1). Hence it is statistically not significant. 8% of first years and 4% of second years, 19% of third and final years were aware about adult stem cells and embryonic stem cells Pearson chi square test shows p value is 0.074 (p value > 0.05). Hence it is statistically not significant. (Fig 2). 3% of first years, 6% of second years, 21% of third years and 14% of final years were aware that dental stem cell banks are present in India. Pearson chi square test shows p value is 0.223 (p value > 0.05). Hence it is statistically not significant. (Fig 3). 3% of first years, 9% of second years, 16% of third years and 22% of final years were aware that dental stem cells can be retrieved from apical papilla of the tooth (Fig 4). 1% of first years, 8% of second years, 12% of third years and 17% of final years were aware that dental stem cells can be used to develop non dental tissues and organs (Fig 5). 6% of first years, 10% of second years, 21% of third years and 24% of final years are willing to practice stem cell based regenerative therapy in their practise (Fig 6). 4% of the first years, 6% of second years, 20% of third years and 23% of final years are willing to invest for dental stem cell banking (Fig 7).




Figure 1: Graph represents the association between the awareness of different types of stem cells and the year of study. The X axis represents the awareness of different types of stem cells and the Y axis represents the number of participants belonging to different years of study. Blue denotes first years, green denotes second years, yellow denotes third years and purple denotes final year participants. 4% of first year students, 8% of second year students, 18% of third year and 22% of final year students were aware about the different types of stem cells. Pearson chi square test shows p value is 0.639 (p value > 0.05). Hence it is statistically not significant.




Figure 2: Graph represents the association between the knowledge on the different types of stem cells and the year of study. The X axis represents the different types of stem cells and the Y axis represents the number of participants belonging to different years of study. Blue denotes first years, green denotes second years, yellow denotes third years and purple denotes final year participants 8% of first years, 4% of second years, 19% of third and final years were aware about adult stem cells and embryonic stem cells. .Pearson chi square test shows p value is 0.074 (p value > 0.05). Hence it is statistically not significant.




Figure 3: Graph represents the association between the awareness of participants regarding the dental stem cell banks in India and the year of study. The X axis represents the awareness of dental stem cell banks in India and the Y axis represents the number of participants belonging to different years of study. Blue denotes first years, green denotes second years, yellow denotes third years and purple denotes final year participants. 3% of first years, 6% of second years, 21% of third years and 14% of final years were aware that dental stem cell banks are present in India. Pearson chi square test shows p value is 0.223 (p value > 0.05). Hence it is statistically not significant.




Figure 4: Graph represents the association between the awareness of students regarding dental stem cells being sourced from apical papilla of the tooth and the years of study. The X axis represents the awareness of students regarding dental stem cells being sourced from the apical papilla of the tooth and the Y axis represents the number of participants belonging to different years of study. Blue denotes first years, green denotes second years, yellow denotes third years and purple denotes final year participants. 3% of first years, 9% of second years, 16% of third years and 22% of final years were aware that dental stem cells can be retrieved from apical papilla of the tooth. Pearson chi square test shows p value is 0.369 (p value > 0.05). Hence it is statistically not significant.




Figure 5: Graph represents the association between the awareness of students regarding dental stem cells being used to develop non dental tissues and organs and the year of study. The X axis represents the awareness of students regarding dental stem cells being used to develop non dental tissues and organs and the Y axis represents the number of participants belonging to different years of study. Blue denotes first years, green denotes second years, yellow denotes third years and purple denotes final year participants. 1% of first years, 8% of second years, 12% of third years and 17% of final years were aware that dental stem cells can be used to develop non dental tissues and organs. Pearson chi square test shows p value is 0.275 (p value > 0.05). Hence it is statistically not significant.




Figure 6: Graph represents the association between the willingness of students to practice dental stem cell therapy and the different years of study. The X axis represents the willingness of students regarding the practice of dental stem cell therapy and the Y axis represents the number of students belonging to different years of study. Blue denotes first years, green denotes second years, yellow denotes third years and purple denotes final year participants. 6% of first years, 10% of second years, 21% of third years and 24% of final years are willing to practice stem cell based regenerative therapy in their practise. Pearson chi square test shows p value is 0.589 (p value > 0.05). Hence it is statistically not significant.




Figure 7: Graph represents the association between the willingness of students to invest for dental stem cell banking and the year of study. The X axis represents the willingness of students regarding the investment of dental stem cell therapy and the Y axis represents the number of students belonging to different years of study. Blue denotes first years, green denotes second years, yellow denotes third years and purple denotes final year participants. 4% of the first years, 6% of second years, 20% of third years and 23% of final years are willing to invest for dental stem cell banking. Pearson chi square test shows p value is 0.188 (p value > 0.05). Hence it is statistically not significant.

DISCUSSION

Stem cells in dentistry can help a great deal by helping the growth of new dental cells in a laboratory, it can help in the research causes of genetic defects in cells, research how diseases occur or why certain cells develop into cancer cells. It can also test new drugs for safety and effectiveness. The present survey was conducted among undergraduate dental students wherein around 150 students took the survey. The collected data was subjected to descriptive statistical analysis and chi square tests in SPSS.

This study reveals that the dental students showed the willingness to know about stem cell potential in dentistry which may be indicative of the acceptance of newer treatment modalities. Participants who believed that stem cell banking in dentistry were also willing to practice stem cell therapy in future. 4% of the first years, 6% of second years, 20% of third years and 23% of final years are willing to invest for dental stem cell banking. Although the majority of them believed that stem cell tissue regeneration will be advantageous in stem cell therapy, most were not sure whether these stem cells could be used to develop non-dental tissues or organs.. Only 1% of first years, 8% of second years , 12 % of third years and 17 % of final years were aware that dental stem cells can be used to develop non dental tissues and organs. There are numerous applications of this progenitor cell such as continued root formation, **regeneration of an immature tooth with extensive pulp damage**, periodontal regeneration, biological tooth, and stem cell-based therapies,

Most of the participants of this survey seemed up to date with the current evolvments in the field of dentistry as they were able to answer most of the questions correctly. In a similar study conducted earlier by Epelman et.al it was found out that 56% of the participants had received continued education regarding stem cells in dentistry and hence had sufficient knowledge regarding the same (33). **However, those students were future endodontists when compared to the present survey were dental undergraduates were tested.** There is a need to spread awareness regarding these guidelines as the opinion of the dental residents may help to reframe them which in turn might facilitate the safety of procedures/treatment done using dental stem cells. Another

study conducted by Utneja et.al showed that regenerative dentistry is a **far better option** than implant dentistry and that maybe in about 20 years teeth can also be replaced using regenerative procedures (34). **Currently only 23% of the dentists in the world use some type of regenerative procedures in their practice.** Future tissues like tissue engineered bone grafts, engineered joints and cranial sutures can be developed with stem cell therapy (35). A team of professionals including stem cell biologists, molecular biologists, geneticists, polymer and materials scientists, mechanical engineers and clinicians with knowledge of oral and maxillofacial disorders is needed to develop the field of craniofacial tissue engineering (36).

CONCLUSION

From the above mentioned results, it can be concluded that the dental students have sufficient knowledge and awareness regarding the different types of dental stem cells. Practice of dental stem cell therapy may be a big step now but is definitely going to be an advantage in the future.

REFERENCES

1. Manipal S, John J, Saravanan S, Meignana Arumugham I. Stem cells in dentistry – A review [Internet]. Vol. 5, Indian Journal of Dentistry. 2014. p. 43–7. Available from: <http://dx.doi.org/10.1016/j.ijd.2013.11.011>
2. Zheng LW, Cheung LK. Stem Cells in Dentistry [Internet]. Stem Cell and Regenerative Medicine. 2012. p. 140–8. Available from: <http://dx.doi.org/10.2174/978160805008611001010140>
3. Shruthi M. IMPORTANCE OF STEM CELLS IN DENTISTRY [Internet]. Vol. 4, annals and essences of dentistry. 2012. p. 75–8. Available from: <http://dx.doi.org/10.5368/aedj.2012.4.2.4.4>
4. Paz AG, Maghaireh H, Mangano FG. Stem Cells in Dentistry: Types of Intra- and Extraoral Tissue-Derived Stem Cells and Clinical Applications [Internet]. Vol. 2018, Stem Cells International. 2018. p. 1–14. Available from: <http://dx.doi.org/10.1155/2018/4313610>
5. Mohamad SA, Milward MR, Hadis MA, Kuehne SA, Cooper PR. Photobiomodulation of mineralisation in mesenchymal stem cells. Photochem Photobiol Sci [Internet]. 2021 May 4; Available from: <http://dx.doi.org/10.1007/s43630-021-00047-5>
6. Chang C-C, Lin T-A, Wu S-Y, Lin C-P, Chang H-H. Regeneration of Tooth with Allogeneous, Autoclaved Treated Dentin Matrix with Dental Pulpal Stem Cells: An In Vivo Study. J Endod. 2020 Sep;46(9):1256–64.
7. Baudry A, Uzunoglu E, Schneider B, Kellermann O, Goldberg M. From pulpal stem cells to

- tooth repair: an emerging field for dental tissue engineering [Internet]. Vol. 1, Evidence-Based Endodontics. 2016. Available from: <http://dx.doi.org/10.1186/s41121-016-0002-x>
8. Analysis of Prevalence of Oral Squamous Cell Carcinoma in Patients with History of Chronic Irritation of Oral Tissues - A Retrospective Study [Internet]. Indian Journal of Forensic Medicine & Toxicology. 2020. Available from: <http://dx.doi.org/10.37506/ijfmt.v14i4.12511>
 9. Risbud MV, Shapiro IM. Stem cells in craniofacial and dental tissue engineering [Internet]. Vol. 8, Orthodontics & Craniofacial Research. 2005. p. 54–9. Available from: <http://dx.doi.org/10.1111/j.1601-6343.2005.00324.x>
 10. Das M, Das A, Barui A, Paul RR. Comparative evaluation of proliferative potential and replicative senescence associated changes in mesenchymal stem cells derived from dental pulp and umbilical cord. Cell Tissue Bank [Internet]. 2021 Apr 26; Available from: <http://dx.doi.org/10.1007/s10561-021-09926-8>
 11. Khodabandeh Z, Haghghat S, Tanideh N, Zare S, Farrokhi F, Karandish M, et al. Comparing the effects of *Elaeagnus Angustifolia*, *Hypericum Perforatum* and *Psidium Guajava* extracts on metabolic activity of dental pulp-derived mesenchymal stem cells. Cell Tissue Bank [Internet]. 2021 Apr 11; Available from: <http://dx.doi.org/10.1007/s10561-021-09923-x>
 12. E A, Aswani E, Gheena S, Pratibha R, Abilasha R, Hannah R, et al. Overexpression of HNRNPA2B1 is Associated with Poor Prognosis in Head and Neck Squamous Cell Carcinoma [Internet]. International Journal of Current Research and Review. 2020. p. 15–8. Available from: <http://dx.doi.org/10.31782/ijcrr.2020.122502>
 13. Princeton B, Santhakumar P, Prathap L. Awareness on Preventive Measures taken by Health Care Professionals Attending COVID-19 Patients among Dental Students. Eur J Dent. 2020 Dec;14(S 01):S105–9.
 14. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: randomized controlled trial. Clin Oral Investig. 2020 Sep;24(9):3275–80.
 15. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. J Oral Pathol Med. 2019 Apr;48(4):299–306.
 16. R H, Hannah R, Ramani P, Ramanathan A, Jancy MR, Gheena S, et al. CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene [Internet]. Vol. 130, Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology. 2020. p. 306–12. Available from: <http://dx.doi.org/10.1016/j.oooo.2020.06.021>
 17. Antony JVM, Ramani P, Ramasubramanian A, Sukumaran G. Particle size penetration rate and effects of smoke and smokeless tobacco products - An invitro analysis. Heliyon. 2021 Mar;7(3):e06455.
 18. Sarode SC, Gondivkar S, Sarode GS, Gadbaile A, Yuwanati M. Hybrid oral potentially malignant disorder: A neglected fact in oral submucous fibrosis. Oral Oncol. 2021 Jun

16;105390.

19. Hannah R, Ramani P, WM Tilakaratne, Sukumaran G, Ramasubramanian A, Krishnan RP. Author response for “Critical appraisal of different triggering pathways for the pathobiology of pemphigus vulgaris—A review” [Internet]. Wiley; 2021. Available from: <https://publons.com/publon/47643844>
20. Chandrasekar R, Chandrasekhar S, Sundari KKS, Ravi P. Development and validation of a formula for objective assessment of cervical vertebral bone age. *Prog Orthod*. 2020 Oct 12;21(1):38.
21. Subramanyam D, Gurunathan D, Gaayathri R, Vishnu Priya V. Comparative evaluation of salivary malondialdehyde levels as a marker of lipid peroxidation in early childhood caries. *Eur J Dent*. 2018 Jan;12(1):67–70.
22. Jeevanandan G, Thomas E. Volumetric analysis of hand, reciprocating and rotary instrumentation techniques in primary molars using spiral computed tomography: An in vitro comparative study. *Eur J Dent*. 2018 Jan;12(1):21–6.
23. Ponnulakshmi R, Shyamaladevi B, Vijayalakshmi P, Selvaraj J. In silico and in vivo analysis to identify the antidiabetic activity of beta sitosterol in adipose tissue of high fat diet and sucrose induced type-2 diabetic experimental rats. *Toxicol Mech Methods*. 2019 May;29(4):276–90.
24. Sundaram R, Nandhakumar E, Haseena Banu H. Hesperidin, a citrus flavonoid ameliorates hyperglycemia by regulating key enzymes of carbohydrate metabolism in streptozotocin-induced diabetic rats. *Toxicol Mech Methods*. 2019 Nov;29(9):644–53.
25. Alsawalha M, Rao CV, Al-Subaie AM, Haque SKM, Veeraraghavan VP, Surapaneni KM. Novel mathematical modelling of Saudi Arabian natural diatomite clay. *Mater Res Express*. 2019 Sep 4;6(10):105531.
26. Yu J, Li M, Zhan D, Shi C, Fang L, Ban C, et al. Inhibitory effects of triterpenoid betulin on inflammatory mediators inducible nitric oxide synthase, cyclooxygenase-2, tumor necrosis factor-alpha, interleukin-6, and proliferating cell nuclear antigen in 1, 2-dimethylhydrazine-induced rat colon carcinogenesis. *Pharmacogn Mag*. 2020;16(72):836.
27. Shree KH, Hema Shree K, Ramani P, Herald Sherlin, Sukumaran G, Jeyaraj G, et al. Saliva as a Diagnostic Tool in Oral Squamous Cell Carcinoma – a Systematic Review with Meta Analysis [Internet]. Vol. 25, *Pathology & Oncology Research*. 2019. p. 447–53. Available from: <http://dx.doi.org/10.1007/s12253-019-00588-2>
28. Zafar A, Sherlin HJ, Jayaraj G, Ramani P, Don KR, Santhanam A. Diagnostic utility of touch imprint cytology for intraoperative assessment of surgical margins and sentinel lymph nodes in oral squamous cell carcinoma patients using four different cytological stains. *Diagn Cytopathol*. 2020 Feb;48(2):101–10.
29. Karunagaran M, Murali P, Palaniappan V, Sivapathasundharam B. Expression and distribution pattern of podoplanin in oral submucous fibrosis with varying degrees of dysplasia – an immunohistochemical study [Internet]. Vol. 42, *Journal of Histotechnology*. 2019. p. 80–6. Available from: <http://dx.doi.org/10.1080/01478885.2019.1594543>
30. Sarode SC, Gondivkar S, Gadmail A, Sarode GS, Yuwanati M. Oral submucous fibrosis and

heterogeneity in outcome measures: a critical viewpoint. *Future Oncol.* 2021 Jun;17(17):2123–6.

31. Raj Preeth D, Saravanan S, Shairam M, Selvakumar N, Selestin Raja I, Dhanasekaran A, et al. Bioactive Zinc(II) complex incorporated PCL/gelatin electrospun nanofiber enhanced bone tissue regeneration. *Eur J Pharm Sci.* 2021 May 1;160:105768.
32. Prithiviraj N, Yang GE, Thangavelu L, Yan J. Anticancer Compounds From Starfish Regenerating Tissues and Their Antioxidant Properties on Human Oral Epidermoid Carcinoma KB Cells. In: PANCREAS. LIPPINCOTT WILLIAMS & WILKINS TWO COMMERCE SQ, 2001 MARKET ST, PHILADELPHIA ...; 2020. p. 155–6.
33. Epelman I, Murray PE, Garcia-Godoy F, Kuttler S, Namerow KN. A practitioner survey of opinions toward regenerative endodontics. *J Endod.* 2009;35(9):1204-10.
34. . Utneja S, Nawal RR, Ansari MI, Talwar S, Verma M. A survey of attitude and opinions of endodontic residents towards regenerative endodontics. *J Conserv Dent.* 2013;16(4):314-8.
35. Britto.R.P, Danraj.M. Stem cell therapy in oral and maxillofacial region. *International Journal of Advancements in Research & Technology.* 2014;3(8).
36. Sunil P, Manikandhan R, Muthu M, Abraham S. Stem cell therapy in oral and maxillofacial region: An overview. *J Oral Maxillofac Pathol.* 2012;16(1):58-63. doi:10.4103/0973-029X.92975