

Original Research Article

PRELIMINARY PHYTOCHEMICAL ANALYSIS AND ASSESSMENT OF ANTIOXIDANT AND ANTI CHOLESTEROL ACTIVITY OF ETHANOLIC LEAF EXTRACT OF *ACALYPHA INDICA* LINN.

ABSTRACT:

Background: *Acalypha indica* Linn. is a weed plant found in shady wetlands. It is commonly called as kuppaimeni in Tamil, belonging to the family *Euphorbiaceae*. It is found to be one of the ~~greatest~~ most important Indian plants with anti-venom, antibacterial, antiviral and antimicrobial properties.

Aim: Preliminary phytochemical analysis and assessment of antioxidant and anti cholesterol activity of ethanolic leaf extract of *Acalypha indica* Linn

Materials and Methods: Phytochemical screening and *in vitro* antioxidant and hypolipidemic potential of *Acalypha indica* Linn. was analysed as per the standard methods. The data were analyzed statistically by one-way analysis of variance (ANOVA) followed by Duncan's multiple range test and it was used to see the statistical significance among the groups. The results with the $p < 0.05$ level were considered to be statistically significant.

Results: Ethanolic leaf extract of *Acalypha indica* Linn exhibited a significant antioxidant and hypolipidemic activity and increased in a dose dependent manner as compared to the standard drug.

Conclusion: Many researchers are working towards replacing synthetic drugs with herbal alternatives. Herbal extracts are natural and will aid in longevity.

keywords : *Acalypha indica*; weed plant; standard drugs; In-vitro analysis of antioxidant and anti cholesterol; innovative technology; novel method.

Running title: Antioxidant and anti cholesterol properties of *Acalypha indica* Linn.

Introduction:

Acalypha indica Linn. is a weed plant found in shady wetlands. It is commonly called as kuppaimeni in Tamil, belonging to the family *Euphorbiaceae*. It is found to be one of the greatest-most important Indian plants with anti-venom, antibacterial, antiviral and antimicrobial properties (Gupta, 2010). Indian medicine like Ayurveda, siddha, utilize the extract to treat diseases like Pneumonia, asthma, wounds, allergies, cleansing liver and kidney etc. As a folk medicine, the extract is used as facial masks and other utilities (Durai Pandian *et al.*, 2007). It is considered as one of the greatest gifts of mother earth.

This plant produces a wide range of bioactive molecules which result in a cure to are used to treat a wide range of diseases. The most important phytoconstituents of the extract are alkaloid, flavonoid, tannin and phenolic compounds (Awadh *et al.*, 2017). These compounds are found in several parts of plants like the roots, shoots, leaves and bark of the plant. These phytochemicals were have been analysed and studied by many researchers in many parts of the world (Pauranik *et al.* 1986) (Rai, Zachariah and Shrivastava, 1989). Catlow (Catlow, 1848), has predicted-determined that the variety of biomolecules produced by the plant can act as an antimicrobial and antioxidant agents (Catlow, 1848) which constitute 80% of most popular and healthy natural remedies.

This plant has been reported to contain various clinical constituents like kaempferol glucoside, clitorin, mauritian, nicotiflorin and borodin. Which were isolated and few in recent studies of flowers and leaves (Shima, Hisada and Inagaki, 1972); (Doughari, 2007).

Hyperlipidemia is a major risk factor causing cardiovascular disease which can result in cardiac arrest; it is responsible for the cardiac risks associated with one third of the world-wide population all over the globe. It is caused by the over deposition of cholesterol and triglyceride, which may have resulted due to the unhealthy food practice which has resulted in the production of free radicals that cause fast aging and decreased physical flexibility (Nelson, 2013); (Ohlsen and Gaughran, 2011).

2020); (Theertha *et al.*, 2020); (Veeraraghavan, Periadurai, *et al.*, 2021). In the current study, *in vitro* antioxidant and hypolipidemic potential of *Acalypha indica* Linn. was analysed.

Materials and Methods:

1. Phytochemical Screening test

Test for phlobatannin

1.0 ml of the extract was treated with 1ml of 1% HCl and boiled for 10 mins. The formation of red color precipitate indicates the presence of phlobatannin.

Test for Carbohydrates

Three to five drops of Molisch reagent was added with 1.0 ml of the extract and then 1.0 ml of concentrated sulphuric acid was added carefully through the side of the test tube. The mixture was then allowed to stand for two minutes and diluted with 5.0 ml of distilled water. The development of a red or dull violet ring at the junction of the liquids showed the presence of carbohydrates.

Test for Flavonoids

Few drops of 1% liquid ammonia were taken in a test tube and along with it 1.0 ml of the extract was added resulting in the formation of yellow color thereby indicating the presence of flavonoids.

Test for Alkaloids

2.0 ml of sample was mixed with 2.0 ml of HCl. Then 6 drops of HCN was added and further 2 drops of picric acid was added that resulted in a creamish pale yellow ppt-precipitate indicating the presence of alkaloids.

Test for Terpenoids

2.0 ml of sample along with 2.0 ml of chloroform and 3.0 ml of ~~concentrated~~ H₂SO₄ was added. A red color ppt-precipitate obtained indicates the presence of terpenoids.

Test for proteins

~~One milliliter~~ 1.0 ml of ninhydrin was dissolved in 1.0 ml of acetone and then a small amount of extract was added with ninhydrin. The formation of purple colour revealed the presence of protein.

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Font color: Auto

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, Font color: Auto

Formatted: Font: (Default) Times New Roman, Font color: Auto

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, Font color: Auto

Comment [MOU1]: How much is a small amount?

Detection of saponins

Foam test: A fraction of the extract was vigorously shaken with water and observed for persistent foam.

Test for steroids

~~One milliliter~~ 1.0 ml of chloroform was mixed with 1.0 ml of extract and then ten drops of acetic anhydride and five drops of concentrated sulphuric acid were added and mixed. The formation of dark red colour or dark pink colour indicates the presence of steroids.

2. DPPH free radical scavenging activity of *Acalypha indica*

Scavenging of 2, 2-Diphenyl-1-picrylhydrazyl (DPPH) radicals was assessed by the method of Hatano et al, (1989). DPPH solution (1.0 ml) was added to 1.0 ml of extract at different concentrations (0.1 to 0. 5mg/ml). The mixture was kept at room temperature for 50 minutes and the activity was measured at 517 nm. Ascorbic acid at the same concentrations was used as standard. The capability to scavenge the DPPH radical was calculated and expressed in percentage (%) using following formula:

$$\text{DPPH radical scavenging (\%)} = \frac{\text{Control OD} - \text{Sample OD}}{\text{Control OD}} \times 100$$

3. In vitro anti-cholesterol activity of *Acalypha indica*

The anti-cholesterol assay was carried out as described as per the kit method (Spinreact, S.A.U-Ctra Santa Coloma, Girona, Spain). Cholesterol was dissolved in chloroform at a concentration of 2.5 mg mL/ml. ~~Ten microliter~~ 10.0 ml of the extract was pipetted into a microtiter plate followed by the addition of 2000 µL of R1 reagent and 10 µL of cholesterol as sample. ~~Twenty microliter~~ 20.0 l of distilled water and 2000 µL of R1 reagent were used as blank. Negative control consisted of 20 µL cholesterol and 2.0 ml R1; standard consisted of 20 µL simvastatin and 2000 µL R1 reagent. The contents were incubated between 0-30 min at room temperature and the absorbance was read at 500 nm in a UV-Vis spectrophotometer against the blank reagent blank. Anti-cholesterol assay of the extract was calculated using the following equation:

Comment [MOU2]: Is this correct?

Comment [MOU3]: Blank? What is the control? R1 Reagent? Water? Something else?

$$\text{Inhibition (\%)} = \frac{\text{Negative Control} - \text{Sample}}{\text{Negative Control}} \times 100$$

$$\text{Inhibition (\%)} = \frac{\text{Negative control} - \text{Sample}}{\text{Negative control}} \times 100$$

Negative control

Comment [MOU4]: Could use Equation Editor in Word to create a better looking equation.

Statistical Analysis

The data were subjected to statistical analysis using one-way analysis of variance (ANOVA) and Duncan's multiple range test to assess the significance of individual variations between the groups. In Duncan's test, significance was considered at the level of $p < 0.05$.


Formatted: Left

Results and Discussion:

Table 1: Phytochemical screening of *Acalypha indica*

Phytochemicals	Leaf extract
Proteins	(-)
Amino acids	(+)(+)(+)
Terpenoids	(+)(+)
Flavonoids	(+)
Alkaloids	(+)
Saponins	(+)
Steroids	(+)

Comment [MOU5]: How does +++ differ from + or ++. What does it take to get three +?



(Wu *et al.*, 2019)

FIGURE 1: The graph depicts the association between *in vitro* antioxidant activity of *Acalypha indica* and ~~standard drug simvastatin~~ (vitamin C). X-axis represents ~~activity concentration~~ of drugs and Y-axis percentage of DPPH scavenging activity. Blue represents ~~the~~ activity of ~~std drug simvastatin~~ Vitamin C and orange represents ~~the~~ activity of *Acalypha indica*. The graph denotes that ~~standard drug simvastatin~~ Vitamin C shows greater activity than that of ethanolic extract of *Acalypha indica*. Each line Represents Mean \pm SEM of 3 independent observations. Significance at $p \leq 0.05$.

Formatted: Font: Italic

Comment [MOU6]: Simvastatin is not Vitamin C

Formatted: Font: (Default) Times New Roman




FIGURE 2: The graph depicts the association between *in vitro* antioxidant activity of *Acalypha indica* and standard drug-simvastatin (vitamin C). X-axis represents activity of drugs and Y-axis percentage cholesterol scavenged. Blue represents activity of the std drug-*Acalypha indica* extract and brown-red represents activity of simvastatin. The graph denotes that standard drug simvastatin shows greater anti-cholesterol activity than that of ethanolic extract of *Acalypha indica*. Each line Represents Mean \pm SEM of 3 independent observations. Significance at $p \leq 0.05$.

Formatted: Font: Italic

Comment [MOU7]: Simvastatin is not Vitamin C

Formatted: Font: (Default) Times New Roman

The result of qualitative phytochemical analysis of *Acalypha indica* leaf extracts are given in table 1. The result reveals a strong presence of flavonoid, alkaloids, saponins, steroids and amino acids. The DPPH radical scavenging activity revealed the antioxidant potential of the plant extract. Antioxidant potential was significantly exhibited by the *Acalypha indica* extract with $Ic_{50} = 290$ μ g/ml and increased in a dose dependent manner as compared to the standard drug Vitamin C (Ic_{50} of Vitamin-C= 210 μ g/ml)(Figure 1)

The result also revealed a strong in vitro anti-cholesterol potential of ethanolic extract of *Acalypha Indica.*, which is slightly lower than that of ~~standard~~-simvastatin. Ic50 of *Acalypha indica* was found to be 300~~ugug~~/ml and the ~~standard drug~~-simvastatin exhibited Ic50 at 260ug/ml. (Figure 2).

Previous studies strongly suggest that the plant extracts which are rich in phenolic compounds can fight free radicals effectively, since aromatic rings in phenolic compounds can easily accept or donate an electron.(Jennings, 1981).

Thus, in the present study, the plant extract was found to be rich in phenolic compounds such as alkaloids and flavonoids, which forms the basis for its strong antioxidant property.

Cholesterol is a steroid which is produced endogenously and also taken from the diet. The biosynthesis of cholesterol is usually regulated by regulating the key enzyme HMG CoA reductase(Nagpal, 2018). Standard anti-cholesterol drugs act by inhibiting these key enzymes. Though the standard drugs are more potent in reducing cholesterol levels, it also exhibits various side effects if consumed for a longer period of time(Huang and Freter, 2016).

The study showed that the in vitro antioxidant property *Acalypha indica* extract is Ic50 = 290 ug/ml which is similar with (Reddy and Reddy Prasad Reddy, 2012; Dwijayanti, Farida and Purwaningsih, 2018; Zulkifliet *al.*, 2018). The study also found that in vitro anti-cholesterol potential of ethanolic extract of *Acalypha i*~~ndica.~~, which is slightly lower than that of standard simvastatin i.e.Ic50 of *Acalypha indica*, which is found to be similar with (R, Jayaprakasam and Ravi, 2013);(El-Shemy, 2017). From the study, it was evident that the plant extract exhibited a strong anti-cholesterol property. Further research needs to be done to explore various medicinal properties of the extract.

Conclusion

The ethanolic extract of *Acalypha indica Linn.* possessed potent antioxidant and anticholesterol activity.~~Which will be natural and may be with or without any~~ It is unknown whether this extract has any of the side effects ~~seen inof std drug~~ simvastatin. Nowadays people are moving towards the use of herbal products, as they are feeling fear about the side effects it may cause. So further research is needed to convert the extract into active drug formulation.

COMPETING INTERESTS DISCLAIMER:

Formatted: Not Highlight

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

Formatted: Not Highlight

NOTE:

The study highlights the efficacy of "herbal", "Ayurveda", "siddha" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

Reference

Awadh, A. *et al.* (2017) 'Does Zika Virus Cause Microcephaly - Applying the Bradford Hill Viewpoints', *PLoS currents*, 9. doi: 10.1371/currents.outbreaks.2fced6e886074f6db162a00d4940133b.

Babu, S. and Jayaraman, S. (2020) 'An update on β -sitosterol: A potential herbal nutraceutical for diabetic management', *Biomedicine & pharmacotherapy = Biomedecine&pharmacotherapie*, 131, p. 110702.

Barma, M. D. *et al.* (2021) 'Inhibition of *Streptococcus mutans*, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish', *Archives of oral biology*, 126, p. 105132.

Catlow, A. (1848) 'Popular field botany; containing a familiar and technical description of the plants most common to the various localities of the British Isles, adapted to the study of either the artificial or natural systems'. doi: 10.5962/bhl.title.134959.

Chen, F. *et al.* (2019) '6-shogaol, a active ~~eonstituents~~constituents of ginger prevents UVB radiation mediated inflammation and oxidative stress through modulating Nrf2 signaling in human epidermal keratinocytes (HaCaT cells)', *Journal of photochemistry and photobiology. B, Biology*, 197, p. 111518.

Doughari, J. H. (2007) 'Antimicrobial Activity of *Tamarindus indica* Linn', *Tropical Journal of*

Pharmaceutical Research. doi: 10.4314/tjpr.v5i2.14637.

Durai Pandian, J. *et al.* (2007) 'Stroke and thrombolysis in developing countries', *International journal of stroke: official journal of the International Stroke Society*, 2(1), pp. 17–26.

Dwijayanti, A., Farida, S. and Purwaningsih, E. H. (2018) 'Comparing Anti Aging Potential Between *Centella asiatica* and *Acalypha indica*: Focus on Forelimb Muscle Strength', *Advanced Science Letters*, pp. 6058–6060. doi: 10.1166/asl.2018.12621.

El-Shemy, H. (2017) *Aromatic and Medicinal Plants: Back to Nature*. BoD – Books on Demand.

Eyob, S. *et al.* (2008) 'Traditional medicinal uses and essential oil composition of leaves and rhizomes of korarima (*Aframomum corrorima* (Braun) P.C.M. Jansen) from southern Ethiopia', *South African Journal of Botany*, pp. 181–185. doi: 10.1016/j.sajb.2007.10.007.

Gothai, S. *et al.* (2018) 'Pharmacological insights into antioxidants against colorectal cancer: A detailed review of the possible mechanisms', *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*, 107, pp. 1514–1522.

Gupta, R. K. (2010) *Medicinal & Aromatic Plants: With Colour Plates (hb)*.

Halliwell, B. (1994) 'Vitamin C: the key to health or a slow-acting carcinogen?', *Redox report: communications in free radical research*, 1(1), pp. 5–9.

Han, X. *et al.* (2019) 'Anticarcinogenic potential of gold nanoparticles synthesized from *Trichosanthes kirilowii* in colon cancer cells through the induction of apoptotic pathway', *Artificial cells, nanomedicine, and biotechnology*, 47(1), pp. 3577–3584.

Huang, C. and Freter, C. (2016) *Cholesterol Lowering Therapies and Drugs*. BoD – Books on Demand.

Ide, H. *et al.* (1990) 'Effects of HMG-CoA Reductase Inhibitor (MK-733) on Plasma Lipids and Steroid Hormones', *The Journal of Japan Atherosclerosis Society*, pp. 767–774. doi: 10.5551/jat1973.18.7-8_767.

Liu Y.-W. *et al.* (2008) '[A test of screening to predict lung cancer among dust-exposed tin miners with sputum imaging cytometry]', *Zhonghua lao dong wei sheng zhi ye bing za zhi = Zhonghualao dong weisheng zhi yebing zazhi = Chinese journal of industrial hygiene and occupational diseases*, 26(4), pp. 203–207.

Li, Z. *et al.* (2020) 'Apoptotic induction and anti-metastatic activity of eugenol encapsulated chitosan nanopolymer on rat glioma C6 cells via alleviating the MMP signaling pathway', *Journal of Photochemistry and Photobiology B: Biology*, p. 111773. doi: 10.1016/j.jphotobiol.2019.111773.

Malaikolundhan, H. *et al.* (2020) 'Anticarcinogenic effect of gold nanoparticles synthesized from *Albizia lebbek* on HCT-116 colon cancer cell lines', *Artificial cells, nanomedicine, and biotechnology*, 48(1), pp. 1206–1213.

Mickymaray, S. *et al.* (2021) 'Rhaponticin suppresses osteosarcoma through the inhibition of PI3K-Akt-mTOR pathway', *Saudi journal of biological sciences*, 28(7), pp. 3641–3649.

Nagpal, M. L. (2018) *Cholesterol: Good, Bad and the Heart*. BoD – Books on Demand.

Nelson, R. H. (2013) 'Hyperlipidemia as a Risk Factor for Cardiovascular Disease', *Primary Care: Clinics in Office Practice*, pp. 195–211. doi: 10.1016/j.pop.2012.11.003.

Ohlsen, R. and Gaughran, F. (2011) 'Schizophrenia: A major risk factor for cardiovascular disease', *British Journal of Cardiac Nursing*, pp. 228–232. doi: 10.12968/bjca.2011.6.5.228.

Rai, G. P., Zachariah, K. and Shrivastava, S. (1989) 'Comparative efficacy of indirect haemagglutination test, indirect fluorescent antibody test and enzyme linked immunosorbent assay in serodiagnosis of typhoid fever', *The Journal of tropical medicine and hygiene*, 92(6), pp. 431–434.

Rajendran, P. *et al.* (2020) 'Consumption of reused vegetable oil intensifies BRCA1 mutations', *Critical reviews in food science and nutrition*, pp. 1–8.

Reddy, T. P. and Reddy Prasad Reddy, T. (2012) 'Exploring the Anti-inflammatory and Anti-cancer compounds from the leaves of *Acalypha indica*', *IOSR Journal of Pharmacy and Biological Sciences*, pp. 1–7. doi: 10.9790/3008-0420107.

R, J., Jayaprakasam, R. and Ravi, T. K. (2013) 'EVALUATION OF ANTI ARTHRITIC ACTIVITY OF THE ROOT EXTRACT OF ACALYPHA INDICA LINN. USING IN VITRO TECHNIQUES', *International Journal of Phytopharmacy*. doi: 10.7439/ijpp.v2i6.36.

Samuel, S. R. (2021) 'Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life?', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 31(2), pp. 285–286.

Samuel, S. R. *et al.* (2021) 'Dental pain, parental SARS-CoV-2 fear and distress on quality of life of 2 to 6 year-old children during COVID-19', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 31(3), pp. 436–441.

Sathya, S. *et al.* (2020) 'An in vitro study on hexavalent chromium [Cr(VI)] remediation using iron oxide nanoparticles based beads', *Environmental Nanotechnology, Monitoring & Management*, 14, p. 100333.

Shima, K., Hisada, S. and Inagaki, I. (1972) 'Studies on the Constituents of Anodendron affine DURCE. IV. : Isolation of Kaempferol, Astragalol and Dambonitol from Leaves', *YAKUGAKU ZASSHI*, pp. 507–509. doi: 10.1248/yakushi1947.92.4_507.

Tang, Y. *et al.* (2021) 'Osteogenic differentiation and mineralization potential of zinc oxide nanoparticles from *Scutellaria baicalensis* on human osteoblast-like MG-63 cells', *Materials Science and Engineering: C*, p. 111656. doi: 10.1016/j.msec.2020.111656.

Teja, K. V. and Ramesh, S. (2020) 'Is a filled lateral canal – A sign of superiority?', *Journal of Dental Sciences*, pp. 562–563. doi: 10.1016/j.jds.2020.02.009.

Theertha, M. *et al.* (2020) 'Innate lymphoid cells: Potent early mediators of the host immune response during sepsis', *Cellular & molecular immunology*, 17(10), pp. 1114–1116.

Veeraraghavan, V. P., Hussain, S., *et al.* (2021) 'A Comprehensive and Critical Review on Ethnopharmacological Importance of Desert Truffles: *Terfeziaclavaryi*, *Terfeziaboudieri*, and *Tirmania nivea*', *Food Reviews International*, pp. 1–20. doi: 10.1080/87559129.2021.1889581.

Veeraraghavan, V. P., Periadurai, N. D., *et al.* (2021) 'Green synthesis of silver nanoparticles from aqueous extract of *Scutellariabarabata* and coating on the cotton fabric for antimicrobial applications and wound healing activity in fibroblast cells (L929)', *Saudi journal of biological sciences*, 28(7), pp. 3633–3640.

Willcox, B. J. *et al.* (2004) 'C-reactive protein, cardiovascular disease and stroke: new roles for an old biomarker', *Expert review of neurotherapeutics*, 4(3), pp. 507–518.

Wu, F. *et al.* (2019) 'Biologically synthesized green gold nanoparticles from Siberian ginseng induce growth-inhibitory effect on melanoma cells (B16)', *Artificial cells, nanomedicine, and biotechnology*, 47(1), pp. 3297–3305.

Yang, Z. *et al.* (2020) 'Piperine loaded zinc oxide nanocomposite inhibits the PI3K/AKT/mTOR signaling pathway via attenuating the development of gastric carcinoma: In vitro and in vivo studies', *Arabian Journal of Chemistry*, 13(5), pp. 5501–5516.

Yin, Z. *et al.* (2021) 'Potential chemotherapeutic effect of betalain against human non-small cell lung cancer through PI3K/Akt/mTOR signaling pathway', *Environmental toxicology*, 36(6), pp. 1011–1020.

Young, I. S. and McEneaney, J. (2001) 'Lipoprotein oxidation and atherosclerosis', *Biochemical Society transactions*, 29(Pt 2), pp. 358–362.

Zulkifli, C. *et al.* (2018) 'Anti-Oxidative Potential of *Acalypha indica* L. Root Extract on Brain-Derived Neurotrophic Factor Levels in Old Sprague-Dawley Rats', *Proceedings of BROMO Conference*. doi: 10.5220/0008357400580061.

Awadh, A. *et al.* (2017) 'Does Zika Virus Cause Microcephaly - Applying the Bradford Hill Viewpoints', *PLoS currents*, 9. doi: 10.1371/currents.outbreaks.2fced6e886074f6db162a00d4940133b.

Catlow, A. (1848) 'Popular field botany; containing a familiar and technical description of the plants most common to the various localities of the British Isles, adapted to the study of either the artificial or natural systems'. doi: 10.5962/bhl.title.134959.

Doughari, J. H. (2007) 'Antimicrobial Activity of *Tamarindus indica* Linn', *Tropical Journal of Pharmaceutical Research*. doi: 10.4314/tjpr.v5i2.14637.

- Durai Pandian, J. et al. (2007) 'Stroke and thrombolysis in developing countries', *International journal of stroke: official journal of the International Stroke Society*, 2(1), pp. 17–26.
- Dwijayanti, A., Farida, S. and Purwaningsih, E. H. (2018) 'Comparing Anti Aging Potential Between *Centella asiatica* and *Acalypha indica*: Focus on Forelimb Muscle Strength', *Advanced Science Letters*, pp. 6058–6060. doi: 10.1166/asl.2018.12621.
- El-Shemy, H. (2017) *Aromatic and Medicinal Plants: Back to Nature*. BoD – Books on Demand.
- Eyob, S. et al. (2008) 'Traditional medicinal uses and essential oil composition of leaves and rhizomes of korarima (*Aframomum corrorima* (Braun) P.C.M. Jansen) from southern Ethiopia', *South African Journal of Botany*, pp. 181–185. doi: 10.1016/j.sajb.2007.10.007.
- Gupta, R. K. (2010) *Medicinal & Aromatic Plants: With Colour Plates* (hb).
- Halliwell, B. (1994) 'Vitamin C: the key to health or a slow-acting carcinogen?', *Redox report: communications in free radical research*, 1(1), pp. 5–9.
- Huang, C. and Freter, C. (2016) *Cholesterol Lowering Therapies and Drugs*. BoD – Books on Demand.
- Ide, H. et al. (1990) 'Effects of HMG-CoA Reductase Inhibitor (MK-733) on Plasma Lipids and Steroid Hormones', *The Journal of Japan Atherosclerosis Society*, pp. 767–774. doi: 10.5551/jat1973.18.7-8_767.
- Liu Y.-W. et al. (2008) '[A test of screening to predict lung cancer among dust-exposed tin miners with sputum imaging cytometry]', *Zhonghua lao dong wei sheng zhi ye bing za zhi = Zhonghualaodongweishengzhiyebingzazhi = Chinese journal of industrial hygiene and occupational diseases*, 26(4), pp. 203–207.
- Nagpal, M. L. (2018) *Cholesterol: Good, Bad and the Heart*. BoD – Books on Demand.
- Nelson, R. H. (2013) 'Hyperlipidemia as a Risk Factor for Cardiovascular Disease', *Primary Care: Clinics in Office Practice*, pp. 195–211. doi: 10.1016/j.pop.2012.11.003.
- Ohlsen, R. and Gaughran, F. (2011) 'Schizophrenia: A major risk factor for cardiovascular disease', *British Journal of Cardiac Nursing*, pp. 228–232. doi: 10.12968/bjca.2011.6.5.228.
- Rai, G. P., Zachariah, K. and Shrivastava, S. (1989) 'Comparative efficacy of indirect haemagglutination test, indirect fluorescent antibody test and enzyme linked immunosorbent assay in serodiagnosis of typhoid fever', *The Journal of tropical medicine and hygiene*, 92(6), pp. 431–434.
- Reddy, T. P. and Reddy Prasad Reddy, T. (2012) 'Exploring the Anti-inflammatory and Anti-cancer compounds from the leaves of *Acalypha indica*', *IOSR Journal of Pharmacy and Biological Sciences*, pp. 1–7. doi: 10.9790/3008-0420107.
- R, J., Jayaprakasam, R. and Ravi, T. K. (2013) 'EVALUATION OF ANTI ARTHRITIC

ACTIVITY OF THE ROOT EXTRACT OF ACALYPHA INDICA LINN. USING IN VITRO TECHNIQUES', International Journal of Phytopharmacy. doi: 10.7439/ijpp.v2i6.36.

Shima, K., Hisada, S. and Inagaki, I. (1972) 'Studies on the Constituents of Anodendron affine DURCE. IV. : Isolation of Kaempferol, Astragalol and Dambonitol from Leaves', YAKUGAKU ZASSHI, pp. 507–509. doi: 10.1248/yakushi1947.92.4_507.

Willcox, B. J. et al. (2004) 'C-reactive protein, cardiovascular disease and stroke: new roles for an old biomarker', Expert review of neurotherapeutics, 4(3), pp. 507–518.

Young, I. S. and McEneny, J. (2001) 'Lipoprotein oxidation and atherosclerosis', Biochemical Society transactions, 29(Pt 2), pp. 358–362.

Zulkifli, C. et al. (2018) 'Anti-Oxidative Potential of Acalypha indica L. Root Extract on Brain-Derived Neurotrophic Factor Levels in Old Sprague-Dawley Rats', Proceedings of BROMO Conference. doi: 10.5220/0008357400580061.

UNDER PEER REVIEW