

COMPARATIVE ANALYSIS OF ANTI GOUT ACTIVITIES OF ETHANOLIC EXTRACT OF *ALTERNANTHERA SESSILIS* AND *MORINGA OLEIFERA*

ABSTRACT :

INTRODUCTION:

Ethanollic extracts of *Alternanthera sessilis* and *Moringa oleifera* were screened for the presence of different classes of phytochemicals. The phytochemical screening revealed the presence of amino acids, flavonoids, alkaloids, terpenoids, saponins and steroids. Gout is the type of arthritis that causes painful inflammation and develops due to overproduction of urate.

AIM: To analyse the anti gout activities of ethanolic extract of *Alternanthera sessilis* and *moringa oleifera*.

MATERIALS AND METHODS:

Ethanollic extract of *Alternanthera sessilis* and *Moringa oleifera* were prepared as per the standard methods and used for the assessment of preliminary phytochemical screening, antioxidants and anti gout activities. The data were analyzed statistically by a one-way analysis of (ANOVA) followed by Duncan's multiple range test was used to see the statistical significance among the group. The results with the $p < 0.05$ level were considered to be statistically significant.

RESULT: Antioxidant and Anti gout potential of the ethanolic extract of *Moringa oleifera* was found to be significantly more than the ethanolic extract of *Alternanthera sessilis*.

CONCLUSION: Both the ethanolic extracts exhibited anti gout activity, further studies are needed to validate the herbal extracts as a drug formulation.

KEYWORDS : Extract, phytochemicals, antioxidants, xanthine oxidase, innovative technology; novel method

RUNNING TITLE : Antigout activities of ethanolic extracts of *Alternanthera sessilis* and *Moringa oleifera*

INTRODUCTION

Plants possess a wide variety of secondary metabolites. These plants are rich in antioxidant activity and other therapeutic values(1). Medicinal plants are used for the treatment of various ailments(2). For drug synthesis it serves as a starting material. *Moringa oleifera* belongs to the Moringaceae family. It is a native to indian subcontinent(2,3). The plant is widely grown in tropical and subtropical countries. *Moringa oleifera* is a highly valued plant for its nutrition and medication. Moringa trees are commonly known as drumsticks. It is a traditional medicinal source (1). The parts of the moringa trees like pods, seeds and leaves are edible. Moringa seeds are used in spice, cosmetic oils, food etc(4). It is used as livestock feed and has several medicinal uses (2). The digestive ability of the plant is affected by the chemical composition of the fibre. (5)

The nutritional value of plants depends on the environment, cultivation method, genetic background(4,6). It shows antipyretic, analgesic and wound healing activities. The leaves contain 14 amino acids and leaves are the natural boosters, vitamin A,C,E, proteins, magnesium, calcium and carbohydrates(7). *Moringa Oleifera* can serve as a wide variety of biological activity and antioxidants for humans and animals (8)

Gout is a type of arthritis which causes painful inflammation. Joint pain is caused due to elevation of uric acid in the blood which triggers the formation of crystals. Xanthine oxidase catalyses the oxidative hydroxylation of hypoxanthine to xanthine to uric acid.(9). The further catalysis is done by enzyme uricase which converts uric acid to highly soluble allantoin which is secreted in urine. Symptoms of gout can be intense joint pains, lingering discomfort, inflammation, and redness(10). Development of gout has several risk factors including hyperuricemia, age, dietary factors, metabolic syndrome, diuretic, chronic renal disease(11). *Alternanthera sessilis* belongs to amaranthaceae family. It is a perennial herb and an aquatic plant. It is found in sterms, wetlands, marshy areas, reservoirs, and river banks. It is also used for gastrointestinal problems and the branches row from the root upto 50cms long (8,12). It is used for the treatment of skin disease, fever, cuts, gonorrhea, diarrhea, burning sensation, liver & spleen disease (8). Aerial part of the plant is the source of vitamins, minerals and antioxidants. It contains bio active phytoconstituents. It is rich in nutrition, and contains iron and vitamin A, serves as food and is also used in medicine(13). There is not much research done on antioxidant and anti gout properties of the plant. Our team has extensive knowledge and research experience that has translate into high quality publications (14),(15),(16),(17),(18),(19),(20),(21),(22),(23),(24),(25),(26),(27),(28),(29),(30),(31),(32),(33). The main aim of the study is to analyse anti gout activities of ethanolic extract of *Alternanthera sessilis* and *Moringa oleifera*.

MATERIALS AND METHODS:

1. Phytochemical Screening test

Test for phlobatannin

1ml of the extract was treated with 1ml of 1% HCl and boiled for 10 mins. The formation of red color precipitate indicates the presence of phlobatannin.

Test for Carbohydrates

Three to five drops of Molisch reagent was added with 1 mL of the extract and then 1 mL of concentrated sulphuric acid was added carefully through the side of the test tube. The mixture was then allowed to stand for two minutes and diluted with 5 mL of distilled water. The development of a red or dull violet ring at the junction of the liquids showed the presence of carbohydrates.

Test for Flavonoids

Few drops of 1% liquid ammonia were taken in a test tube and along with it 1ml of the extract was added resulting in the formation of yellow color thereby indicating the presence of flavonoids.

Test for Alkaloids

2ml of sample was mixed with 2ml of HCl. Then 6 drops of HCN was added and further 2 drops of picric acid was added that resulted in a creamish pale yellow ppt indicating the presence of alkaloids.

Test for Terpenoids

2 ml of sample along with 2ml of chloroform and 3ml of con. H₂SO₄ was added. Red color ppt obtained indicates the presence of terpenoids.

Test for proteins

One milliliter of ninhydrin was dissolved in 1 mL of acetone and then a small amount of extract was added with ninhydrin. The formation of purple colour revealed the presence of protein.

Detection of saponins

Foam test: A fraction of the extract was vigorously shaken with water and observed for persistent foam.

Test for steroids

One milliliter of chloroform was mixed with 1 mL of extract and then ten drops of acetic anhydride and five drops of concentrated sulphuric acid were added and mixed. The formation of dark red colour or dark pink colour indicates the presence of steroids.

2. DPPH free radical scavenging activity of *Alternanthera sessilis* and *Moringa oleifera*:

Scavenging of 2, 2-Diphenyl-1-picrylhydrazyl (DPPH) radical was assessed by the method of Hatano et al, (1989). The DPPH solution (1.0 ml) was added to 1.0 ml of extract at different concentrations (0.1 to 0.5mg/ml). The mixture was kept at room temperature for 50 minutes and the activity was measured at 517 nm. Ascorbic acid at the same concentrations was used as standard. The capability to scavenge the DPPH radical was calculated and expressed in percentage (%) using following formula:

$$\text{DPPH radical scavenged (\%)} = \frac{\text{Control OD} - \text{Sample OD}}{\text{Control OD}} \times 100$$

3. In Vitro Xanthine Oxidase Inhibitory Activity of *Alternanthera sessilis* and *Moringa oleifera*:

In vitro Xanthine oxidase inhibitory of the extract was assessed as per the method of (Nguyen et al, 2004; Umamaheswari et al., 2007). Briefly, the assay mixture consisted of 1 ml of the fraction (0.1 to 0.5g/ml), 2.9 ml of phosphate buffer (pH 7.5) and 0.1 ml of xanthine oxidase enzyme solution (0.1 units/ml in phosphate buffer, pH 7.5), which was prepared immediately before use. After preincubation at 25°C for 15 min, the reaction was initiated by the addition of 2 ml of the substrate solution (150 M xanthine in the same buffer). The assay mixture was incubated at 25°C for 30 min. The reaction was then stopped by the addition of 1 ml of 1N hydrochloric acid and the absorbance was measured at 290 nm using a UV spectrophotometer. Allopurinol (0.1 to 0.5mg/ml), a known inhibitor of XO, was used as the positive control. One unit of XO is defined as the amount of enzyme required to produce 1 mmol of uric acid/min at 25°C. XO activity was expressed as the percentage inhibition of XO in the above assay system calculated as percentage of inhibition as follows.

Inhibitory activity (%) = $(1 - A_s/A_c) \times 100$ Where,

A_s – absorbance in presence of test substance, A_c – absorbance of control


4. Statistical Analysis

The data were subjected to statistical analysis using one – way analysis of variance (ANOVA) and Duncan's multiple range test to assess the significance of individual variations between the groups. In Duncan's test, significance was considered at the level of $p < 0.05$.


RESULTS AND DISCUSSION :

Table 1 : The table depicts the phytoconstituents of *Alternanthera sessilis* and *Moringa oleifera*

PHYTOCONSTITUENTS	<i>Alternanthera sessilis</i>	<i>Moringa oleifera</i>
Protein	Absent	+
Amino acid	++	+
Flavonoids	++	+
Terpenoids	++	++
Sapponents	+	+
Steroids	+	+
Alkaloids	++	++



Graph 1: The bar graph depicts the In vitro antioxidant activity of *Alternanthera sessilis* and *Moringa oleifera*. The X axis represents the concentration and the Y axis represents the percentage of inhibition. Blue represents the activity of *Moringa oleifera* and orange represents the activity of *Alternanthera sessilis*. Each line represents mean \pm SEM of 3 independent observations. Significance at $p \leq 0.05$.



Graph 2: The bar graph depicts the In vitro xanthine oxidase inhibitory activity of *Alternanthera sessilis* and *Moringa oleifera*. The X axis represents the concentration and the Y axis represents the percentage of inhibition. Blue represents the activity of *Moringa oleifera* and orange represents the activity of *Alternanthera sessilis*. Each line represents mean \pm SEM of 3 independent observations. Significance at $p \leq 0.05$.

The phytochemical screening of the ethanolic extract of *Alternanthera sessilis* and *Moringa oleifera* showed the presence of different phytoconstituents such as Amino acid, Flavonoids, Alkaloids, Terpenoids, Saponins, steroids and proteins. In *Alternanthera sessilis* amino acids, flavonoids, alkaloids, terpenoids are richly present whereas saponins and steroids are fairly present. In *Moringa oleifera* proteins, amino acids, flavonoids, saponins, steroids are fairly present and alkaloids, terpenoids are richly present.

The percentage of radical scavenging activity of plant extract was evaluated at different concentrations. The Ic_{50} value of *Alternanthera sessilis* was found to be $380 \mu\text{g/ml}$ and the Ic_{50} value of *Moringa oleifera* was estimated as $290 \mu\text{g/ml}$, which is significantly higher than *Alternanthera sessilis*. xanthine oxidase inhibitory potential of *Alternanthera sessilis* and *Moringa Oleifera* was analysed. *Moringa oleifera* showed higher xanthine oxidase inhibitory activity ($Ic_{50} = 300 \mu\text{g/ml}$) than *Alternanthera sessilis*.

Medicinal plants contain many bioactive compounds and antioxidant potential. Herbal extracts can be used as alternative medicines for the treatment of gout. Xanthine oxidase inhibitory assay

is considered as standard to study the anti gout potential of medicinal plants. Some plants and their phytochemicals can act as xanthine oxidase inhibitors(34,35). The phytochemical screening of *Alternanthera sessilis* and *Moringa oleifera* also revealed the presence of various phytonutrients.

The previous study revealed the presence of strong antioxidant potential in both *Moringa oleifera* and *Alternanthera sessilis* extract(34). Xanthine oxidase inhibitory activity was observed for the extracts and compared with the standard drug allopurinol. Though the extracts exhibited a lesser anti gout potential, side effects exhibited by the synthetic drugs needs to be considered. The most prevalent side effects of the synthetic drugs are skin allergies, fever, rashes etc. Thus, further research is required to utilize these herbal extracts into drug formulations.

CONCLUSION:

Phytochemical screening of ethanolic extract of *Alternanthera sessilis* and *Moringa oleifera* showed the presence of phytoconstituents. Both the plant extracts exhibited antioxidant activity. Xanthine oxidase inhibitory activity was observed for the extracts and the standard drug allopurinol. It is concluded that the extracts of *Alternanthera sessilis* and *Moringa oleifera* showed antigout activities and thus it can be used as a natural phytochemical in the treatment of gout.

The study highlights the efficacy of " Herbal " which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

REFERENCES :

1. Harborne AJ. Phytochemical Methods A Guide to Modern Techniques of Plant Analysis. Springer Science & Business Media; 1998. 302 p.
2. Siew YY, Zareisedehizadeh S, Seetoh WG, Neo SY, Tan CH, Koh HL. An ethnobotanical survey on the usage of fresh herbs [Internet]. Vol. 81, Planta Medica. 2015. Available from: <http://dx.doi.org/10.1055/s-0035-1556318>
3. Bukar A, Uba A, Oyeyi T. Antimicrobial profile of moringa oleifera lam. Extracts against some food – borne microorganisms [Internet]. Vol. 3, Bayero Journal of Pure and Applied Sciences. 2010. Available from: <http://dx.doi.org/10.4314/bajopas.v3i1.58706>
4. Anwar F, Latif S, Ashraf M, Gilani AH. Moringa oleifera: a food plant with multiple medicinal uses [Internet]. Vol. 21, Phytotherapy Research. 2007. p. 17–25. Available from: <http://dx.doi.org/10.1002/ptr.2023>
5. Moyo S, Bussmann H, Mangwendeza P, Dusara P, Gaolathe T, Mine M, et al. Validation of

- A Point-of-Care Lactate Device For Screening At-Risk Adults Receiving Combination Antiretroviral Therapy In Botswana. *J Antivir Antiretrovir*. 2011 Oct;3(4):45–8.
6. Brisibe EA, Umoren UE, Brisibe F, Magalhães PM, Ferreira JFS, Luthria D, et al. Nutritional characterisation and antioxidant capacity of different tissues of *Artemisia annua* L [Internet]. Vol. 115, *Food Chemistry*. 2009. p. 1240–6. Available from: <http://dx.doi.org/10.1016/j.foodchem.2009.01.033>
 7. Kumbhare MR, Sivakumar T, Udavant PB, Dhake AS, Surana AR. In vitro antioxidant activity, phytochemical screening, cytotoxicity and total phenolic content in extracts of *Caesalpinia pulcherrima* (Caesalpiniaceae) pods. *Pak J Biol Sci*. 2012 Apr 1;15(7):325–32.
 8. Hossain AI, Faisal M, Rahman S, Jahan R, Rahmatullah M. A preliminary evaluation of antihyperglycemic and analgesic activity of *Alternanthera sessilis* aerial parts. *BMC Complement Altern Med*. 2014 May 24;14:169.
 9. Nile SH, Kumar B, Park SW. In vitro evaluation of selected benzimidazole derivatives as an antioxidant and xanthine oxidase inhibitors. *Chem Biol Drug Des*. 2013 Sep;82(3):290–5.
 10. Teh CL, Chew KF, Ling GR. Acute Gout in Hospitalized patients in Sarawak General Hospital. *Med J Malaysia*. 2014 Jun;69(3):126–8.
 11. Roddy E, Doherty M. Gout. *Epidemiology of gout* [Internet]. Vol. 12, *Arthritis Research & Therapy*. 2010. Available from: <http://dx.doi.org/10.1186/ar3199>
 12. Archana A, Ramesh Babu K. Nutrient composition and antioxidant activity of gonads of sea urchin *Stomopneustes variolaris* [Internet]. Vol. 197, *Food Chemistry*. 2016. p. 597–602. Available from: <http://dx.doi.org/10.1016/j.foodchem.2015.11.003>
 13. Kota S, Govada VR, Anantha RK, Verma MK. An Investigation into phytochemical constituents, antioxidant, antibacterial and anti-cataract activity of *Alternanthera sessilis*, a predominant wild leafy vegetable of South India [Internet]. Vol. 10, *Biocatalysis and Agricultural Biotechnology*. 2017. p. 197–203. Available from: <http://dx.doi.org/10.1016/j.bcab.2017.03.008>
 14. Wu F, Zhu J, Li G, Wang J, Veeraraghavan VP, Krishna Mohan S, et al. Biologically synthesized green gold nanoparticles from Siberian ginseng induce growth-inhibitory effect on melanoma cells (B16). *Artif Cells Nanomed Biotechnol*. 2019 Dec;47(1):3297–305.
 15. Chen F, Tang Y, Sun Y, Veeraraghavan VP, Mohan SK, Cui C. 6-shogaol, a active constituents of ginger prevents UVB radiation mediated inflammation and oxidative stress through modulating Nrf2 signaling in human epidermal keratinocytes (HaCaT cells). *J Photochem Photobiol B*. 2019 Aug;197:111518.
 16. Li Z, Veeraraghavan VP, Mohan SK, Bolla SR, Lakshmanan H, Kumaran S, et al. Apoptotic induction and anti-metastatic activity of eugenol encapsulated chitosan nanopolymer on rat glioma C6 cells via alleviating the MMP signaling pathway [Internet]. Vol. 203, *Journal of Photochemistry and Photobiology B: Biology*. 2020. p. 111773.

Available from: <http://dx.doi.org/10.1016/j.jphotobiol.2019.111773>

17. Babu S, Jayaraman S. An update on β -sitosterol: A potential herbal nutraceutical for diabetic management. *Biomed Pharmacother*. 2020 Nov;131:110702.
18. Malaikolundhan H, Mookkan G, Krishnamoorthi G, Matheswaran N, Alsawalha M, Veeraraghavan VP, et al. Anticarcinogenic effect of gold nanoparticles synthesized from *Albizia lebbek* on HCT-116 colon cancer cell lines. *Artif Cells Nanomed Biotechnol*. 2020 Dec;48(1):1206–13.
19. Han X, Jiang X, Guo L, Wang Y, Veeraraghavan VP, Krishna Mohan S, et al. Anticarcinogenic potential of gold nanoparticles synthesized from *Trichosanthes kirilowii* in colon cancer cells through the induction of apoptotic pathway. *Artif Cells Nanomed Biotechnol*. 2019 Dec;47(1):3577–84.
20. Gothai S, Muniandy K, Gnanaraj C, Ibrahim IAA, Shahzad N, Al-Ghamdi SS, et al. Pharmacological insights into antioxidants against colorectal cancer: A detailed review of the possible mechanisms. *Biomed Pharmacother*. 2018 Nov;107:1514–22.
21. Veeraraghavan VP, Hussain S, Balakrishna JP, Dhawale L, Kullappan M, Ambrose JM, et al. A Comprehensive and Critical Review on Ethnopharmacological Importance of Desert Truffles: *Terfezia claveryi*, *Terfezia boudieri*, and *Tirmania nivea* [Internet]. *Food Reviews International*. 2021. p. 1–20. Available from: <http://dx.doi.org/10.1080/87559129.2021.1889581>
22. Sathya S, Ragul V, Veeraraghavan VP, Singh L, Niyas Ahamed MI. An in vitro study on hexavalent chromium [Cr(VI)] remediation using iron oxide nanoparticles based beads. *Environmental Nanotechnology, Monitoring & Management*. 2020 Dec 1;14:100333.
23. Yang Z, Pu M, Dong X, Ji F, Priya Veeraraghavan V, Yang H. Piperine loaded zinc oxide nanocomposite inhibits the PI3K/AKT/mTOR signaling pathway via attenuating the development of gastric carcinoma: In vitro and in vivo studies. *Arabian Journal of Chemistry*. 2020 May 1;13(5):5501–16.
24. Rajendran P, Alzahrani AM, Rengarajan T, Veeraraghavan VP, Krishna Mohan S. Consumption of reused vegetable oil intensifies BRCA1 mutations. *Crit Rev Food Sci Nutr*. 2020 Oct 27;1–8.
25. Barma MD, Muthupandiyani I, Samuel SR, Amaechi BT. Inhibition of *Streptococcus mutans*, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish. *Arch Oral Biol*. 2021 Jun;126:105132.
26. Samuel SR. Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life? *Int J Paediatr Dent*. 2021 Mar;31(2):285–6.
27. Samuel SR, Kuduruthullah S, Khair AMB, Shayeb MA, Elkaseh A, Varma SR. Dental pain, parental SARS-CoV-2 fear and distress on quality of life of 2 to 6 year-old children during COVID-19. *Int J Paediatr Dent*. 2021 May;31(3):436–41.

28. Tang Y, Rajendran P, Veeraraghavan VP, Hussain S, Balakrishna JP, Chinnathambi A, et al. Osteogenic differentiation and mineralization potential of zinc oxide nanoparticles from *Scutellaria baicalensis* on human osteoblast-like MG-63 cells [Internet]. Vol. 119, Materials Science and Engineering: C. 2021. p. 111656. Available from: <http://dx.doi.org/10.1016/j.msec.2020.111656>
29. Yin Z, Yang Y, Guo T, Veeraraghavan VP, Wang X. Potential chemotherapeutic effect of betalain against human non-small cell lung cancer through PI3K/Akt/mTOR signaling pathway. *Environ Toxicol*. 2021 Jun;36(6):1011–20.
30. Veeraraghavan VP, Periadurai ND, Karunakaran T, Hussain S, Surapaneni KM, Jiao X. Green synthesis of silver nanoparticles from aqueous extract of *Scutellaria barbata* and coating on the cotton fabric for antimicrobial applications and wound healing activity in fibroblast cells (L929). *Saudi J Biol Sci*. 2021 Jul;28(7):3633–40.
31. Mickymaray S, Alfaiz FA, Paramasivam A, Veeraraghavan VP, Periadurai ND, Surapaneni KM, et al. Rhaponticin suppresses osteosarcoma through the inhibition of PI3K-Akt-mTOR pathway. *Saudi J Biol Sci*. 2021 Jul;28(7):3641–9.
32. Teja KV, Ramesh S. Is a filled lateral canal – A sign of superiority? [Internet]. Vol. 15, Journal of Dental Sciences. 2020. p. 562–3. Available from: <http://dx.doi.org/10.1016/j.jds.2020.02.009>
33. Theertha M, Sanju S, Priya VV, Jain P, Varma PK, Mony U. Innate lymphoid cells: Potent early mediators of the host immune response during sepsis. *Cell Mol Immunol*. 2020 Oct;17(10):1114–6.
34. Pathak I, Budhathoki R, Yadav N, Niraula M, Kalauni SK. Phytochemical Screening, Cytotoxic and Antioxidant Activity of *Alternanthera sessilis* and *Moringa oleifera* [Internet]. Vol. 1, Amrit Research Journal. 2020. p. 65–71. Available from: <http://dx.doi.org/10.3126/arj.v1i1.32456>
35. Chong S, Loh KE. Xanthine Oxidase Inhibitory Activity of Methanolic Extract of *Alternanthera sessilis* [Internet]. Vol. 49, Sains Malaysiana. 2020. p. 405–10. Available from: <http://dx.doi.org/10.17576/jsm-2020-4902-19>