

Pharmacological activity through synthesizing and evaluating in the form of Gene expression Study

Comment [M1]: Title: revised because it is not specific with the contents of the manuscript. Suggestion: study of the effectiveness of chitosan encapsulated iron oxide nanoparticles

Abstract

Comment [M2]: Abstract: major revision because there is no purpose and conclusion

The magnetic iron oxide nanoparticles have been synthesized by using co-precipitation approach. Chitosan coated nanoparticles compare the effect of surface coating on the stability and toxicity of nanoparticles. Invitro cytotoxicity of the nanoparticles are evaluated by three-(four-5) dimethyl thiazole-2-yl and -two,5, diphenyl thiazoniam bromide (MTT assay) along with glide cytometry examine for cell viability. Does dependent reduction was observed in SKMEL cancer cells administered with different concentration of the sample. I₅₀ value obtained for the Iron nanoparticle is 217.75µg/MI. The fluorescence primarily based real time opposite transcription PCR(RT-PCR) is widely used for the quantification of constant state mRNA ranges and is a crucial device for primary research molecular medicine and biotechnology. FTIR amino companies were acted as capping sites for the iron nanoparticles stabilization. In antibacterial studies Gentamycin 80mcg showed the higher activity of Iron nanoparticles.

Keywords: Chitosan, MTT assay, mRNA, FTIR and Antimicrobial activity.

Comment [M3]: revised according to topic or title

1. Introduction

Chitosan has been widely used for different biological and biomedical applications due to its unique properties. For instance, it can be used in water treatment (1), wound healing materials (2) pharmaceutical excipient or drug carrier (3) obesity treatment (4) and as a scaffold for tissue engineering (5). It is commonly used as an excipient in tablet formulation for oral medication. High molecular weight chitosan is more viscous and delays the release of active ingredient prolongs the duration of drug activity improves therapeutic efficiency as well as reducing the side effects of oral tablets (6). Gene therapy has drawn vast interest over the past many years for the treatment of inherited and acquired diseases. Major studies efforts have focussed on designing suitable provider vectors that

compact and defend oligonucleotides for gene remedy. DNA or RNA is large, hydrophilic macromolecule with negative charge very unstable in physiological environment and do not cross biological membranes effectively since both the gens and cell surfaces are negatively charged spontaneously entry of unprotected genes inside cells is difficult (7). The MTT assay is the most widely used cell viability assay and several modifications of the original method have been described (8). In the present study, an attempt has been made to prepare the Iron oxide nanoparticles through co-precipitation method that has been explain the gene expression study, effective antimicrobial activity and MTT assay.

Comment [M4]: added the novelty of this research

2. Materials and Methods

Comment [M5]: adjusted with suggestion title

2.1 Preparation of chitosan

Crab shells had been accumulated from Rameswaram seashore. The shells were washed with sea water and again with sparkling water and dried. The nicely dried crab shell wastes are powdered by crushing it in a pestle and mortar and sieved. Then subjected to demineralisation, deproteinization deacetylation and eventually the chitosan powder become obtained.

2.2 Preparation of iron oxide nanoparticles

0.1g of Iron (III) chloride hexahydrate $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ and 0.50g of Iron (II) sulphate hexahydrate $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ have been taken in a 250ml beaker. To this delivered 20ml of Millipore water (double distilled). The mixture changed into stirred in a magnetic stirrer for 10 minutes. To this 7% freshly prepared ammonia solution was added to make the mixture more basic. The addition is continued till the PH of reaction mixture reached 11. This mixture was kept for drying in oven at 110° . Finally black precipitate of iron oxide nanoparticles was obtained. It became collected and washed numerous times with water.

2.3 Synthesis of chitosan coated iron oxide nanoparticles

The iron oxide nanoparticles were dried and made proper right into a fine powder. 200mg of chitosan changed into made proper into a gel the use of 2-5 ml of formaldehyde beneath magnetic stirrer for one hour. Finally, a homogenous combination of chitosan coated iron

oxide particles received which have been filtered, dried and saved for addition assessment.

3. Result and Discussion

3.1 FTIR Spectra Analysis

FTIR study was finished to verify the functional group of materials. The spectra of natural Fe nanoparticles indicated that the manifested absorption peaks at approximately 3419, 3049.25, 2360.71, 1592.13, 873 and 553.53cm^{-1} (Fig.1). The surprisingly severe height at 3419 cm^{-1} corresponded to the stretching vibrations of free OH and CH groups (Pawlak et al, 2003). O H vibrations obtained for the peak of 3049 cm^{-1} . The peak at 2360.71 cm^{-1} corresponded to N-H bending vibrations. The peak at 873 cm^{-1} belonged to the C-BR stretch of alkyl halides given in table-1. The end result sincerely indicated the involvement of primary amino companies inside the interaction with metal surface and the amino groups have been acted as capping web sites for the Iron nanoparticles stabilization.




Fig. 1. FTIR spectra of Chitosan encapsulated Iron nanoparticles

3.2 Antibacterial activity

Comment [M6]: Reference sources or citations of chitosan preparations have not been listed.

Comment [M7]: demineralization, deproteination, deacetylation processes have not been specifically described.

The chitosan nanoparticles pretence to have properly bactericidal activity. The synthesized chitosan with nanoparticles has been tested in opposition to two extraordinary pathogenic micro-organisms which include *Escherichia coli*, *Staphylococcus aureus*. From this examine, the chitosan act as an antibacterial agent against *Escherichia coli* and *Staphylococcus aureus*. When the concentration of widespread Gentamycin at 80mcg zone of inhibition for each *S.aureus* and *E.coli* were 25mm. In addition Gentamycin at 400mcg zone of inhibition for *S.aureus* and *E.coli* were identical of 12mm while when gentamycin at 800 mcg zone of inhibition for *S.aureus* is 14 mm and for *E.coli* thirteen (Fig.2). Comparing this region of inhibition for standard Gentamycin 80mcg showed the higher activity than the others.




Fig.2. Antimicrobial activity of chitosan coated Iron oxide nanoparticles

3.3 MTT Assay

MTT Assay used to measure cell metabolic activity as an indicator of cell viability proliferation and cytotoxicity. This colorimetric assay is primarily based the reduction of a yellow tetrazonium salt (three-(four, five-dimethylthiazol-2-yl)-2, five-diphenyl tetrazolium bromide or MTT) to pink formazan crystals by using metabolically active cells. The cytotoxic effect of the chitosan, mobile viability become observed in SKMEL most cancer cells at a wavelength of 570nm administrated with distinctive concentration of 6.25, 12.5, 25, 50, and 100µg/ml. Percentage of viability obtain for chitosan coated iron nanoparticle is 99.39, 96.66, 94.01, 90.02 and 76.81%. The cell viability decreases as per increase in the concentration of chitosan. This showed that the percentage of cancer cells decreased by increases the volume of test solution (Joseph 2012). This experiment was carried out in triplicates and the mean values are depicted in table (1) The IC₅₀ value was obtained for chitosan coated iron nanoparticle 217.75 µg/ml.

Table 1. Viability of Chitosan coated Iron nanoparticle

Concentration(µg/mL)	6.25	12.5	25	50	100
Percentage of viability	99.39	96.66	94.01	90.02	76.81
IC ₅₀	217.75 µg/mL				

The IC₅₀ values were calculated using the equation for slope ($y=mx+C$) obtained by plotting the

absorbance of the different concentrations of the sample (6.25 µg/mL) (Fig 3)




Fig.3.Cytotoxicity detection-MTT assay

3.4 mRNA-agarose electrophoresis

It is mainly used for the purification of DNA fragments. The RNA quality was checked by 1% agarose gel electrophoresis by loading 5µl of the four differ samples. They are HER 1 109, HER2 192, HER1 218 and HER2 96.RNA was obtained was quantitated at 260nm and purity was judged with 260/280 nm. The image of the picture was taken using Medicare, Gelstan geldoc system and is represented in Fig.4




Fig.4 Lane 1: DNA Ladder, lane 2: lane 3: HER2 96, Lane 4: HER1 109, Lane 5: HER2 192, Lane 6: HER1 218

Table 2. mRNA Quantification

Samples	OD at 260	OD at 280	260/280
Control	0.2659	0.1466	1.81
HER296	3.113	1.1935	1.60
HER1 109	2.815	1.52	1.85
HER2 192	2.908	2.525	1.15
HER1 218	3.089	1.625	1.90

Result of this component is RNA sample received as a good integrity and purity. So, before cDNA synthesis attempt to quantify the samples in nanodrop and test the integrity via electrophoresis on a denaturing gel. A right RNA sample will have a 260/230 ratio ranging 2. Zero - 2.2 and 260/280 ratio ranging 1.8 - 2. Zero. All the four samples mRNA purified exact great. They are non-poisonous, in the main used for DNA and RNA separation.

3.5 Real time analysis

Analysis of expression levels of BCL2 and beta-actin in the treated samples with respect to control. The primers used for the real time PCR were BCI2 F- GAACTGGGGGAGGATTGTGG and BCI2 R-ACTTCACTTGTGGCCAGAT. β-Actin selected as a control gene which was detected as the reference gene using the senser primer. They are Beta actin F- ACTATTGGCAACGAGCGGTT and Beta actin R- AGGGTAAAACGCAGCTCA. All gene will amplify around 300bp.

Comment [M8]: Discussion about the results of IC 50 can be added with supporting reference citations

3.6 Average CT values

Quantitative analysis was performed by the measurement of threshold cycle CT value during the exponential phase of amplification. In these studies, HER2 96, HER1 109, HER2 192, and HER1 218 are used as a control. Average CT values for BCl2 are 36.64, 37.61, 38.02, and 35.09. When the CT value is lowering higher in viral loaded. **Lowest CT vales** are 35.09 and **highest** value is 38.02. The obtained CT values for β -actin are 28.16, 32.08, 23.72, and 31.13. Lower the CT value is 23.72 which are lightly affected by viral gene. Higher the CT value is 32.08 which is moderately affected by **viral** gene. These are given in **table: 3**. Comparing this **Bcl2** showed the **good positive results**.

Table 3. Relative Expression of BCl2 with respect to Actin

Samples	BCI2	Actin
CNTL	38.89	30.19
96	36.64	28.16
109	37.61	32.08
192	38.2	23.72
218	35.09	31.13

4. Conclusion

This study shows that chitosan is a promising polymer for biomacromolecules delivery **especially** for gene therapy purposes. Invitro cytotoxicity assays, MTT

assay **used** to measure cell viability. The cell viability decreases as per **increase** in the **chitosan coated** iron nanoparticles. A precise RNA pattern will have a 260/230 ratio ranging 2.0 - 2.2 and 260/280 ratio ranging 1. Eight - 2.0. All **the four** samples **mRNA** properly first-rate. They are non-toxic, mostly used for DNA and RNA separation. When the CT value is lowering higher in viral loaded. Average CT values for BCl2 are 36.64, 37.61, 38.02, and 35.09. These are all higher the value so it is good quality.

5. References

1. E, Onsoyeny, Sksaugrud, O.Metal recovery using chitosan. J.Chem.Technol, 18(1990) 395-404.
2. T, Chandy, Sharma, C. P. Chitosan as a biomaterial, Biomate. Artif. Cells Artif. Organs, 18(1990)1-24.
3. O. Felt, P. Buri Gurny, R. Chitosan; A unique polysaccharide for delivery. Drug Dev. Ind. Pharm, 24,979-993.
4. L.K Han, Y. Kimuru, Okuda, Reduction in fat storage during chitin-treatment in mice fed a high-fat diet. Int. j.obes. Relat. Metab. Disord. 23(1999) 174-179.
5. Y. Zhang, M. Zhan, Three - dimensional microporous calcium phosphate bio-ceramics with nested chitosan sponges for load-bearing bone implants. J. Biomed. Master. Res, 61(2002)1-8.
6. Kofuji, k. Qian, C.J. Nishimura, M; Sugiyama, I, Kawashima, S. Relationship between, physicochemical

Comment [M11]: The number of references is added and ancient references are updated (no. 1,2,3,4,5,8,9), at least the last 5 - 10 years

Comment [M9]: Discussion can be added....

Comment [M12]: Publication year...???

Comment [M10]: conclusion: adjusted for research purposes

characteristics and functional properties of chitosan. Eur. Polym. J, 41(2005)284-2791.

- 7. Murali Ramamoorthi, Aparna Narvekar; Nonviral vectors in Gene therapy, J clin. Diagn Res, 9 (2015), 0973-709X.**
- 8. Edmondson, Armstrong, Martin, Arapid and simple MTT-based spectrometric assay for determining drug sensitivity in monolayer cultures. J Tissue Cult Methods. 11(1998) 15-7.**
- 9. Pawlak and Maria Mucha, Thermogravimetric and FTIR studies of chitosan blends, Thermochemical Acta, 396(2003): 153-166.**
- 10. Manu M Joseph, S.R Aravind, Sheeja Varghese & T.T Sreelekha, Evaluation of antioxidant, antitumor and immunomodulatory properties of polysaccharide isolated from fruit rind of punicagranatum, Molecular medicine reports, 5(2012) 489-96.**

UNDER PEER REVIEW