

Frequency of immediate post-operative gynecologic and obstetric complications in a tertiary care center in Karachi

ABSTRACT

Objective: Postoperative complications are commonly observed during gynecologic and obstetric surgeries that could be controlled by timely management. The factors that involve in these complications could be related with patient or surgeon as well. Therefore, the aim of this study was to determine the frequency of immediately reported postoperative complications in gynecologic and obstetric surgeries in a tertiary care center.

Methodology: A retrospective study was conducted in the Department of Obstetrics and Gynecology of Hamdard hospital, Karachi. The duration of this study was 1 year from July 2018 – June 2019. A total of 200 adult female patients of 18 to 65 years having elective or emergency surgical procedure were included in the study. Data was analysed by using SPSS version-16. Mean standard deviation was calculated for age and hospitalization stay. Frequencies and percentages were calculated for presence of comorbidities such as obesity, anaemia and diabetes mellitus and postoperative complications.

Results: The study results showed the mean age was reported 32.04 ± 7.79 years and mean hospital stay was observed 3.38 ± 1.08 days. Comorbidities revealed that 26(13.0%) were obese, 89(44.5%) were anemic and 6(3.0%) had diabetes. Postoperative Nausea and Vomiting was the most common complication observed in 110(55.0%) women followed by fever that was reported in 69(34.5%) cases. Three patients (1.5%) were found to have Paralytic Ileus, 23(11.5%) reported abdominal distention, postoperative gastritis was reported in 15(7.5%) cases, bleeding per vagina observed in 7(3.5%) women. Wound dehiscence was found in 2(1.0%) cases and Wound Infection in 10(5.0%) women.

Conclusion: The present study concluded that post-operative nausea and vomiting were observed in most of the patients followed by fever subsequent to gynecologic and obstetric surgeries. Additionally, the incidence of wound infection and wound dehiscence were observed low.

Keywords: Nausea, Vomiting, Postoperative complication, Diabetes, Anemia

INTRODUCTION

Risks of complications exist in any type of surgical procedure. The chances of postoperative complications depend on independent distinctive features such as age, coexisting illnesses and functional position.(1) These complications can happen both during operation and after operation. Ureters, bowels, nerves and vessels are in immediate contact to the uterus that makes them liable to injury. Complications that occur intraoperatively can be restored instantly if they are quickly recognized during the surgical treatment. It is predicted that gynecological surgeries are accounted for 75% of iatrogenic injuries to the urinary tract (2). Urinary tract injuries can complicate almost 0.2-1% of gynecological surgeries (2). Hysterectomy, salpingectomy, and cystectomy are the most common surgical procedures executed in gynaecology(3).

On the other hand, postoperative complications can occur at any time subsequent to surgery. These complications can be varied included pyrexia, upper respiratory illnesses, urinary tract infections, urinary retention, surgical site infections (SSIs), and abdominal distension as well. It is predicted that SSIs are accounted for 0.5-15% of the total postoperative complications (4). World Health Organization (WHO) stated that 11.8 out of 100 surgical procedures in lower- and middle socioeconomic countries are affected by the surgical site infections (5). The causes that are responsible for these unwanted complications could be either surgeon-related or patient-related. The patient's age, comorbidities, weight, compliance level, hygienic condition, nourishment, and functional status may demonstrate to be direct or indirect contributory factor. (4).

A center for Disease Control and Prevention (CDC) declared that any infections happening in 30 days following surgical operation is referred to surgical site infection (SSI) (6). The time period of the hospital stay and the frequency of hospital re-admission will increase due to having SSI that will lead to higher mortality and morbidity rate (7). Furthermore, obesity; high body mass index (BMI) during pregnancy (25 and over) have a greater risk for postoperative complications following Cesarean Section (8,9). Additionally, pregnant women with other coexisting illness like anemia, diabetes mellitus and hypertension have higher chances to develop SSI following Cesarean Section.(10)

Internationally, the number of cesarean sections performed annually is rising at a striking rate. As a result, postoperative care of these women needs attention. Cesarean delivery (CD) is one of the most frequent surgical procedures carried out in the United States, is accounted for 32% of all deliveries (11). Almost 1.3 million CDs were carried out in 2014

(11). Wound seroma or hematoma that is expressed in 2-5% of women following CD can cause wound dehiscence leading to wound infection (12,13). Wound dehiscence is described as disconnection of incision and it complicates 2-7% following CD (12,13). Moreover, wound infection shows by discharge, erythema, and abnormal hardening of the incision that complicates 2-7% of cases and usually grows 4 to 7 days following CD (14).

As far as comorbidities are concerned with respect to wound infection that includes diabetes mellitus, blood transfusion, obesity, smoking, age, malnutrition, ineffectiveness of immune system, immunosuppressive treatment and longer interval of preoperative hospitalization stay (15). Postoperative complications particularly associated with C-sections are deficient prenatal care, many pregnancies, history of previous C-section, chorioamnionitis, pre-labor breakage of the fetal membranes, labor dystocia, emergency/labored delivery, and obstetrical service carried out in the training hospitals (16).

Gastrointestinal troubles are one of the common postoperative complications (17). Intestinal gas retention, nausea and abdominal pain, abdominal distension is caused by Ileus that occurs due to decreased peristalsis, intestinal manipulation and immobility of intestine (18,19). Moreover, flatulence, (20) nausea and vomiting (21) leads to mother's disappointment and make longer hospitalization stay (22).

There is an inadequate data available regarding immediate postoperative complications with coexisting diseases in women underwent gynecologic and obstetric surgeries. Hence, the aim of this study was to assess the frequencies of immediate and late gynecological and obstetric complications in a tertiary care hospital in order to overcome these complications.

METHODOLOGY

A retrospective study was conducted in the Department of Obstetrics and Gynecology of Hamdard hospital, Karachi by using non-probability, purposive sampling technique. Ethical approval was taken from the ethical review board of Hamdard University, Karachi. Study was carried out over a period of one year from July 2018 – June 2019. A total of 200 women were selected for this study. Adult female patients of 18 to 65 years having elective or emergency procedure were included in the study whereas minor patients admitted due to other pathology, pathological fractures, other benign diseases, Patients who simultaneously having any other pathology at the time of admission, medical or obstetric complications, inter-operative or immediate major postoperative complica-

tions (for instance the need for blood transfusion for any reason, intolerance to oral fluid) were excluded from the study.

A standardized **Performa** was prepared for data collection. Retrospective Data was obtained from departmental records. Demographic information such as age, hospital stay, religion and socioeconomic status were documented. Type of comorbidities (obesity, anaemia and diabetes mellitus) **and postoperative** complications were also documented.

Data was analysed by using SPSS version-16. Mean standard deviation was calculated for age and hospitalization stay. Frequencies and percentages were calculated for type of comorbidities such as obesity, anaemia and diabetes mellitus and postoperative complication.

RESULTS

A total of 200 women were selected for this study their mean age was reported 32.04 ± 7.79 years and mean hospital stay was observed 3.38 ± 1.08 days, as shown in Table I.

As far as religion is concerned, 182(91.0%) were Muslims, 15(7.5%) women were Hindu and 3(1.5%) women were Christian. Socioeconomic status showed 11(5.5%) belonged to lower class, 117(58.5%) belonged to middle class and 72(36.0%) belonged to upper class, as shown in Table II.

Comorbidities revealed that 26(13.0%) were obese, 89(44.5%) were anemic and 6(3.0%) had diabetes. Postoperative Nausea and Vomiting was the most common complication observed in 110(55.0%) women followed by fever that was reported in 69(34.5%) cases. Three patients (1.5%) were found to have Paralytic Ileus, 23(11.5%) reported abdominal distention, postoperative gastritis was reported in 15(7.5%) cases, bleeding per vagina observed in 7(3.5%) women. As far as the surgical site infection is concerned, wound dehiscence was found in 2(1.0%) cases and Wound Infection in 10(5.0%) women, as shown in Table III.

Table I: Mean Demographic characteristics of patients. (n=200) Frequency of religion and socioeconomic status of patients

Variable		Mean±SD n(%)
Age (years)		32.04±7.79
Hospital Stay (Days)		3.38±1.08
Religion	Islam	182(91.0%)
	Hindu	15(7.5%)
	Christian	3(1.5%)

Socio Economic Status Household income	Lower Class	11(5.5%)
	Middle Class	117(58.5%)
	Upper Class	72(36.0%)

Table II: Frequency of Comorbidities and Postoperative complications.

Variable		n	%
Co-Morbid Obesity	Yes	26	13.0
	No	174	87.0
Co-Morbid Anemia	Yes	89	44.5
	No	111	55.5
Co-Morbid Diabetes	Yes	6	3.0
	No	194	97.0
Fever	Yes	69	34.5
	No	131	65.5
Postoperative Nausea and Vomiting	Yes	110	55.0
	No	90	45.0
Paralytic Ileus	Yes	3	1.5
	No	197	98.5
Abdominal Distension	Yes	23	11.5
	No	177	88.5
Wound Dehiscence	Yes	2	1.0
	No	198	99.0
Wound Infection	Yes	10	5.0
	No	190	95.0
Bleeding Per Vagina	Yes	7	3.5
	No	193	96.5
Post Operative Gastritis	Yes	15	7.5
	No	185	92.5

UNDER PEER REVIEW

DISCUSSION

Postoperative complications after surgical procedure are a forthcoming challenge to the operating surgeon. The achievement in the surgical field with respect to innovation in surgical skill and the simple accessibility of surgery as a treatment preference have been ruined due to the rising frequency of wound dehiscence. Various external and internal factors affects wound healing which is a complicated process. The internal factors represent comorbidities such as hypertension and diabetes whereas external factors constitute wound infection, the general suturing techniques, suturing materials, and postoperative wound care.(23)This study demonstrated the instant postoperative complications that happens after gynecologic and obstetric surgical procedures.

One of the studies reported the average interval of hospitalization stay following gynecologic and obstetric surgery was 10.79 ± 7.91 days because of major complications. (4) Our study was inconsistent with the above reported study and revealed that mean hospital stay was 3.38 ± 1.08 days following surgery due to minor complications that was reflected in our study.

Similarly, one research reported the higher frequency of nausea that was 73.6% after 2 hours following cesarean delivery in spinal anesthesia.(24) This is due to the fact that high level of progesterone causes relaxation of smooth muscle, enhance gastrin secretion, reduction in gastrointestinal motility, and lower esophageal sphincter tones that leads to develop more frequently emetic symptoms..(25) Our study was in accordance with above mentioned study and revealed that postoperative nausea and vomiting was reportedly high in 110(55.0%) women.

Likewise, the frequency of Postoperative nausea and vomiting (PONV) in one research was relatively low (26) contrast to some researches from most parts of the globe (27-30)such as Asia, North America and Europe. This is regardless of the fact that a considerable number of these patients peri-operatively received opioids. The present study was inconsistent with the above reported studies and revealed that incidence of postoperative nausea and vomiting was high 110(55.0%) due to insufficient administration of opioids peri-operatively.

Postoperative wound infection is of huge significance for both patient and surgeon. Another research that was conducted in hospital of Nepal, determined the frequency of postoperative wound infection and to assess the risk factors that leads to wound infection in obstetrics and gynecology surgery. The incidence of wound infection was found in 5.87% cases. 70% patients had preoperative hospital stay that ranges to 0-2 days while 30% had observed 3-18 days (31) The present study was in agreement with the above reported study and showed that only 5.0% had postoperative wound infection and mean hospital stay was 3.38 ± 1.08 days.

Frequency of wound dehiscence following obstetrical and gynecological surgeries can be considered as surgical crisis that can prolong the illness period and causes unnecessary anxiety to patients and can also raise the hospitalization expenses. Therefore, these complications can be prevented by handling the factors included in the progression of wound dehiscence.(32) One research demonstrated the frequency of wound dehiscence in their hospital was observed to be 3.05% and secondary suturing are required for all of these patients. They found the commonest coexisting morbidities were diabetes and hypertension. Among comorbidities, high frequency of wound dehiscence was observed in diabetes and hypertension. One more secondary effect was the usual stay of patients in hospital with wound dehiscence that reflected overall morbidity in these patients. Patients reflected prolong stay in the hospital who developed wound dehiscence.(33) As far as the present study is concerned, coexisting disease such as diabetes was reported only in 6(3.0%) cases and wound dehiscence was observed only in 2(1.0%) cases. Therefore, it was proved by our study that patients had low incidence of wound dehiscence reflecting shorter duration in hospital stay regardless of existence of co-morbidity like diabetes.

Post-operative fever is frequently observed following most of the gynecologic surgeries. Though mostly fever is physiological following surgery with self-resolution while few need thorough investigations.(34) Therefore, one retrospective study assessed the frequency and risk factors for postoperative febrile morbidity in 199 women subsequent to laparoscopic-assisted vaginal hysterectomy (LAVH). Their mean age was reported 46 ± 6 years, a mean overall hospital stay of 5 ± 2 days, and a mean hospital stay of 3 ± 2 days postoperatively. Postoperative fever was recorded in 31(15.6%) cases thereby reflecting moderate frequency of febrile morbidity after LAVH (35). The present study was inconsistent with the above cited research and reported that mean age of studied women was 32.04 ± 7.79 years and mean postoperative hospital stay of 3.38 ± 1.08 days. Postoperative fever was documented in 69 (34.5%) cases thereby reflecting higher frequency of febrile morbidity.

CONCLUSION

The present study concluded that post-operative nausea and vomiting were observed in most of the patients followed by fever subsequent to gynecological and obstetrical surgeries. Even though postoperative fever is common in gynecologic patients, the incidence of wound infection and wound dehiscence were observed low.

REFERENCES

1. Erekson EA, Yip SO, Ciarleglio MM, Fried TR. Postoperative complications after gynecologic surgery. *Obstet Gynecol.* 2011 Oct;118(4):785-93. doi: 10.1097/AOG.0b013e31822dac5d.
2. Pal DK, Wats V, Ghosh B. Urologic complications following obstetrics and gynecological surgery: Our experience in a tertiary care hospital. *Urol Ann.* 2016 Jan-Mar;8(1):26-30. doi: 10.4103/0974-7796.158502.
3. Fuentes MN, Rodríguez-Oliver A, NaveiroRilo JC, Paredes AG, Aguilar Romero MT, Parra JF. Complications of laparoscopic gynecologic surgery. *JLS.* 2014 Jul-Sep;18(3):e2014.00058. doi: 10.4293/JLS.2014.00058.
4. Bahadur A, Mundhra R, Kashibhatla J, Chawla L, Ajmani M, Sharma S, et al. Intraoperative and Postoperative Complications in Gynaecological Surgery: A Retrospective Analysis. *Cureus.* 2021 May 7;13(5):e14885. doi: 10.7759/cureus.14885.
5. World Health Organization 2016: Global guidelines on the prevention of surgical site infection. [Jun;2020];<http://www.who.int/gpsc/ssi-prevention-guidelines/en/2016>
6. Consensus paper on the surveillance of surgical wound infections. *Infect Control HospEpidemiol.* 1992;13:599-605. 10.1086/646435
7. Shikha SS, Latif T, Moshin M, Hossain MA, Akter H, Shamsi S, et al. Evaluation of Surgical Site Infection among Post Cesarean Patients in Mymensingh Medical College Hospital. *Mymensingh Med J.* 2018 Jul;27(3):480-486.
8. Molla M, Temesgen K, Seyoum T, Melkamu M. Surgical site infection and associated factors among women underwent cesarean delivery in Debretabor General Hospital, Northwest Ethiopia: hospital based cross sectional study. *BMC Pregnancy Childbirth.* 2019 Aug 29;19(1):317. doi: 10.1186/s12884-019-2442-0.
9. Gong SP, Guo HX, Zhou HZ, Chen L, Yu YH. Morbidity and risk factors for surgical site infection following cesarean section in Guangdong Province, China. *J*

ObstetGynaecol Res. 2012 Mar;38(3):509-15. doi: 10.1111/j.1447-0756.2011.01746.x.

10. Zejnullahu VA, Isjanovska R, Sejfiija Z, Zejnullahu VA. Surgical site infections after cesarean sections at the University Clinical Center of Kosovo: rates, microbiological profile and risk factors. *BMC Infect Dis.* 2019 Aug 28;19(1):752. doi: 10.1186/s12879-019-4383-7.
11. MacDorman M, Declercq E, Menacker F. Recent trends and patterns in cesarean and vaginal birth after cesarean (VBAC) deliveries in the United States. *ClinPerinatol.* 2011 Jun;38(2):179-92. doi: 10.1016/j.clp.2011.03.007.
12. Mackeen AD, Khalifeh A, Fleisher J, Vogell A, Han C, Sendekci J, et al. Suture compared with staple skin closure after cesarean delivery: a randomized controlled trial. *Obstet Gynecol.* 2014;123(6):1169–1175. doi: 10.1097/AOG.0000000000000227.
13. Peleg D, Eberstark E, Warsof SL, Cohen N, Ben Shachar I. et al. Early wound dressing removal after scheduled cesarean delivery: a randomized controlled trial. *Am J Obstet Gynecol.* 2016;215(3):388.e1–388.e5. doi: 10.1016/j.ajog.2016.03.035.
14. Suarez-Easton S, Zafran N, Garmi G, Salim R. Postcesarean wound infection: prevalence, impact, prevention, and management challenges. *Int J Womens Health.* 2017 Feb 17;9:81-88. doi: 10.2147/IJWH.S98876.
15. deAraújo Madeira MZ, Trabasso P. Surgical site infections in women and their association with clinical conditions. *Rev Soc Bras Med Trop.* 2014 Jul;47(4):457–61.
16. Kawakita T, Landy HJ. Surgical site infections after cesarean delivery: epidemiology, prevention and treatment. *Mater Health NeonatolPerinatol.* 2017;1(3):1–9.
17. Adeli M, Razmjoo N, Tara F, Ebrahimzade S. Effect of early post cesarean feeding on gastrointestinal complications. *Nurs Midwifery Stud.* 2013 Jun;2(2):176-81. doi: 10.5812/nms.10184.
18. Chantarasorn V, Tannirandorn Y. A comparative study of early postoperative feeding versus conventional feeding for patients undergoing cesarean section; a randomized controlled trial. *J Med Assoc Thai.* 2006;89Suppl 4:S11–6.

19. Gocmen A, Gocmen M, Saraoglu M. Early post-operative feeding after caesarean delivery. *J Int Med Res.* 2002;30(5):506–11.
20. Izbizky GH, Minig L, Sebastiani MA, Otano L. The effect of early versus delayed postcaesarean feeding on women's satisfaction: a randomised controlled trial. *BJOG.* 2008;115(3):332–8. doi: 10.1111/j.1471-0528.2007.01591.x.
21. Teoh WH, Shah MK, Mah CL. A randomised controlled trial on beneficial effects of early feeding post-Caesarean delivery under regional anaesthesia. *Singapore Med J.* 2007;48(2):152–7.
22. Bar G, Sheiner E, Lezerovitz A, Lazer T, Hallak M. Early maternal feeding following caesarean delivery: a prospective randomised study. *ActaObstetGynecol Scand.* 2008;87(1):68–71. doi: 10.1080/00016340701778849.
23. Al Jama FE. Risk factors for wound infection after lower segment caesarean section. *Qatar Med J* 2012;12(2):26–31.
24. Jabalameli M, Honarmand A, Safavi M, Chitsaz M. Treatment of postoperative nausea and vomiting after spinal anesthesia for cesarean delivery: A randomized, double-blinded comparison of midazolam, ondansetron, and a combination. *Adv Biomed Res.* 2012;1:2. doi: 10.4103/2277-9175.94424.
25. Klauser CK, Fox NS, Istwan N, Rhea D, Rebarber A, Desch C, et al. Treatment of severe nausea and vomiting of pregnancy with subcutaneous medications. *Am J Perinatol.* 2011 Oct; 28(9):715-21.
26. Ugochukwu O, Adaobi A, Ewah R, Obioma O. Postoperative nausea and vomiting in a gynecological and obstetrical population in South Eastern Nigeria. *Pan Afr Med J.* 2010;7:6. doi: 10.4314/pamj.v7i1.69111.
27. Harmon D, Ryan M, Kelly A, Bowen M. Acupressure and prevention of nausea and vomiting during and after spinal anaesthesia for caesarean section. *Br J Anaesth.* 2000 Apr;84(4):463–7.
28. Van den Bosch JE, Moons KG, Bonsel GJ, Kalkman CJ. Does measurement of preoperative anxiety have added value for predicting postoperative nausea and vomiting? *AnesthAnalg.* 2005 May;100(5):1525-1532. doi: 10.1213/01.ANE.0000149325.20542.D4.
29. Mraovic B, Simurina T, Sonicki Z, Skitarelic N, Gyan TJ. The dose response of nitrous oxide used in postoperative nausea in patients undergoing gynecologic laparoscopic Surgery – A preliminary study. *AnesthAnalg.* 2008 Sep;107(3):818–23.
30. Choi DH, Ko SJ, Ahn HJ, Kim JA. A Korean predictive model for postoperative nausea and vomiting. *J Korean Med Sci.* 2005;20(5):811–815.

31. Suwal A, Shrivastava VR, Giri A. A prospective study of wound infections after laparotomy in obstetrics and gynaecology department. *Nepal Med Coll J.* 2012 Dec;14(4):267-70.
32. Narang R, Nandmer GK, Sapkal R. Factors affecting post operative wound gaping and their outcome in obstetrical and gynecological abdominal surgeries. *Int J ReprodContraceptObstetGynecol* 2017;6(4):1530–1533.
33. Metgud MC, Kataria A, Nadipally SR, Patil K, et al. Incidence of Wound Dehiscence Following Obstetric and Gynecological Surgeries at a Tertiary Care Hospital: A Retrospective Study. *J South Asian FederObstGynae* 2020;12(2):73–78.
34. Pile JC. Evaluating postoperative fever: a focused approach. *Cleve Clin J Med.* 2006 Mar;73Suppl 1:S62-6. doi: 10.3949/ccjm.73.suppl_1.s62.
35. Wongpia I, Thinkhamrop J, Seejorn K, Buppasiri P, Luanratanakorn S, Temtanakitpaisan T, et al. Incidence of and risk factors for febrile morbidity after laparoscopic-assisted vaginal hysterectomy. *Int J Womens Health.* 2014;6:385-388<https://doi.org/10.2147/IJWH.S57521>.