

## Review Form 1.6

|                          |  |
|--------------------------|--|
| Journal Name:            | <a href="#">Journal of Pharmaceutical Research International</a> |
| Manuscript Number:       | Ms_JPRI_86517  |
| Title of the Manuscript: | <b>Biliary Atresia: GGT vs histopathology as diagnostic tool</b> |
| Type of the Article      | <b>Original article</b>  |

### **General guideline for Peer Review process:**

This journal's peer review policy states that **NO** manuscript should be rejected only on the basis of '**lack of Novelty**', provided the manuscript is scientifically robust and technically sound. To know the complete guideline for Peer Review process, reviewers are requested to visit this link:

(<https://www.journaljpri.com/index.php/JPRI/editorial-policy>)

### **PART 1: Review Comments**

|                                     | <b>Reviewer's comment</b>  | <b>Author's comment (if agreed with reviewer, correct the manuscript and highlight that part in the manuscript. It is mandatory that authors should write his/her feedback here)</b>  |
|-------------------------------------|--|---|
| <b>Compulsory</b> REVISION comments | exceptionally consistent with the gold standard i.e. histopathology<br>The current gold standard for diagnosis of biliary atresia is histopathological examination of Liver Biopsy specimen, with reported diagnostic accuracy as high as 94%<br>Introduction is too lengthy .... Make in brief  | Gold standard diagnostic method is Intraoperative cholangiogram   |
| <b>Minor</b> REVISION comments      | In the case of biliary atresia, most infants are full-term, although a higher incidence of low birthweight may be observed<br><br>, whereas GGT levels are almost invariably determined only by liver and bile duct disorders<br><br>, The Children's Hospital & Institute of Child's Health, Lahore in the duration of 6 months (2018-2019) | most infants with BA are full-term, although a higher incidence of low birthweight may be observed<br>\<br>whereas GGT levels are almost invariably ELEVATED only by liver and bile duct disorders<br><br>, The Children's Hospital & Institute of Child's Health, Lahore between (2018-2019) |
| <b>Optional/General</b> comments    | Introduction Should Be Brief And Discussion Should Be Crisp  |   |

### **PART 2:**

|  | <b>Reviewer's comment</b>  | <b>Author's comment (if agreed with reviewer, correct the manuscript and highlight that part in the manuscript. It is mandatory that authors should write his/her feedback here)</b> |
|--|--|--|
| Are there ethical issues in this manuscript? | <u>(If yes, Kindly please write down the ethical issues here in details)</u> |  |

### **Reviewer Details:**

|   |  |
|---|--|
| <b>Name:</b>                                | <b>Raashid Hamid</b>                               |
| <b>Department, University &amp; Country</b> | <b>Kashmir Institute of Medical Science, India</b> |