

**Trypanocidal activity of aqueous and ethanolic extracts of *Leptadenia hastata* on white mice infected with *Trypanosoma congolense***

## ABSTRACT

**Aims:** The aims of this study is to evaluate the trypanocidal activity of the plant extracts on mice infected with the trypanosome strain.

**Study Design:** The retrospective study was conducted from June to November 2020 at the laboratory of animal Biology, Department of Biological Sciences, and University of Ngaoundéré, Cameroon.

**Methodology:** The trypanocidal activity of *Leptadenia hastata* aqueous and ethanolic extracts on *Trypanosoma congolense* was evaluated during ten days of treatment. Thus, the smear was used to monitor the parasitaemia of mice treated with *Leptadenia hastata* aqueous and ethanolic extracts by gavage at doses of 250, 500, 1000 mg/kg (test groups); with 10 ml/kg of distilled water (negative control) and 1mg/kg of isometamidium chloride (positive control).

**Results and Discussion:** The results of the pharmacological studies on the trypanocidal potential of the plant showed that On day ten (D<sub>10</sub>) a mortality rates of 48 %, 68.75 % and 59.18 % respectively for the doses 250, 500 and 1000 mg/kg with the ethanolic extract while the aqueous extract induced a mortality rates of 35 %, 60 % and 41 % respectively. Isometamidium chloride induced the highest mortality rate of 85 %. The lethal doses were 271.37 and 229.07 mg/kg for the aqueous and ethanolic extracts respectively.

**Conclusion:** All these results justify at least in part the use of this plant in traditional medicine for the treatment of trypanosomiasis.

**Keywords:** Ethanolic extracts; Aqueous extracts; Trypanocidal potential; Trypanosomes; Inhibition of parasitaemia.

**Abbreviation:** HAT: Human African Trypanosomiasis, AAT: African Animal Trypanosomiasis, DL<sub>50</sub>: Lethal dose 50, WHO: World Health Organization, ANOVA: Analysis of Variance, T: Morality rates, TN: Negative Control, TP: Positive Control, W: Weight, V: Volume, mg: milligram, kg: kilogram.

## INTRODUCTION

Trypanosomoses are parasitic diseases caused by protozoa belonging to the genus *Trypanosoma* and the family Trypanomastidae, which multiply in blood plasma, lymph and various tissues, including the heart muscle and central nervous system of mammals [1]. Tsetse flies are the pathogens of both HAT and AAT [2]. They therefore play a key role in the epidemiology of trypanosomiasis through their central role in the transmission of trypanosomes to vertebrate hosts [3].

The development of cattle breeding has been consistent with the control of this disease. Despite these available results, trypanocides remain the most widely sold drugs [4]. Currently available treatments (Suramin and pentamidine in the blood stage, melarsoprol and eflornithine in the advanced stage) are outdated, some of them even proving to be toxic to patients [5]. Several approaches to control the vector, such as breeding trypan tolerant animals, using biotopes unfavorable to vector development and using trypanocides are used [6]. Vector control using insecticides of varying degrees of persistence are also used, but these are involved in the destruction of the useful insects such as bees, which play an important role in pollination and honey production. The limitations of these different control methods and the appearance of trypan-resistance have been demonstrated worldwide [7]. The chemo resistance of trypanosomes to the current drugs used since 1940 is becoming an alarming problem, due to the lack of therapeutic alternatives that seriously compromise the control of this parasite [8]. The need for new, less expensive and non-toxic trypanocides has been pressing for almost a century. It is also known that locally used plants are an important source of new drugs [9]. Thus, in view of the poverty afflicting African countries, the evaluation of the trypanocidal activity of medicinal plants remains an important area of research.

*Leptadenia hastata* is a typical forest and fruit species. It is a thorny shrub with a beautiful appearance that reminds one of the jujube tree in terms of its foliage and thorns. *Leptadenia hastata* is a well-known strong medicinal plant. It is presented as an anti-venomous serum,

very good antispasmodic, anti-diarrheal and excellent purgative. It is also used as an anti-malarial, galactogen, against nerve problems and dermatosis

Medicinally, *Leptadenia hastata* has many applications. The latex is applied to wounds and inserted into the nose for headaches. Decoctions and macerations of roots and leaves are applied (alone or in combination with preparations of other plants) against abdominal pain such as constipation, urethral discharge, gonorrhoea, stomachache and diarrhoea. In veterinary medicine, the plant is used against colic in horses and cattle.

This study aims to validate the trypanocidal traditional use of *Leptadenia hastata*, the activity of the aqueous and ethanolic extracts was evaluated on mice infested with *Trypanosoma congolense*.

## 1. MATERIALS AND METHODS

### 1.1. Plant material

The ethnobotanical survey was carried out among the traditional healers, and the recipes and plants most commonly used for the treatment of the disease were selected for the experimental studies in the laboratory. Thus, the leaves of *Leptadenia hastata* were chosen. The leaves of *Leptadenia hastata* were collected during the months of May and June 2020 in Léré (southwestern Chad), a locality located 35 km from Figuil (Cameroon). The identification was carried out at the National Herbarium of Yaoundé/Cameroon where the voucher was kept under specimen number: 40786/SRF. The leaves were washed with water and then dried under artificial ventilation, free from direct sunlight and dust. Once dried, the leaves were grind into powder and sealed in airtight bags.



**Figure 1:** *Leptadenia hastata* [10]

## 1.2. Animal material

The study was carried out on white mice *Mus musculus* Swiss, of both sexes weighing between 20 and 32 g. They were supplied by the National Veterinary Laboratory (LANAVET of Garoua (Cameroon), and then acclimatized for 7 days in the Animal Physiology Laboratory of the University of Ngaoundéré. The animals were fed with pellets supplied by LANAVET and *adlibitum* water.



**Figure 2:** *Mus musculus* Swiss mouse

## 1.3. Trypanosomal strain

The *Trypanosoma congolense* strain used in this study to infect mice are originated from the Faculty of Veterinary Medicine, University of Nigeria Nsukka.

## 1.4. Chemical substance

Isomethamidium chloride (Trypamidium®) which was used as a reference substance in this study was purchased at Veterinary pharmacy in Moundou, Chad. It is in the form of a red powder, which is soluble in water.

## 1.5. Preparation of extracts

One hundred and fifty grams of the crushed material was macerated in 500 ml of distilled water. The maceration was carried out for 24 hours under magnetic stirring, protected from light by covering the beakers with aluminum foil. The macerate was then collected and centrifuged. The supernatant was filtered, frozen and freeze-dried. After 72 hours of

freeze-drying, a powder was obtained with a percentage yield of 24.6%. The lyophilisate was kept in a desiccator to avoid any humidification.

$$\text{Percentage yield} = \frac{\text{Mass of powder extract}}{\text{Mass of concentrated filtrate}} \times 100$$

### **1.5.1. Preparation of ethanolic extracts of *Leptadenia hastata***

One gram of lyophilisate was taken and dissolved into 10 ml of 70 % ethanol. The stock solution of concentration 100 mg/ml was obtained and corresponded to the dose of 1000 mg/kg (D<sub>1</sub>). This solution was dissolved into ½ and ¼ in distilled water and yielded the respective doses of 500 (D<sub>2</sub>) and 250 mg/kg (D<sub>3</sub>).

### **1.5.2. Preparation of aqueous extracts of *Leptadenia hastata***

One gram of dry extract of *Leptadenia hastata* was taken and dissolved into 10 ml of distilled water. The stock solution of concentration 100 mg/ml was obtained and corresponded to the dose of 1000 mg/kg (D<sub>1</sub>). This solution was dissolved into ½ and ¼ in distilled water, and yielded the respective doses 500 (D<sub>2</sub>) and 250 mg/kg (D<sub>3</sub>).

## **1.6. Qualitative analysis of phytochemical constituents**

Three solvents of different polarity were used: Water, ethanol, and diethyl ether. For both extracts, extraction was done by decoction of the plant material at 10 % (W/V: 10/100) in the solvent for 30 minutes. The mixture was filtered and then subjected to various phytochemical tests to determine the content of tannins, saponosides, triterpenes, flavonoids and alkaloids [11]:

### **1.6.1. Determination of tannins**

Two drops of 2 % FeCl<sub>3</sub> solution was added to 2 ml of the test solution and the mixture was left to stand for 5 minutes. A positive test was revealed by the appearance of a blue-black coloration and a precipitate, which confirms the presence of tannins [12].

### **1.6.2. Determination of saponosides**

Five millimeters of three solvents: etheric, ethanolic and aqueous were thoroughly mixed with 10 ml of distilled water for 2 minutes. There was the formation of a persistent foam after 15 minutes, which confirmed the presence of saponosides [12].

### **1.6.3. Determination of triterpenes**

Five millimeters of each extract was mixed with 2 ml of chloroform and 3 ml of concentrated sulphuric acid. A brownish-red color of the interface layer indicated the presence of heterosidic triterpenes [13].

### **1.6.4. Determination of flavonoid**

Five millimeters of *Leptadenia hastata* extracts were mixed with a few drops of concentrated HCl and a quantity of magnesium turnings was added (leaving to act). The presence of aglycone flavones was confirmed by the appearance of a red or orange color [12].

### **1.6.5. Determination of alkaloid**

A few drops of Mayer's reagent were added to 1 ml of *Leptadenia hastata* extracts. The formation of a white precipitate indicated that the test was positive.

## **1.7. Distribution and infestation of mice**

Mice of about 8 to 10 weeks old were selected homogeneously according to their weight (20 to 32 g). These mice marked on their tails were distributed into 14 groups of 6 mice each in polystyrene cages: 1 negative control group, 1 positive control group and 12 test groups. After one week of acclimatization in the laboratory, the mice were inoculated by an intraperitoneal injection of about 0.05 ml of blood containing about 1000 strains of *Trypanosoma congolense* trypanosomes. After injection, the animals were placed in cages with the same environmental conditions as before. Parasitemia was checked on day 4 post-infection, and then the mice were treated on day 5 with the different doses of the extracts (250, 500 and 1000 mg/kg) and isometmidium chloride (1mg/kg). As from the 7th day, a cut was made on the terminal part of the tail of each animal every 2 days during 10 days. After the cutting of the tail, a drop of blood was deposited between slide and coverslip for the observation under the microscope and the number of parasites per field was recorded. The evolution of mice body weight was also evaluated. The pronouncement of clinical symptoms (coat, behavior, noticeable emaciation) and mortality during the infection were recorded.

## **1.8. Treatment of infected mice**

The mice divided as described above were treated with different treatments. The negative control group was treated with distilled water (by oral pathway), the positive control group was receive a dose of 1mg/kg of isometamidium while the 12 test groups were treated orally with different doses (250, 500 and 1000 mg/kg) of the aqueous and ethanolic extracts. After treatment, mice were monitored for 10 days and fresh blood was drawn (50 µl) by cutting off the terminal part of the tail of each mouse every 2 days for parasitaemia analysis.

### **1.9. Body weight and food intake measurements**

Every animal in each group were weighed before blood collection every two days sing an electronic kitchen scale. The animal food intake in each group was also recorded daily. The mice were weighed every other day.

### **1.10. Realization of blood smears**

A drop of blood from the tip of the tail of each mouse was placed on a slide. Another slide was used to spread the drop. A smear was prepared following the usual hematology procedure. The smear was air-dried and then fixed with methanol for one minute. Excess methanol was removed by turning the smear downwards onto a staining tray. Using a 20 ml syringe and a blunt-tipped needle, the Giemsa was diluted 1:10 with buffered distilled water. After mixing, the Giemsa was removed by air. Using the needle and syringe, the Giemsa solution was introduced under the slide, taking care to avoid trapping large air bubbles.

The whole set was left for 30 minutes. At the end of the staining time, the slides are rinsed briefly with running water and then left to dry in an upright position. Any parasites were observed with an immersion objective for further detail of their morphology. Immersion objectives x 100 were particularly useful in the preliminary examination. Several microscopic fields are scanned to determine the presence or absence of trypanosomes.

### **1.11. Statistical analysis:**

Blood parasite densities (D) were determined using the following formula from [14]:

$$D = \frac{\text{Number of parasites}}{\text{Number of fields read}} \times 50$$

The number of parasites corresponds to the number of trypanosomes in 100 slide fields.

Mortality rates (T) are determined by the following formula from [15]:

$$T = \frac{(A-B)}{A} \times 100$$

A and B are the parasite densities before and after treatment respectively, the same animal being considered as a control before treatment. The difference between the positive control and treated groups was statistically analyzed by the analysis of variance (ANOVA) method followed by Dennett's multiple comparison tests, using Graph Pad Instate software. P-values of less than 5% ( $P < 0.05$ ) are considered statistically significant. The lethal doses 50 ( $LD_{50}$ ) were calculated using the equation of the linear regression line expressed as follows  $Y = A x + B$

x is the value of the decimal logarithm of the doses, assuming that for  $LD_{50}$ ,  $Y = 5$ , then  $x = (5 - B) / A$ .

## 2. RESULTS

### 2.1. Phytochemical characterization tests


**Table 1: Phytochemical composition of *Leptadenia hastata***

Families of chemical compounds	Decoction
Alkaloids	+
Saponosides	+
Flavonoids	+
Tannins	-
Triterpenes	-

**Legends:** +: Present; -: Absent

### 2.2. Overall parasite densities before treatment

Observation of parasite densities before treatment ranged from 250 to 800 parasites/ $\mu$ l (Figure 3). We found out that, the 84 mice inoculated with the *Trypanosoma congolense* strain, where all parasitic. More than 70 % of the mice showed a high parasite density [400-650]. Mice with low parasite densities ranged from [250-350] and those with very low parasite densities were less numerous [700-800].




**Figure 3:** Overall parasite densities from 0 to 4 days before treatment with plant extracts and isometamidium chloride.

### 2.3. Effects of *Leptadenia hastata* on parasite numbers

#### 2.3.1. Effect of aqueous extract on parasite density


In each batch, the different treatments resulted in a progressive decrease in parasite densities as the time after treatment increased. During the ten days ( $D_{10}$ ) after treatment, the 250 and 500 mg/kg dose had a statistically similar trypanocidal effect. The 1000 mg/kg dose on day six ( $D_6$ ) right up to day eight ( $D_8$ ), induced a statistically decrease in density which is similar to that of isometamidium chloride (Figure 4). On day ten ( $D_{10}$ ), there was a significant difference between the 1000 mg/kg dose and isometamidium chloride.



**Figure 4:** Effect of aqueous extract of *Leptadenia hastata* on the daily evolution of *Trypanosoma congolense* numbers.

### 2.3.2. Variations in parasite mortality rate in relation to the effect of the aqueous extract

The different results showed that the high dose (1000 mg/kg) has a potential effect compared to the low doses (250 and 500 mg/kg). On day ten ( $D_{10}$ ) of treatment, parasite mortality was induced by 35 % by the 250 mg/kg dose, 60 % by the 500 mg/kg dose and 41 % by the 1000 mg/kg dose of the aqueous extract of *Leptadenia hastata* in mice infested with *Trypanosoma congolense* (Figure 5). On day ten ( $D_{10}$ ) of treatment, an increase in mortality rate was observed depending on the dose used. The 250 and 500 mg/kg doses showed a mortality rate of less than 50 %. On the other hand, the 1000 mg/kg dose and isometamidium chloride (), a positive control, induced a mortality rate of 65 % and 85 % respectively.




**Figure 5:** Mortality rate of the effect of the aqueous extract of *Leptadenia hastata* on the parasite density of *Trypanosoma congolense*

**2.3.3. Lethal Dose 50 (LD<sub>50</sub>) of the aqueous extract of *Leptadenia hastata***

The equation of the linear regression line obtained with the mortality probits at day 10 of treatment is  $y = 0.307x + 2.209$  (Figure 6).


The LD<sub>50</sub> of the aqueous extract of *Leptadenia hastata* is 229.07 mg/kg on day 10.



**Figure 6:** Regression line of *Trypanosoma congolense* mortality versus the decimal logarithm of the doses of the aqueous extract of *Leptadenia hastata*.

### 2.3.4. Effect of ethanolic extract on parasite density

In figure 7, the effect of the *Leptadenia hastata* extract was a function of the treatment days and the doses administered. The decrease was progressive over time. The 1000 mg/kg dose induced a significant decrease compared to the doses (250 and 500 mg/kg) (Figure 7). At day 0 (D<sub>0</sub>) the parasite density was 500 parasites/ $\mu$ l and at day 10 (D<sub>10</sub>) it was reduced to 250 parasites/ $\mu$ l in the 1000 mg/kg treated group. Until day 6, the effect of isometamidium chloride was comparable to that of the 1000 mg/kg dose (Figure 7). The 250 and 500 mg/kg doses showed a statistically similar trypanocidal effect from day 2 until day 10. The 1000 mg/kg dose and isometamidium chloride have a statistically similar effect on day 2 until day 3 and then its effect differs significantly on day 6 until day 10.




**Figure 7:** Effect of ethanolic extract of *Leptadenia hastata* on the daily evolution of *Trypanosoma congolense* numbers

### 2.3.5. Variations in parasite mortality rate by the effect of ethanolic extract

On the tenth day (D<sub>10</sub>) of treatment, all doses induced a mortality rate higher than 50 %. The activity of this extract is explained by its composition: alkaloids, tannins, terpenes and flavonoids that have an antiparasitic activity. Parasite mortality on the tenth day of treatment

was induced by 48 % by the 250 mg/kg dose, 68.75 % by the 500 mg/kg dose and 59.18 % by the 1000 mg/kg dose. Isometamidium chloride, a positive control, induced a mortality of 85 % (Figure 8).




**Figure 8:** Mortality rate of the effect of ethanolic extract of *Leptadenia hastata* on the parasite density of *Trypanosoma congolense*

### 2.3.6. Lethal dose 50 (LD<sub>50</sub>) of ethanolic extract of *Leptadenia hastata*

The equation of the linear regression line obtained with the mortality probits at day 10 of treatment is  $y = 0.322x + 2.193$  (Figure 9).

The LD<sub>50</sub> of the ethanolic extract of *Leptadenia hastata* is 271.37 mg/kg on day 10.



**Figure 9:** Regression line of *Trypanosoma congolense* mortality versus the decimal logarithm of the doses of ethanolic extract of *Leptadenia hastata*.

## DISCUSSION

More than 80 % of the population in developing countries use medicinal plants for their health problems. Despite this extensive use, very little scientific work is undertaken to verify the safety and efficacy of remedies derived from these plants [16]. In this study, ethanolic and aqueous extracts of *Leptadenia hastata* foliage were tested on mice infected with *Trypanosoma congolense* to demonstrate competition between different doses of isomethamidium chloride and the doses of 250 and 500 mg/kg of *Leptadenia hastata*. Between these doses, aqueous and ethanolic extracts of *Leptadenia hastata* showed trypanocidal activity.

The results of the overall parasite densities before treatment of the 84 mice inoculated with the *Trypanosoma congolense* strain showed that more than 70 % of the mice showed high parasite density. White mice are very sensitive hosts to trypanosomes in general and *Trypanosoma congolense* in particular. Several authors have used this parasite in white mice. This is the case of Matovu et al [17] who showed the role of IgM and B-cells antibodies in *Trypanosoma congolense* infected mice and Balé Bayala et al [18].

The effect of the aqueous extract treatments on the parasite density of mice in each batch resulted in a progressive decrease in parasite densities as the time after treatment increased. Ten days after treatment, the 250 and 500 mg/kg doses had a statistically similar

trypanocidal effect. These results are comparable to those obtained by several authors [6, 19, 18] on the competitive effect of aqueous extracts of the plant and testosterone propionate on impuberate-castrated mice. They obtained a similar effect of two concentrations; this can be explained by the proximity of the doses which induce a similar effect (250 and 500 mg/kg) compared to the 1000 mg/kg dose. These authors show that apart from the low toxicity for the plant, the 1000 mg/kg dose is the most effective.

The different results of the variation of the parasite mortality rate in relation to the effect of the aqueous extract showed that the higher dose (1000 mg/kg) has a potential effect compared to the lower doses (250 and 500 mg/kg). On the tenth day of treatment, the increase in mortality rate was relative to the dose used. Thus, the 250 and 500 mg/kg doses had a mortality rate of less than 50 %. In contrast, the 1000 mg/kg dose and the positive control isometamidium chloride induced a mortality of 65 % and 85 % respectively. The lethal dose 50 of the aqueous extract of *Leptadenia hastata* obtained on day 10 was 218.07 mg/kg. This inhibitory action of the different doses of the extract on *Trypanosoma congolense* parasites could be explained by the presence of active substances such as alkaloids, which have antiparasitic activity [20]. This activity could also be explained by the trypanocidal activity of alkaloids and flavonoids. Mamoudou [7] have shown that the accumulation of diminazene rapidly and specifically in the kinetoplast, a parasite organelle containing DNA, suggests selective inhibition of parasite DNA synthesis.

The effect of the ethanolic extract on the parasite density of the *Leptadenia hastata* extract was relative to the days of treatment and the doses administered. The decrease was progressive over time. The 1000 mg/kg dose induced a significant decrease compared to the doses (250 and 500 mg/kg). This effect is comparable to that of the 1000 mg/kg dose revealed by isometamidium chloride. Indeed, whatever the concentration used the parasite densities decrease when the time increases. This decrease would be due to the concentration of the active principle [21]. We can say that this decrease is caused by the dilution, which decreases the effectiveness of the extract. The less active effects of the 250 and 500 mg/kg doses are comparable to those obtained by Vitouley [19] on the study of the trypanocidal potential of aqueous plant extracts for the treatment of trypanosomiasis. He showed that some plants used at a low dose, have a weak action.

Variations in the mortality rate of parasites by the effect of the ethanolic extract on the tenth day ( $D_{10}$ ) of treatment induced a mortality rate higher than 50 %. The activity of this

extract could be explained by its composition: alkaloids, tannins, terpenes and flavonoids that have antiparasitic activity [19]. Parasite mortality on the tenth day of treatment was increasing with increasing doses. Ethanolic extract of *Leptadenia hastata* induced a mortality of 59.18% at a dose of 1000 mg/kg. These results are statistically comparable to the 85% mortality rate induced by the isometamidium chloride positive control. The lethal dose 50 of ethanolic extract of *Leptadenia hastata* was obtained at 229.07 mg/kg on day 10 (D<sub>10</sub>), which could be explained by the presence of active substances such as alkaloids with anti-parasitic activity.

## CONCLUSION

At the end of this study, it was found that extracts of *Leptadenia hastata* showed trypanocidal activity, which confirms the concern of farmers about the decline in animal fertility after consumption of this plant. Thus, we noted mortality rates of 48%, 68.75% and 59.18% respectively for doses of 250; 500 and 1000 mg/kg of the ethanolic extract of *Leptadenia hastata* while the aqueous extract of *Leptadenia hastata* induced mortality rates of 35%, 60% and 41% respectively for the doses 250; 500 and 1000 mg/kg. The lethal doses 50 of ethanolic and aqueous extracts of *Leptadenia hastata* noted were 229.07 and 271.37 mg/kg on *Trypanosoma congolense* respectively.

### Availability of data and materials

All data generated and/or analyzed during this study are included in the article and its additional files.

### COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

### REFERENCES

1. Nimpaye H, Njiokou F, Njine T, Njitchouang GR, Cuny G, Herder S, et al. *Trypanosoma vivax*, *Typanosoma congolense* forest type and *T. simiae* : prevalence in domestic animals of sleeping sickness foci of Cameroon. *Parasite*. 2011; 18 : 171-179.
2. Hu C, Askoy S. Les réponses immunitaires innées régulent l'infection par le parasite trypanosome de la glossine, *Glossina morsitans morsitans*. *Molecular Microbiology*. 2006; 60: 1194-1204. French.
3. Matovu E, Kitibwa A, Picado A, Biéler S, Bessell PR, Ndungu JM. Serological tests for gambiense human African trypanosomiasis detect antibodies in cattle. *Parasites & Vectors*. 2017; 10: 546-552.

4. Barrett MP, Fairlamb AH. The biochemical basis of arsenical-diamidine crossresistance in African trypanosomes. *Parasitol Today*. 2007; 15(4): 136-40.
5. Kennedy PG. "The continuing problem of human African trypanosomiasis (sleeping sickness)." *Ann Neurol*. 2008; 64(2): 116-26.
6. Nikiéma JB. Effects of antiinflammatory Triterpènes isolated from *Leptadenia hastata* latex on keratinocyte proliferation. *Phytother Res*. 2001; 15(2): 131-134.
7. Mamoudou A, Zoli A, Mbahin N, Tanenbe C, Bourdanne A, Clausen PH, et al. Prevalence and incidence of bovine trypanosomosis on the Adamaoua Plateau in Cameroon 10 years after the tsetse eradication campaign. *Vetinary parasitology*. 2006; 142:16-22.
8. Diallo D, Doumbia O, Sanogo F, Mahamoud M. Contribution to the phytochemical study of some medicinal plants of Gourma. Annex N° 21, SSE progress report wild plants. Doctoral thesis in pharmacy from the University of Gourma Mali. 1992; 140p.
9. Boukef MK. Traditional medicine and pharmacopoeia. Plants in traditional Tunisian medicine. 1986; p350.
10. Boullard B. Medicinal plants of the world, realities and beliefs. Editions Estems. 2001; p636.
11. Hennebelle T, Sahpaz S, Bailleul F. Plant polyphenols, sources, uses and potential in the fight against oxidative stress. *Phytotherapie*. 2004; N°1: 3-6.
12. Karumi Y, Onyeyili PA, Ogugbuaja VO. Identification of active principles of *M. balsamina* (Balsam Apple) leaf extract. *J Med Sci*. 2004; 4(3):179-182.
13. Edeoga HO, Okwu DE, Mbaebie BO. Phytochemical constituents of some Nigerian medicinal plants. *African Journal of Biotechnology*. 2005; 4(7): 685-688.
14. WHO. Control and surveillance of African trypanosomiasis. Report of a Who Expert Committee. Who technical Report Series. 1998; n°881.
15. Tamboura HH. Ecological distribution, morphological characteristics and acute toxicity of aqueous extracts of *Holarrhena floribunda* (G. Don) Durand and Schinz, *Leptadenia hastata* (Pers.) Decne and *Cassia sieberiana* (DC) used by veterinary healers in Burkina Faso. *Afr. J. Trad. Complementary Altern. Med*. 2004; 2: 11-22.
16. Hilaly JE, Israili ZH, Lyoussi B. Acute and chronic toxicological studies of *Ajuja iva* in experimental animals. *J. Ethnopharmacol*. 2004; 91: 43-50.

17. Matovu E, Stewart ML, Geiser F, Brun R, Maser P. Mechanisms of arsenical and diamidine uptake and resistance in *Trypanosoma brucei*. *Eukaryot Cell*. 2003; 2(5): 1003-1008.
18. Balé B, Maria T, Rubio P, Moussa Z, Benoit M, Laya S. Anti-androgenic activity of *Leptadenia hastata* (Pers.) Dene: competitive effect of aqueous extracts of the plant and testosterone propionate on impuberate castrated rats. 2010; p229.
19. Vitouley SH. Study of the trypanocidal potential of aqueous extracts of medicinal plants for the treatment of African animal trypanosomoses. 2005; p63-81.
20. Trail JCM, Ieteren GDM, Feron AL. Effect of trypanosome infection, control of parasitemia and control of anernla development on productivity of N'Dama cattle. *Acta Tropica*. 1991; 48: 37-45.
21. Tamboura HH. Ecological distribution, morphological characteristics and acute toxicity of aqueous extracts of *Holarrhena floribunda* (G. Don) Durand & Schinz, *Leptadenia hastata* (Pers.) Decne and *Cassia sieberiana* (DC) used by veterinary healers in Burkina Faso. *Afr. J. Trad. Complementary Altern. Med.* 2004; 2 : 11-22.