


Original Research Article

Investigation of antioxidant potential in *Ocimum basilicum* flower.

GRAPICAL ABSTRACT



ABSTRACT

Aims: The present work is particularly focused on antioxidant properties of flower of *Ocimum basilicum* plant.

Study design: Study is basically designed on Column chromatography of extracts.

Place and Duration of Study: Sample collection and all experimental work was done in Chemistry Department Government College University, Lahore. The study comprises duration of 6 months.

Methodology: The flower of *Ocimum basilicum* were collected, dried and grinded. It was soaked in methanol-water (70:30) in dark bottle for a week. Followed by a scheme (column chromatography). After TLC of extracts, three activities were done. Phosphomolybdate, Ferric thiocyanate (FTC), and Folin-Ciocalteu (FC reagent) for determination of antioxidant capacity, peroxidation, determination of total phenols respectively.

Results: The sample OC2 and crude have maximum absorbance at the concentration of 100µl, 200µl and 300µl. The results show that crude has maximum antioxidant capacity. The phenolic contents are in the increasing order of fraction OC2, OC5, and crude. The maximum phenolic contents are present in crude. Reference has the maximum ability for peroxidation for ferric thiocyanate complex by giving red colour.

Conclusion: Overall it is concluded that *Ocimum basilicum* flower has antioxidant capacity as good as a standard antioxidant. It is recommended in food/medicine as natural herbal product.

Keywords: [*Ocimum basilicum*;flower;antioxidant capacity;Ferric thiocyanate; Folin-Ciocalteu]

1. INTRODUCTION

Ocimum basilicum belongs to family Lamiaceae. This annual herb having more than 150 species of the genus *Ocimum*, grows in several regions of world. The origin of *O. basilicum* commonly known as basil is native to India, Afghanistan, Pakistan, Northern India and Iran, now this plant is cultivated worldwide [1].

Ocimum basilicum called sweet basil, is botanically described as a branched plant that grows between 0.3 and 1.3 m height. It has light green silky leaves which grow in opposite directions and contain many oily glands that store essential oils. The flowers are colored, from white to purple and arranged in a terminal spike. [2]

Ocimum species are not only characterized by an abundance of compounds such as phenolic acids, but also volatile oils. [3]

Foods based on plants are considered, as they possess natural antioxidants like flavonoids polyphenols, vitamin C and vitamin E. The antioxidants have minimized risk of cardiovascular, chronic diseases and certain types of cancer [4].

It has been found that essential oils from *Ocimum basilicum* act as a strong antiviral agent against DNA viruses like herpes simplex viruses, adenoviruses and hepatitis B virus. Besides RNA viruses like coxsackievirus and enterovirus. [5,6,7]

The complex composition of *Ocimum* species extract determines its strong anti-inflammatory, antibacterial and antiviral activity. The most powerful therapeutic species of *Ocimum*, are *Ocimum basilicum*, *Ocimum gratissimum* and *Ocimum sanctum*. [8]

Studies have shown that hydroalcoholic extract of *Ocimum basilicum* has antiosteoporotic effect, bone protection against osteoporosis induced by glucocorticoids. [9]

Reported research work is novel by considering flower of *Ocimum basilicum*. Previous studies were restricted to leaves, stem and other aerial parts of this plant only.

The aim of this study is determination of antioxidant properties particularly, in flower of *Ocimum basilicum*.

2. MATERIAL AND METHODS

EXPERIMENTAL DETAILS

IR spectrum was recorded using KBr disk with a Perkin-Elmer 735B spectrometer. UV spectra were recorded on UV/VIS spectrophotometer 2300 (Shimadzu, Kyoto, Japan). Heidolph, Laborota 400, rotary evaporator was used to evaporate solvents from samples. Silica gel 60 (0.063-0.200 mm) for column chromatography and TLC silica gel 60 F254 aluminum sheets (20 x 20 cm) from Merck. Solvents of analytical grade were purchased from Panreac (Spain). All other chemicals and reagents of analytical grade were from Merck (Germany). Standard deviation (\pm SD) of repeated measurements was calculated using Microsoft Excel 2007.


METHODOLOGY

The flower of *Ocimum basilicum* was collected, dried and grinded. It was soaked in methanol-water (70:30) in dark bottle for a week.

As organic solvents like methanol, ethanol, and hydro-alcoholic mixtures are mostly used for the extraction of flavonoid and phenolic compounds. [10]

To the dried flower extract of *O.basilicum* which was prepared in methanol different solvents were added to check its solubility. The order of solvents was Hexane 100%, hexane: ethyl acetate (1:2), ethyl acetate 100%, ethyl acetate: methanol (1:2). It was found flower extract was soluble in ethyl acetate. So, it was dissolved in ethyl acetate and water. For removal of aqueous contents the solvent extraction was done. The organic layer(OL) and aqueous layers(AQ) were completely separated. The organic layer(OL) was further processed. Ethyl acetate was evaporated from the extract(EE) and 4% sodium carbonate was added. This further gave two layers organic and aqueous, organic layer (ethyl acetate) was placed as neutral fraction(NF) and dilute HCl was added to the aqueous layer of sodium carbonate and pH was maintained at 3. Two layers were obtained, solvent extraction was done again and finally two layers obtained, one was aqueous fraction(AF) and other as acidic fraction(AF) of ethyl acetate. (Scheme 1)

TLC was done for both the fractions. The spot in neutral fraction was more clearly compared to acidic fraction. Mixture of hexane and ethyl acetate was found best for neutral fraction (NF). The neutral fraction was made clearer by adding ethyl acetate, separating and filtering. Slurry of neutral fraction was prepared and column chromatography was done. (Table 1)



Scheme 1 Extraction and processing of flower extract of *Ocimum basilicum*

Table No.1 Different fractions of *Ocimum basilicum* by Column Chromatography

SR.NO	Solvent system		Fraction code	Volume ml
1	n- Hexane	100%	OC-1	500ml
2	Ethyl acetate:n-hexane	5%	OC-2	500ml
3	Ethyl acetate:n-hexane	20%	OC-3	500ml
4	Ethyl acetate:n-hexane	50%	OC-4	500ml
5	Ethyl acetate:n-hexane	80%	OC-5	500ml
6	Chloroform	100%	OC-6	500ml
7	Chloroform :methanol	1%	OC-7	500ml
8	Chloroform:methanol	5%	OC-8	500ml
9	Chloroform:methanol	10%	OC-9	500ml
10	Chloroform:methanol	20%	OC-10	500ml
11	Methanol	100%	OC-11	500ml

TLC was done for different fractions of column. TLC for fraction 1,3,4,5 was run in pure n-hexane. TLC for fraction 2,6,7,8,9,10,11 was run in 1:1 ethyl acetate –n-hexane. Spots were observed and TLC cards were sprayed with ceric sulphate reagent. Fractions showing same spots were mixed together. The fractions 5%,20%,50% and 80% (OC 2, OC 3, OC 4, OC 5) were separated and their TLC was again run in 2% ethyl acetate:n-hexane(2ml:8ml). These fractions were selected for carrying out anti-oxidant activities.

ANTIOXIDANT ACTIVITIES

PHOSPHOMOLYBDATE METHOD

To the 300,200 and 100 μ l of sample including fractions from the column and crude extract 2ml of phospho ammonium molybdate reagent was added and incubated at 95°C for 60 minutes. After cooling absorbance was read at 695nm against blank(phosphomolybdate reagent)[11].

DETERMINATION OF TOTAL PHENOLS BY FOLIN-CIICALTEU (FC REAGENT)

Standard Gallic acid: To the 100 μ l of the standard 50 μ l Folin-Ciocalteu (FC reagent) and 100 μ l of 10%Na₂CO₃ was added and volume was raised up to 2.5ml with methanol. It was kept for 30 minutes and absorbance was read at 765 nm. Different dilutions were done by taking 1ml from standard and making volume up to 2.5ml by adding methanol. Gallic acid calibration curve was drawn. The equation of the curve was $y=2322.1x$

Samples preparation: To different fraction and crude extract of volume 100 μ l, Folin-Ciocalteu (FC reagent) 50 μ l, Na₂CO₃ 100 μ l was added volume was raised up to 2.5 ml with methanol. Stayed for 30 minutes and reading for absorbance was taken at 765 nm. [12]

FERRIC THIOCYANATE (FTC) METHOD

To the 1mg of fractions and extract ethanol was added.1000 μ l was taken and mixed with 1000 μ l of 2.52% linolenic acid prepared in absolute ethanol.1000 μ l of 0.05M phosphate buffer (pH 7) and 1000 μ l of distilled water was added.(Phosphate buffer was prepared by dissolving0.05M disodium hydrogen phosphate di-hydrate in distilled water and pH was maintained by adding sodium hydroxide solution.)The mixture solution with screw cap was placed in dark oven at 40°C for 10 minutes.

The peroxide value was determined by using thiocyanate as a coloring agent. After 10 minutes 100 μ l of solution was taken and 100 μ l of 30 % ammonium thiocyanate with 100 μ l of 0.02 M ferrous sulphate in 3.5%HCl was added and absorbance was read at 532nm. [13]

Table No.2 Antioxidant activity by phosphomolybdate method using 300 μ l sample

S.No	Samples code	Average absorbance at 300 μ l	(+-)Standard deviation
1	OC 2	1.1	0.08544
2	OC3	0.953333	0.066583
3	OC4	0.363667	0.072947
4	OC5	0.915667	0.010504
5	Crude fraction	2.777667	0.018475

Table No.3 Antioxidant activity by phosphomolybdate method using 200 μ l sample

S.No	Samples code	Average absorbance at 200 μ l	(+-)Standard deviation
1	OC 2	1.457667	0.448487
2	OC 3	0.638	0.008
3	OC 4	0.304	0.076236
4	OC 5	0.308	0.020664
5	Crude fraction	3.215	0

Table No.4 Antioxidant activity by phosphomolybdate method using 100 μ l sample

S.No	Samples code	Average absorbance at 100 μ l	(+)-Standard deviation
1	OC 2	1.893333	0.00611
2	OC 3	1.276	0.003
3	OC 4	0.145	0.091263
4	OC 5	0.698333	0.022053
5	Crude fraction	3.029333	0.0

3. RESULTS AND DISCUSSION

The phosphomolybdenum method is used for antioxidant capacity. In this method the reduction of Mo (VI) to Mo (V) by the antioxidant compound takes place. The formation of a green phosphate/Mo (V) complex is at the maximum absorption of 695 nm. [14] The results of phosphomolybdate activity at different concentrations are same. (Table 2-3-4) The sample OC2 and crude have maximum absorbance at the concentration of 100 μ l, 200 μ l and 300 μ l. However, crude has the much higher absorbance compared to OC2 at all three concentrations. The results lead to the conclusion that the raw material has maximum antioxidant capacity. (Fig 1)




Fig 1. Phosphomolybdate method bar graph showing absorbance at 300,200 and 100 μ l

The Folin-Ciocalteu reagent (FC) is used for the colorimetric determination of phenolic and polyphenolic compounds. The total phenolic contents from different fractions were found by plotting standard calibration curve of Gallic acid by the equation $y=2322.1x$. (Fig 2)




Fig 2. Determination of total phenols by standard calibration curve Gallic acid

Chemically, phenols constitute the largest and most important class of antioxidants. Phenolic contents are in the increasing order of fraction OC2, OC5, and crude. The maximum phenolic contents are present in crude. (Table 5)

Table No.5 Total phenols mg/g equivalent of Gallic acid

S.No	Samples code	Total phenols mg/g equivalent of Gallic acid	(+-)Standard deviation
1	OC 2	4.49×10^{-7}	0.068418
2	OC 3	3.04×10^{-7}	0.143962
3	OC 4	4.35×10^{-7}	0.067471
4	OC 5	6.70×10^{-7}	0.069573
5	Crude fraction	8.87×10^{-8}	0.01

Thiocyanate method is used for the determination of peroxidation. The graph was plotted between absorbance and time for each fraction. Tocopherol was used as a reference. **Fig 3** shows (FTC) ferric thiocyanate activity. Different samples shows different absorbance at different intervals of time - zero hour, second hour and twenty fourth hour. It has been found with increase in interval of time there is increase in absorbance. Therefore, the maximum absorbance has been shown by the samples at twenty fourth hour. (Table 6)

Table No.6 Ferric thiocyanate (FTC) method showing absorbance of samples and reference at different hours

S. No	Samples code	absorbance			(+/-) Standard deviation		
		0 hour	2 nd hour	24 th hour	0 hr	2 nd hr	24 th hr
1	OC 2	0.156	0.19	0.201333	0.001	0.001	0.000577
2	OC 3	0.242	0.339333	0.547	0.001	0.004041	0.002646
3	OC 4	0.284333	0.371667	0.540667	0.00404	0.013429	0.015044
4	OC 5	0.270333	0.371	0.536667	0.00251	0.001	0.008083
5	Reference	0.312	0.479333	0.650333	0.001	0.007638	0.012014




Fig 3. Ferric thiocyanate (FTC) bar graph showing absorbance of samples and reference at different hours

Maximum peroxidation shown at twenty fourth hour is by the reference. Among samples peroxidation is in the increasing order of fraction OC3, OC5, and maximum by OC4.

The results of graphs showed that extract of flower have shown significant antioxidant activity and these results are comparable with the activity of tocopherol.

As **reference** has maximum peroxidation. So, reference has been selected for calculating percentage inhibition. Percentage inhibition calculated is 69% .

The results of all activities proof that high content of antioxidant compounds are present in flower of this plant. Flower extract can also be used in future like leaves extracts, for the radiance and resistance of the skin and slowing down of the aging process by maintaining estrogen levels. [15]

4. CONCLUSION

This investigation suggest that the flower extract of *Ocimum basilicum* possess **antioxidants** compounds, phenols, polyphenols and show peroxidation. So, it is concluded *Ocimum basilicum* flower depicts as satisfactory results as leaf extract [16] like an standard antioxidant itself. On basis of antioxidant potential flower extract is **recommended for** inhibition of diseases like obesity and metabolic disorder.

In future essential oils (linalool, estragole, methyl cinnamate, bicyclosesquiphellandrene, eucalyptol, α -bergamotene, eugenol, γ -cadinene and ger-macrene) [17] can be extracted from flower extract like leaves and stem.

DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

- [1] Telci, I; Bayram, E; Yilmaz, G; Avci, B; Variability in essential oil composition of Turkish basils (*Ocimum basilicum* L.) *Biochem. Syst.Ecol.*, (2006) 34: 489-497.
- [2] Sestili, P; Ismail, T; Calcabrini, C; Guescini, M; Catanzaro, E; Turrini, E; Layla, A; Akhtar, S; Fimognari, C; The Potential Effects of *Ocimum Basilicum* on Health: A Review of *Pharmacological and Toxicological Studies. Expert Opin. Drug Metab. Toxicol.*, (2018) 14, 679–692.
- [3] De Araújo Couto, H.G.S; Blank, A.F; de Oliveira e Silva, A.M; de Lima Nogueira, P.C; de Fátima Arrigoni-Blank, M; de Castro Nizio, D.A; de Oliveira Pinto, J.A; Essential Oils of Basil Chemotypes: Major Compounds, Binary Mixtures, and Antioxidant Activity. *Food Chem.*, (2019) 293, 446–454.
- [4] Choi, Y; Jeong, H. S; Lee, J; Antioxidant activity of methanolic extracts from some grains consumed in Korea. *Food Chemistry*, (2007)103: 130_138.
- [5] Sakkas, H; Papadopoulou, C; Antimicrobial Activity of Basil, Oregano, and Thyme Essential Oils. *J. Microbiol. Biotechnol.*, (2017) 27, 429–438.
- [6] Araújo Silva, V; Pereira da Sousa, J; de Luna Freire Pessôa, H; Fernanda Ramos de Freitas, A; Douglas Melo Coutinho, H; Beuttenmuller Nogueira Alves, L; Oliveira Lima, E; *Ocimum Basilicum*: Antibacterial Activity and Association Study with Antibiotics against Bacteria of Clinical Importance. *Pharm. Biol.*, (2016) 54, 863–867.
- [7] Chiang, L.-C; Ng, L.-T; Cheng, P.-W; Chiang, W; Lin, C.-C; Antiviral Activities of Extracts and Selected Pure Constituents of *Ocimum Basilicum*. *Clin. Exp. Pharmacol. Physiol.*, (2005) 32, 811–816.
- [8] Zahran, E; Abdelmohsen, U; Khalil, H; Desoukey, S; Ahmed, M; Kamel, M; Diversity, Phytochemical and Medicinal Potential of the Genus *Ocimum* L. (Lamiaceae). *Phytochem. Rev.*, (2020)19.
- [9]. Hozayen, W.G; El-Desouky, M.A; Soliman, H.A; Ahmed, R.R; Khaliefa, A.K; Antiosteoporotic Effect of *Petroselinum Crispum*, *Ocimum Basilicum* and *Cichorium Intybus* L. in Glucocorticoid-Induced Osteoporosis in Rats. *BMC Complement. Altern. Med.*, (2016) 16, 165.

- [10] Shahrajabian, M.H; Sun, W; Cheng, Q; Chemical Components and Pharmacological Benefits of Basil (*Ocimum Basilicum*): A Review. *Int. J. Food Prop.*, (2020) 23, 1961–1970.
- [11] Huda-Faujan, N; Noriham, A; Norrakiah, A.S; Babji, A.S; Antioxidant activity of plants methanolic extracts containing phenolic compounds. *African Journal of Biotechnology*, (2009) Vol. 8 (3):484-489.
- [12] Shahwar, D;Raza, M.A; Mirza, A.S.M; Abbasi, M.A; Ahmad, U.V; Comparative Study on the Antioxidant and Antimicrobial Activities of Stem-Bark Extract of Litchi chinensis and its Organic Fractions. *Journal- Chemical Society of Pakistan* (2010)32(3):357-362.
- [13]Osawa,T;Namiki,M.A; Novel Type of Antioxidant Isolated from Leaf Wax of Eucalyptus leaves. *Agricultural and Biological Chemistry*, (1981) Vol 45, (3):735–739.
- [14] Prieto, P; Pineda, M; Aguilar, M; Spectrophotometric quantitation of antioxidant capacity through the formation of a Phosphomolybdenum Complex: Specific application to the determination of vitamin E. *Analytical Biochemistry*, (1999)269: 337-341.
- [15] Antonescu, A.I; Miere, F; Fritea, L; Ganea, M; Zdrinca, M; Dobjanschi, L; Antonescu, A; Vicas, S.I; Bodog, F; Sindhu, R.K; Perspectives on the Combined Effects of *Ocimum basilicum* and *Trifolium pratense* Extracts in Terms of Phytochemical Profile and Pharmacological Effects. *Plants*, (2021), 10, 1390. <https://doi.org/10.3390/plants10071390>.
- [16] Patil, D.D;Mhaske, D.K;Wadhawa, G.C;Antibacterial and Antioxidant study of *Ocimum basilicum* Labiatae (sweet basil).*Journal of Advanced Pharmacy Education & Research*,(2011) 2: 104-112.
- [17] Ahmed, A.F; Fatma, A.K.A; Zhenhua, L; Changqin, L; Jinfeng, W; Wenyi, K; Antioxidant activity and total phenolic content of essential oils and extracts of sweet basil (*Ocimum basilicum* L.) plants. *Food Science and Human Wellness* 8, (2019) 299–305.