

Chemical analysis of a traditional herbal decoction use in Sri Lanka for snake bite treatments

Abstract

Snake bite is considered as a major occupational health problem in Sri Lanka. A traditional decoction consists of nine medicinal plants clinically proven its efficacy for snake bites. In the present study, an attempt was done to carry out chemical analysis of the decoction. Chemical analyses were carried out for the decoction in terms of (a) phytochemical screening (b) quantification of total phenols and total flavonoids and (c) *in vitro* antioxidant activities. Present study, revealed the presence of alkaloids, flavonoids, phenols, saponins, steroids and tannins in the decoction. Moreover, total polyphenol and total flavonoid contents of the decoction were 66.03 ± 2.74 mg gallic acid equivalents/g of extract and 18.93 ± 0.90 mg quercetin acid equivalents/g of extract respectively. ORAC value was 3.51 ± 0.23 mg trolox equivalents/g of extract and dose dependent ($R^2 = 0.9788$) DPPH radical scavenging ability was observed. IC_{50} of the decoction for the DPPH assay was 4.58 ± 0.12 μ g/ml. In conclusion, traditional decoction which used to treat snake bites consists of many important phytochemical classes and exhibit potent *in vitro* antioxidant activity.

Key words: antioxidant potential, phytochemicals, snake bites, traditional decoction

1. Introduction

Snake bite is regarded as a serious occupational health issue. Various active young people, especially those involved in farming and plantation work, have died or become disabled as a result of it in many regions of the world, particularly in the Southeast Asian region [1]. Sri Lanka has a lengthy history of dealing with snake bites. History books state that even the king “Dutugemunu” has been bitten by snakes and had been cured. Management of snake bites is mostly done by traditional physicians. These physicians usually have a family history of snake bite management. Methods of snake bite treatment are gifted from one generation to another and those are practiced and protected very seriously by the members of each generation. Each family has an identical set of management methods which are different from the others. Traditional snake bite treatment practitioners in Sri Lanka have accurate antidotes for dangerous bites. Different plant components, such as leaves, fruits, stem bark, tubers, and roots, have been used as antidotes in the form of paste, powder, juice, infusion, decoction, and crude form. Other ingredients, such as goat milk, butter milk, lime juice, and ghee, are sometimes added to these plant parts[2].

Even today, most of the people living in rural areas in Sri Lanka believe traditional treatment for snake envenomation. Snake bites are treated with medicinal plants in indigenous medicine systems. There is a vast repository of plants that have been reported to have anti-snake venom activity[3, 4]. Investigation of therapeutic potential of plants used for snake bites shows the presence of different phytochemicals such as phenols, flavonoids, tannins and saponins, etc[5,6]. Globally, traditional snake bite treatment practitioners are practicing herbal medicine to cure snake bites. Few investigations have been conducted to evaluate the pharmacologically active ingredients against snake bites[7]. Therefore, present study was aimed to analyze a traditional decoction consists of nine medicinal plants (Table 1) which is given by Ayurvedic physicians for snake bites as a promising treatment.

2. Material and methods

2.1. Plant materials

All the plant materials were collected from Western Province, Sri Lanka during August 2021 to September 2021 and authenticated by a Senior Lecturer, Institute of Indigenous Medicine, University of Colombo, Rajagiriya, Sri Lanka. Voucher specimen of each plant material (Table 1) was deposited at Institute of Indigenous Medicine.

2.2. Preparation of the decoction

All the plant parts cleaned, washed and dried at 40 °C in an oven. Then equal amounts (approx. 7 g) were taken from each plant and pulverized into a coarse powder using a blender (Kenwood, model: BL440, made in China). Then 60 g of the herbal mixture was boiled in water (1920 ml) until the final volume reduced up to 240 ml. Finally, decoction was filtered and concentrated under reduced pressure using a rotary evaporator (yield 6.5% w/w).

2.3.Phyto-chemical screening

Phyto-chemical screening was carried out as described by Karunakaran et al. [8] and Dahanayake et al. [9] Phytochemical screening was carried out based on the presence of color, precipitate or interface.

2.3.1. Test for alkaloids

(a) Picric acid test: Picric acid (1 ml) was added to 1 ml of decoction. Yellow crystalline precipitate indicates the presence of alkaloids.

(b) Mayer's test: Few drops of Mayer's reagent were added to 1 ml of decoction. Yellow precipitate indicates the presence of alkaloids.

(c) Tannic acid test: Tannic acid (1 ml) was added to 1 ml of decoction. Yellow crystalline precipitate indicates the presence of alkaloids.

2.3.2. Test for flavonoids

(a) Conc. H_2SO_4 and ammonia solution: Ammonia solution (1 ml) was added to the 2 ml of decoction. Finally, few drops of conc. H_2SO_4 were added to the solution via the test tube. Yellow coloration indicates the presence of flavonoids.

(b) Aluminum chloride test: A few drops of Aluminum chloride were added to 1 ml of decoction. Yellow color indicates the presence of flavonoids.

2.3.3. Test for phenols

(a) Folin reagent test: A few drops of Folin reagent were added to 1 ml of decoction. Blue color indicates the presence of phenols.

(b) Ferric chloride test: A few drops of Ferric chloride were added to 1 ml of decoction. Blue color indicates the presence of phenols.

(c) Lead acetate test: Lead acetate (1 ml from 10% solution) was added to 1 ml of decoction.

Yellow precipitate indicates the presence of phenols.

2.3.4. Test for saponins

(a) Frothing test: Water (5 ml) was added to the decoction (2.5 ml) and shaken vigorously.

Persistence of stable froth indicates the presence of saponins

2.3.5. Test for steroids

(a) Acetic Anhydride (1 ml) was added to decoction (2 ml). Then conc. H_2SO_4 (1 ml) was

added to the mixture. A color change from violate to blue or green indicates the presence

of steroids

2.3.6. Test for tannins

(a) Lead acetate test: Lead acetate (1 ml from 10% solution) was added to 1 ml of decoction.

Yellow precipitate indicates the presence of tannins.

(b) Ferric chloride Test: A few drops of Ferric chloride were added to 1ml of decoction.

Blackish blue color indicates the presence of tannins.

(c) Vanillin test: Few drops of 10% Vanillin in Ethyl alcohol and conc. HCl were added to 2

ml of decoction. Red color indicated the presence of tannins.

2.3.7. Test for Sesquiterpenes

(a) Test for Sesquiterpenes: Decoction (1 ml) was mixed with conc. H_2SO_4 (0.5 ml). A

brown green, red or blue color indicates the presence of sesquiterpenes.

2.4. *In vitro* antioxidant activities

Different concentrations were made out by dissolving water extract in a mixture of water and methanol (1:1 w/w) and subjected for quantification of (a) total phenolic content and scavenging ability of (b) DPPH (1,1 – diphenyl-2-picryl hydrazyl) radical (c) ABTS [2,2-azino-bis (3ethylbenzothiazoline-6-sulfonicacid) diammonium] radical.

2.4.1. Total phenolic content

Total phenolic content was determined as Singleton and co-workers [10] and gallic acid used as the reference compound.

2.4.2. Total flavonoid content

Total phenolic content was determined as Siddhuraju and Becker [11] and quercetin used as the reference compound.

2.4.3. DPPH (1,1 – diphenyl-2-picryl hydrazyl) assay

DPPH assay was performed in 96-well micro-plates according to the method described by Blois [12] using the water extract of the drug. Inhibition activity percentage (I%) was calculated as follows:

$$I\% = 100 \times \frac{(A_0 - A_1)}{A_0}$$

A_0

Where A_0 is the absorbance of the control sample and A_1 is the absorbance of the test extract. Extract concentration providing 50% inhibition (IC_{50}) was calculated from the graph plotting inhibition percentage against extract concentration.

2.4.4. Oxygen radical absorbance capacity (ORAC) assay

The ORAC radical scavenging assay was performed in 96-well microplates according to the method described by Ou and co-workers [13] using the water extract of the drug.

2.5. Statistical analysis

Statistical analysis was performed using statistical software origin pro 8. All data were expressed as Mean \pm SEM.

3. Results and Discussion

Manny farmer communities in Asian and African countries using plant base treatments (eg. decoctions, pates, juices) to minimize the venom effects such as haemorrhage and edema [14,15]. The toxic compounds of the venom can be neutralized by active compound/s in the snake venom via several mechanisms including (a) precipitation or inactivation of proteins [16], inactivation or enzyme inhibition [17,18], antioxidant activity [19] or combination of one or more activities. Present study, revealed the presence of alkaloids, flavonoids, phenols, saponins, steroids and tannins (Table 2) in the decoction. More than one screening tests were performed to detect the phytochemicals in the decoction. Presence of these active compound/s may responsible for one or multiple mechanisms that neutralize the snake venom.

Total polyphenol and total flavonoid contents of the decoction were 66.03 ± 2.74 mg gallic acid equivalents/g of extract and 18.93 ± 0.90 mg quercetin acid equivalents/g of extract respectively.

. Polyphenols in the water extract react with Folin-Ciocalteu reagent to form a blue complex that can be quantified by visible-light spectrophotometry at 765 nm [20]. A blue chromophore is formed by the reaction, which is made up of a phosphotungstic-phosphomolybdenum complex [20, 21]. The alkaline solution and the concentration of phenolic compounds [20] in the extract determine the maximum absorption of the chromophores. One of the widely followed methods for the determination of TFC in plant extracts is the aluminum chloride colorimetric assay, where Al(III) is utilized as a complexing agent. The method is based on the formation of chelates of Al(III)-flavonoids due to their many oxo and hydroxyl groups. Furthermore, flavonoids bind metal ions such as Al(III) with a high affinity, usually at a 1:1 ratio, depending on experimental conditions such as pH [22]. It is well documented that phenols and flavonoids act against snake venom [5, 6, 23, 24]. This may be one of the reasons that the present traditional decoction has promising effects for snake bites. Moreover, plant extracts which have potent antioxidant potential also exhibit anti-venom effects [23,24]. In the present study, *in vitro* antioxidant potential of the traditional decoction was investigated via ORAC and DPPH assays. ORAC value was 3.51 ± 0.23 mg trolox equivalents/g of extract and dose dependent ($R^2 = 0.9788$) DPPH radical scavenging ability was observed (Figure 1) along with a IC_{50} value of 4.58 ± 0.12 μ g/ml. When DPPH radical react with an antioxidant, its purple colour is disappeared and gives pale yellow colour at 517 nm and measures only the hydrophilic antioxidants [25] while the ORAC test measures the splitting ability of the radical chain reaction by antioxidants through monitoring the inhibition of the oxidation of the peroxy radicals [26].

4. Conclusion

Traditional decoction which used to treat snake bites consists of many important phytochemicals and exhibit potent in vitro antioxidant activity.

Disclosure of conflict of interest

The authors declare that they have no conflicts of interest

References

- [1] World Health Organization (WHO). Guidelines for the Prevention and Clinical Management of Snakebite in Africa. Brazzaville. 2010.
- [2] Liyanarachchi, S.K. Sithiyamsahitha Vish Vaidya Chinthamani, Sri Lanka, Part 11, 1957.
- [3] Schioldann E, Mahmood MA, Kyaw MM, Halliday D, Thwin KT, Chit NN, et al. Why snakebite patients in Myanmar seek traditional healers despite availability of biomedical care at hospitals? Community perspectives on reasons. PLoS Negl Trop Dis. 2018;12:e0006299.
- [4] Krah E, de Kruijf J, Ragno L. Integrating Traditional Healers into the Health Care System: Challenges and Opportunities in Rural Northern Ghana. J Community Health. 2018; 43: 157–163.
- [5] Kulatunga WMSSK, Arawwawala LDAM. Phytochemical analysis of an anti-venom traditional herbal preparation for snake-bite. South Asian Res J Nat Prod. 2019; 2(4): 1-6.
- [6] Kulatunga WMSSK, Arawwawala LDAM. Chemical analysis of anti-venom herbal paste use in Sri Lanka. South Asian Res J Nat Prod. 2021; 4(4): 1-7.

- [7] Wood D, Webb C, Demeyer J. Severe snakebites in northern KwaZulu-Natal: Treatment modalities and outcomes. *South African Med J.* 2009; 99: 814–818.
- [8] Karunakaran R, Thabrew MI, Thammitiyagodage GM, Galhena BP, Arawwawala LDAM. The gastroprotective effect of ethyl acetate fraction of hot water extract of *Trichosanthes cucumerina* Linn and its underlying mechanisms. *BMC Complement Altern Med.* 2017;17:312.
- [9] Dahanayake JM, Perera PK, Galappatty PG, Perera HDSM, Arawwawala LDAM. Comparative phytochemical analysis and antioxidant activities of Tamalakyadi decoction with its modified dosage forms. *Evid Based Complement Altern Med.* 2019;Article ID 6037137.
- [10] Singleton VL, Orthofer R, Lamuela-Raventos RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Meth Enzymol.* 1999; 299: 152-178.
- [11] Siddhuraju P, Becker K. Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *J Agric Food Chem.* 2003; 51: 2144–2155.
- [12] Blois MS. Antioxidant determination by use of stable free radical. *Nature.* 1958;181: 1199-1200.
- [13] Ou B, Hampsch-Woodill M, Prior RL. Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe: *J Agric Food Chem.* 2001; 49: 4619–4626.
- [14] Gomes A, Das R, Sarkhel S, Mishra R, Mukherjee S, Bhattacharya S, Gomes A. Herbs and herbal constituents active against snakebite. *Indian J Exp Biol.* 2010; 48: 865–878.

- [15] Castro O, Gutiérrez JM, Barrios M, Castro I, Romero M, Umaña E. Neutralization of the hemorrhagic effect induced by *Bothrops asper* (Serpentes: Viperidae) venom with tropical plant extracts. *Revista de Biología Tropical*. 1999; 47: 605–616.
- [16] Vale LHF, Mendes MM, Hamaguchi A, Rodrigues VM, Homsí-Brandeburgo MI, Soares AM. Neutralization of pharmacological and toxic activities of *Bothrops* snake venoms by *Schizolobium parahyba* (Fabaceae) aqueous extract and its fractions. *Basic Clin Pharmacol Toxicol*. 2008; 103: 104–107.
- [17] Hung YC, Sava V, Hong MY, Huang G. Inhibitory effects on phospholipase A2 and antivenin activity of melanin extracted from *Thea sinensis* Linn. *Life Sci*. 2004; 74: 2037–2047.
- [18] Alam MI, Gomes A. Adjuvant effects and antiserum action potentiation by a (herbal) compound 2-hydroxy-4-methoxy benzoic acid isolated from the root extract of the Indian medicinal plant “sarsaparilla” (*Hemidesmus indicus* R. Br.). *Toxicon*. 1998; 36: 1423–1431.
- [19] Chatterjee I, Chakravarty A, Gomes A. *Daboia russellii* and *Naja kaouthia* venom neutralization by lupeol acetate isolated from the root extract of Indian sarsaparilla *Hemidesmus indicus* R.Br. *J. Ethnopharmacol*. 2006; 106: 38–43.
- [20] Schofield P, Mbugua DM, Pell AN. Analysis of condensed tannins: A review. *Anim. Feed Sci. Tech*. 2001; 91, 21–40.
- [21] Gülçin I, Sat IG, Beydemir S, Elmastas M, Küfrevioğlu ÖI. Comparison of antioxidant activity of clove (*Eugenia caryophyllata* Thunb) buds and lavender (*Lavandula stoechas* L.). *Food Chem*. 2004; 87: 393–400.

[22] Shraim, AM, Ahmed TA, Raham MM, Hijji YM. Determination of total flavonoid content by aluminium chloride assay: A critical evaluation. LWT Food Sci Technol. 2021; 150:111932.

[23] Salama WH, Abdel-Aty AM, Fahmy AS. Rosemary leaves extract: Anti-snake action against Egyptian Cerastes cerastes venom. J Tradit Complement Med. 2018; 8 (4): 465 – 475.

[24] Abdel-Aty AM, Salama WH, Hamed MB, Fahmy AS, Mohamed SH. Phenolic antioxidant capacity of Mango seed kernels: Therapeutic effect against Viper venom. Rev Bras Farmacogn 2018; 28: 594-601.

[25] Marc F, Davin A, Deglene-Benbrahim L, Ferrand C, Baccaunand M et al. Studies of several analytical methods for antioxidant potential evaluation in food. Med Sci. 2004; 20: 458-463.

[26] Munteanu IG, Apetrei C. Analytical methods used in determining antioxidant activity: A review. Int J Mol Sci. 2021; 22: 3380.

Table 1. Plant ingredients of the traditional decoction use for snake bite treatment

Botanical name	Voucher Number	Family	Part of the plant
<i>Terminalia chebula</i> Retz	TC_K-1	Combretaceae	Fruit
<i>Terminalia bellirica</i> (Gaertn.) roxb	TB_K-2	Combretaceae	Fruit
<i>Phyllanthus emblica</i> Linn	PE_K-3	Phyllanthaceae	Fruit
<i>Azadirachta indica</i> A. Juss	AI-K-4	Meliaceae	Bark

<i>Rubia cordifolia</i> Linn	RC_K-5	Rubiaceae	Whole Plant
<i>Acorus calamus</i> Linn	AC_K-6	Araceae	Rhizome
<i>Picrorrhiza kurroa</i> Royle ex. Benth	BK_K-7	Plantaginaceae	Stem
<i>Coscinium fenestratum</i> (Goetgh.) Colebr	CS_K-8	Menispermaceae	Stem
<i>Tinospora cordifolia</i> (Thunb.) Meiers	TC_K-9	Menispermaceae	Stem

Table 2. Phytochemical screening of the traditional decoction use for snake bite treatment

Phytochemicals	Test/s	Results
Alkaloids	Picric acid Test	Positive
	Mayer's Test	Positive
	Tannic acid Test	Negative
Flavonoids	NH ₃ + H ₂ SO ₄	Positive
	1% Aluminum chloride	Positive
Phenols	Folin Reagent Test	Positive
	Ferric chloride Test	Positive
	Lead Acetate Test	Positive
Saponins	Frothing Test	Positive
Steroids	Acetic anhydride	Positive
Tannins	Lead acetate Test	Positive

	0.1% FeCl ₃	Positive
	Vanillin Test	Negative
Sesquiterpenes	Conc.H ₂ SO ₄	Positive




Figure 1. Dose response relationship of DPPH free radical scavenging activity of the water extract

$I (%) = 100 \times (A_0 - A_1) / A_0$, where A_0 is the absorbance of the control sample and A_1 is the absorbance of the test compound. Extract concentration providing 50% inhibition (IC₅₀) was calculated from the graph plotting inhibition percentage against extract concentration.

