

## Evaluating *in vitro* compatibility of pesticides with *Metarhizium anisopliae* (Metchnikoff) Sorokin and its pathogenicity against *Aphis craccivora* Koch.

**Abstract:** An *in vitro* trial was conducted in the Department of Entomology, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, Ratnagiri, Maharashtra; to determine the compatibility of green muscardine fungus, *Metarhizium anisopliae* (Metchnikoff) Sorokin with pesticides and its pathogenicity against *A. craccivora*. Azadirachtin, Buprofezin and Sulphur were the most suitable pesticides as it showed significantly superior growth on potato dextrose agar media over untreated control. However, Emamectin benzoate, Azadirachtin, Profenophos, Indoxacarb, Thiophanate, Paraquat Dichloride and Glyphosate showed significantly higher mean mycelial weight over the untreated control on potato broth medium. At the same time, amongst pesticides used in the experiment; only insecticides (Emamectin benzoate, Azadirachtin and Deltamethrin) reported synergistic increase in conidial count with the untreated control, while fungicides and herbicides reported drastic reduction in conidial count. The pathogenicity of *M. anisopliae* harvested from the pesticide treatments against *A. craccivora* revealed that all the pesticides investigated had reduced its pathogenicity in comparison to untreated control. However Emamectin benzoate, Indoxacarb and Paraquat dichloride showed maximum per cent mortality and can be used in combination with *M. anisopliae*.

**Keywords:** *Metarhizium anisopliae*, pesticides, bioassay, virulence, conidia, *Aphis craccivora*

### Introduction

The genus *Metarhizium* is a parasite of a great number of arthropods, occurring in more than 200 species of insects and acaridae. These fungi are also adapted to live as saprophytes as well as symbionts in the plant rhizosphere (Gang and Leger, 2002), which make the soil their major habitat. Conidial survival can be effected by interaction with pesticides, environmental factor or by bio-pesticide and/or chemical product used to protect plants. The pesticides may antagonize or synergize efficacy and potential insecticidal activity of *M. anisopliae* and may disrupt natural epizootics of this pathogen.

The knowledge of the compatibility between the EPF and pesticides may facilitate the choice of these combinations in Integrated Pest Management (IPM) programs (Singh *et al.* 2014; Johnson *et al.*, 2020). Combined utilization of selective insecticides in association with fungal pathogens can increase the efficiency of control by reduction in the amount of applied insecticides, minimizing environmental contamination hazards and pest resistance (Quintela and McCoy 1998). There are numerous examples where the application of chemical pesticides has enhanced the efficacy of entomopathogens against insect pests (Kaakeh *et al.* 1997; Gardner and Kinard 1998). *Lecanicillium muscarium*, an insect pathogen that is used commercially to control greenhouse pests and is a

candidate species for the control of *B. tabaci*, was suggested to be applied sequentially with imidacloprid. IPM Strategy (Duarte *et al.* 1992) pointed out the importance in considering the antagonistic effect of the pesticides on all developmental phases of entomopathogenic fungi since these products may affect the bio-insecticide potential as well as the occurrence of epizootics. Several studies have contributed information for the **choice of pesticides with more selective action on the entomopathogens, and most of them have been** conducted under laboratory conditions (Silva *et al.* 1993).

Conidial germination is more important than vegetative growth and sporulation on cadavers, because the former corresponds to the first step that triggers an epizootic, and the fungus relies on it to infect the host successfully. Thereby, if a pesticide causes substantial decrease in conidial germination, it may reduce the effectiveness of the entomopathogen toward its target. As mycelial growth develops inside the insect host and the **concentration of agrochemicals, especially those with systemic mode of action, are usually found in low titre in the hemolymph, there is little chance of this developmental stage to be negatively affected.** Keeping this in view, a laboratory study was conducted to assess the compatibility between the different chemical pesticides with *M. anisopliae*.

#### **Materials and methods**

*M. anisopliae* was isolated from previously maintained lab culture on Agar plates and multiplied on Potato Dextrose Agar (PDA) media **in petriplates as well as Potato Dextrose Broth media in conical flasks at Post Graduate Laboratory, Department of Agricultural Entomology, Dapoli, Ratnagiri, Maharashtra, India. All *In vitro* studies were carried out aseptically in a laminar air flow chamber.** The experiments were conducted under well-defined conditions of culture room maintained at  $25 \pm 2^{\circ}\text{C}$  temperature, uniform light (1600 Lux) provided by fluorescent tubes (7200 K) over a light and dark cycle of 16/8 hours. The insecticides, fungicides and herbicides chosen for the compatibility study are the most frequently used by vegetable growers. The concentrations of all agrochemicals were tested following the label rates recommended by the IRAC. When the concentration of a product indicated minimum and maximum doses, the average concentration was used. The concentration tested for each product was calculated based on the spray volume rate of  $500\text{-}1000 \text{ Lha}^{-1}$  of water.

#### ***Effect of pesticides on radial growth of M. anisopliae***

The effect of all pesticide preparations was tested on growth of *M. anisopliae* by poisoned food technique. With the help of sterile cork borer, discs of 3mm diameter were cut from actively growing culture of the fungus and transferred in the center of petridishes containing desired pesticides together with the medium. The inoculated dishes were incubated and observations on radial growth of colony (cm) were recorded on 4<sup>th</sup>, 6<sup>th</sup>, 8<sup>th</sup>, 10<sup>th</sup> and 12<sup>th</sup> day **after inoculation on PDA medium.**

#### ***Effect of pesticides on biomass of M. anisopliae (mycelial dry weight)***

The effect of pesticides was assessed on biomass of mycelial growth. Each insecticide concentration was mixed with 25 ml of Potato Dextrose Broth (PDB) media in conical flasks of 100 ml capacity. Circular bits of diameter 3mm were made by sterilized cork borer and the conical flasks

were inoculated under laminar air flow. Each treatment (mixture of fungal conidia + pesticide) was replicated three times (three flasks per treatment). After the period of incubation, the mycelial growth was determined when control expressed mycelial mat covering complete surface of substrate. Afterwards, the entire mycelial mat was harvested on properly weighed handmade filter paper by filtration from the PDB medium and fresh weight of mycelial mat was recorded which was later kept in hot air oven at 55-60°C until constant decrease in weight was observed .

#### ***Effect of pesticides on conidial germination of *M. anisopliae****

The appropriate concentrations of each pesticide were added to 25 ml of cooled (45°C) PDB. This treated was directly inoculated with 1 ml conidial suspension of *M. anisopliae* containing  $10^6$  spore  $\text{ml}^{-1}$  that diluted in sterile distilled water amended with 0.01% Tween 80. The same aliquot of sterile distilled water, standard spore suspension and 0.01% Tween 80, without the pesticides, served as control. The same were incubated for 8 days. After incubation by haemo-cytometer, conidia were counted. The data was used to calculate per cent (%) germinated or non-germinated spores.

#### ***Conidia production assessment***

The sterile PDA medium cooled to  $40 \pm 5^\circ\text{C}$  and pre-established concentrations of insecticides were added with the antibiotic streptomycin ( $0.5 \text{ l}^{-1}$ ). Approximately 20 ml of each concentration was poured into 9 cm petridish. Medium without insecticide but containing streptomycin served as control. After solidification, the medium was inoculated with *M. anisopliae*.

The dishes were incubated for 8 days. After 8 days, colony discs were selected randomly for conidial production assessment. Each disc was placed in a screw cap tube containing 10 ml of sterile water with (0.02%) Tween 20. The tubes were agitated until the conidia were completely released and the conidial concentration was counted using Neubauer Chamber.

#### ***Effect of pesticides on pathogenicity of compatible *M. anisopliae* against Aphid, *A. craccivora* under laboratory conditions***

Aphids were reared on brinjal plants grown in plastic pots. Twenty aphids of uniform growth and of same age were selected for experiment. The aphids were placed on a leaf disc in a Petri dish and sprayed with samples of suspension. Suspensions were prepared in distilled water by using 5g of aliquot per litre of water. The aphids were observed for mortality at an interval of 24 hrs. The data thus obtained on mortality was converted into per cent mortality. One control was maintained without any spray.

#### **Data analysis**

The data sets for each experiment were subjected to a two way analysis of variance (ANOVA) in Randomized Block Design and treatment means were compared by Fisher's least significant difference (LSD,  $\alpha = 0.01$ ) for statistical analysis.

#### **Results and Discussion**

##### **Effect of insecticides on *M. anisopliae***

The *in vitro* effect of insecticides on mean radial growth, mean mycelia weight, mean conidial germination and pathogenicity of *M. anisopliae* culture compatible with insecticides was recorded (Table 2). The pathogenicity test was conducted against cowpea aphid, *A. craccivora* *in vitro* and percent mortality was recorded.

### Mean radial growth

#### 4<sup>th</sup> day after inoculation

The results in the present investigations showed that there were significant differences among various treatments. The mean radial growth on 4<sup>th</sup> day after inoculation varied from 0.40cm (T<sub>6</sub>-Deltamethrin) to 6.36cm (T<sub>4</sub>-Buprofezin). The latter was at par with the treatments T<sub>8</sub>- Azadirachtin (6.16cm) and T<sub>10</sub>- untreated control (5.56cm) and significantly superior over rest of the treatments; also supported good growth of the EPF.

#### 6<sup>th</sup> day after inoculation

The mean radial growth on 6<sup>th</sup> day after inoculation varied from 0.56cm (T<sub>6</sub>-Deltamethrin) to 6.86cm (T<sub>4</sub>-Buprofezin). The latter was at par with the treatments T<sub>8</sub>- Azadirachtin (7.02cm) and T<sub>10</sub>- untreated control (5.9cm) and significantly superior over rest of the treatments.

#### 8<sup>th</sup> day after inoculation

The mean radial growth on 8<sup>th</sup> day after inoculation varied from 1.26cm (T<sub>6</sub>-Deltamethrin) to 9.0cm (T<sub>4</sub>-Buprofezin). The latter was at par with the treatments T<sub>8</sub>- Azadirachtin (7.02cm) and T<sub>10</sub>- untreated control (5.9cm) and significantly superior over rest of the treatments. The results of data on 10<sup>th</sup> and 12<sup>th</sup> day after inoculation were similar to the readings after 8 days of inoculation.

In general, the results indicated that amongst various insecticides tested, Buprofezin was found to be the most suitable for *M. anisopliae* growth followed by Azadirachtin. Mycelial growth was reduced in all insecticide treatments when compared to the untreated control except Buprofezin which allowed normal fungal growth. Lowest *M. anisopliae* mycelial growth was observed with Deltamethrin indicating high fungal toxicity. It is possible that those products compatible with *M. anisopliae*, especially the insecticides, might be used at sublethal dosages in combination with the fungus in further studies aimed at improving its virulence. Pachamuthu and Kamble (2000) also observed a significant interaction similar to present findings, between the EPF (*M. anisopliae*) and commercial pesticide (chlorpyrifos) at LT<sub>50</sub> values. The present findings on mycelial growth at the recommended dose of insecticides were contradictory with those of Puzari *et al.* (2006) who observed 100 per cent of mycelial growth when the entomopathogen was grown in presence of Dichlorvos at recommended dose. Cypermethrin and deltamethrin at recommended doses showed lowest growth inhibition against *M. anisopliae*, may be due to species or strain variation; although the biological activity of *M. anisopliae* can be enhanced by integrating it with sub lethal doses of chemical insecticides (Zurek *et al.*, 2002). Schumacher and Poehling (2012) also mentioned similar findings that the insecticides fipronil, permethrin, imidacloprid, neemazal, and amitraz at different concentrations (0.32 to 200 ppm) were found compatible against two strains of the entomopathogenic

fungus *M. anisopliae* (MA) while only fipronil at higher dose rates of 40 and 200 ppm as moderately toxic. The growth rate of *M. anisopliae* in presence of spinosad at low concentrations remains unaffected, but reduces at concentrations of 192 ppm or higher (Ericsson *et al.*, 2007). The insecticide cypermethrin is slightly toxic to all the isolates of *M. anisopliae* and fungicides, carbendazim and hexaconazole are not compatible with *M. anisopliae* isolates (Johnson *et al.*, 2020).

#### **Mean mycelial weight of *M. anisopliae***

Mycelial growth was reduced in all insecticide treatments when compared to the untreated control except T<sub>8</sub> (Azadirachtin 0.15EC), T<sub>2</sub> (Indoxacarb 14.5SC) and T<sub>5</sub> (Profenophos 50EC) which allowed normal fungal growth of 6.211g, 3.787g and 3.784g respectively. White coloured unsporulated mycelial growth was observed with T<sub>8</sub> (Azadirachtin 0.15EC) which was different from the other treatments. The treatment T<sub>7</sub> (Dichlorvos 76EC) strongly inhibited the mycelial growth and recorded lowest mycelial weight 2.386g while other treatments *viz.*, T<sub>4</sub> (Buprofezin 25EC), T<sub>1</sub> (Cypermethrin 25EC), T<sub>3</sub> (Chlorantraniliprole 18.5SC) and T<sub>6</sub> (Deltamethrin 2.8EC) reported 3.484g, 3.292g, 3.187g and 3.105g of mycelial mat respectively. At the same time, control reported 3.605g weight of mycelial mat.

The present findings are similar with the findings of Singh *et al.* (2014) who reported that insecticides are compatible with EPF, though it shows noticeable reduction in biomass on broth media, they can be applied in combination with EPF at half of mean concentration on field evaluation. Babu *et al.* (2013) also found that *M. anisopliae* displays an enhancement in the vegetative growth with imidacloprid (2%) and HIT (2-18%). Among botanicals, Azadirachtin is reported to enhance the vegetative growth and spore germination of *B. bassiana* and *M. anisopliae* (Khan *et al.*, 2012). Rashid *et al.* (2010) observed that the insecticides *viz.*, Fipronil at concentration 10, 40 and 120 ppm, Pyriproxyfen at concentration 10, 500 and 1000 ppm and Hexaflumuron at concentration 20, 50 and 120 ppm on SDA medium; Hexaflumuron at the concentration of 120 ppm reduce the vegetative growth of *M. anisopliae* to zero per cent compared to Pyriproxyfen (24.59%) and Fipronil (24.31%).

#### **Mean conidial germination of *M. anisopliae***

All insecticides affected conidial germination in contrast to the untreated control. The highest (8.00) detrimental effect on conidial germination was observed in T<sub>3</sub> (Chlorantraniliprole 18.5SC). While other treatments *viz.*, T<sub>7</sub> (Azadirachtin 0.15EC), T<sub>4</sub> (Buprofezin 25EC), T<sub>6</sub> (Dichlorvos 76EC), T<sub>2</sub> (Indoxacarb 14.5SC) and T<sub>1</sub> (Cypermethrin 25EC) reported mean conidial germination 39.66, 29.33, 20.00, 19.00 and 13.00 respectively. The treatment T<sub>9</sub> (Emamectin benzoate 5SG) did not affect conidial germination and reported a maximum of mean conidial germination 92.00 while T<sub>5</sub> (Deltamethrin 2.8EC) with mean conidial germination 49.33 was at par with T<sub>9</sub> (Emamectin benzoate). At the same time untreated control reported 33.79 mean conidial germination.

The similar findings were also reported by James and Elzen (2001) that insecticides slightly inhibit the conidial germination and the toxicity is dose dependent i.e. it increases with increase in concentration of insecticides. The use of insecticides, in most cases, had no negative effect on conidia

germination, conidia production and vegetative growth of *B. bassiana*, *M. anisopliae* and *Paecilomyces* sp (Neves *et al.*, 2001). The application of the EPF *M. anisopliae* var. *acridum* and *M. anisopliae* var. *anisopliae* with the Neem oil on PDA medium still liquid ( $\pm 45^{\circ}\text{C}$ ), inhibited conidial production (Santos *et al.*, 2009). Imidacloprid 17.8SL and neem oil 3 percent were found significantly compatible and safe to conidial germination and growth of \_\_\_\_\_??. (Hemalatha *et al.*, 2015). No reports are available in literature regarding the synergistic effect recorded by Emamectin benzoate and Deltamethrin.

#### **Percent mortality of cow pea aphid (*A. craccivora*) by *M. anisopliae* suspension from different treatments**

The highest (80.00%) mortality was recorded in the treatment T<sub>2</sub> (Indoxacarb 14.5SC) and was significantly superior over other treatments. While other treatments viz., T<sub>7</sub> (Emamectin benzoate 5SG), T<sub>6</sub> (Azadirachtin 0.15EC), T<sub>4</sub> (Buprofezin 25EC), T<sub>3</sub> (Chlorantraniliprole 18.5SC) and T<sub>1</sub> (Cypermethrin 25EC) reported 76.65 percent, 66.65 percent, 58.30 percent, 51.65 percent and 40.00 percent mortality, respectively. The lowest (28.30%) per cent mortality was recorded in T<sub>5</sub> (Deltamethrin 2.8EC). At the same time, untreated control recorded 86.65 per cent mortality.

The present findings are in agreement with the findings of Araujo *et al.* (2009) who reported that neem based insecticides expressed 90 per cent mortality in comparison with untreated control against aphid *Lipaphis erysimi* (Kalt.). The synergism exists between the botanical insecticide, azadirachtin and destruxin, a mycotoxin extracted from *M. anisopliae* in a joint action against cotton aphid, *Aphis gossypii* (Fei Yi *et al.*, 2012). The mortality of *B. tabaci* treated with combination of *B. bassiana* and 3 EFAM fungicides is lower than or similar to the mortality with application of *B. bassiana* alone (Ling *et al.*, 2016). The fungus-insecticide bait formulation (termiticide fipronil with a biocontrol agent *M. anisopliae*) showed the greatest compatibility and synergistic effect that increased subterranean termite (*C. curvignathus*) mortality as well as reduced the lethal time at a sublethal dose of fipronil (Jing *et al.*, 2016). The insecticide imidacloprid showed the best compatibility (95.2%) with a positive toxicity/virulence to the *Metarhizium* spp. (strain PDRL526); causing adult cotton mealy bug mortality after (LT<sub>50</sub>) 13.8 and 19.6 days by using  $6.3 \times 10^{12}$  spores/acre, under screen house and field conditions, respectively (Ujjain *et al.*, 2015).

#### **Effect of fungicides on *M. anisopliae***

Similar to the insecticides, the *in vitro* effect of fungicides on *M. anisopliae* was recorded (Table 3).

#### **Mean radial growth**

##### **4<sup>th</sup> day after inoculation**

The result showed that there were significant differences among various treatments. The highest (7.33cm) diameter was recorded in the treatment T<sub>10</sub> (untreated control). The second best treatment was T<sub>4</sub> (Sulphur 80WDG) with diameter 7.16cm and was significantly superior over all other treatments. While other treatments viz., T<sub>9</sub> (Metalaxyl 35WS) and T<sub>5</sub> (Thiophanate methyl 70WP) recorded mean diameter 2.53cm and 1.2cm respectively. On the contrary, the lowest or no

growth was recorded in treatments viz., T<sub>1</sub> (Carbendazim 50WP), T<sub>2</sub> (Mancozeb 75WP), T<sub>3</sub> (Copper oxychloride 50WP), T<sub>6</sub> (Hexaconazole 5EC), T<sub>7</sub> (Difenconazole 25EC) and T<sub>8</sub> (Captan 50WP) showing complete inhibition of germination and multiplication of *M. anisopliae*.

#### **6<sup>th</sup> day after inoculation**

The highest (9.0cm) diameter was recorded in the treatment T<sub>4</sub> (Sulphur 80WP) and was significantly superior over all other treatments. The second best treatment was T<sub>10</sub> (untreated control) with diameter 7.5cm and was significantly superior over all other treatments. While other treatments viz., T<sub>9</sub> (Metalaxyl 35WS) and T<sub>5</sub> (Thiophanate methyl 70WP) recorded mean diameter 3.3cm and 1.26cm respectively. On the same contrary the lowest or no growth was recorded in treatments viz., T<sub>1</sub> (Carbendazim 50WP), T<sub>2</sub> (Mancozeb 75WP), T<sub>3</sub> (Copper oxychloride 50WP), T<sub>6</sub> (Hexaconazole 5EC), T<sub>7</sub> (Difenconazole 25EC) and T<sub>8</sub> (Captan 50WP) showing complete inhibition of germination and multiplication of *M. anisopliae*. The results of data on 8, 10 and 12 days after inoculation were similar to the readings after 6 days of inoculation except treatment T<sub>9</sub> (Metalaxyl 35WS) which showed slight increase in radial growth of fungi up to 4.23cm.

In general, the results indicated that the fungicides were the most harmful products to all biological stages of *M. anisopliae* and they should not be applied together with this EPF. All fungicides negatively affected *M. anisopliae* growth compared to the controls except T<sub>4</sub> (Sulphur 80WP). The present findings were concurrent with Patil *et al.* (2014) and Kotwal *et al.* (2012) who reported that carbendazim and mancozeb resulted in complete growth inhibition (100%) of EPF. Sulphur displayed high compatibility to all isolates at higher concentration except for B57 and Copper oxy-chloride which were compatible to all the isolates at lower concentration but showed toxicity at higher levels. This positive effect of sulphur on radial growth of EPF was also reported by Autkar (2004), Jayraman (2005), Kotwal *et al.* (2012) and Usha *et al.* (2014) The fungicides (carbendazim and hexaconazole) showed complete inhibitory action on the growth of *M. anisopliae* (Joshi *et al.*, 2018). The copper fungicides are classified into the upper half of the highly toxic (T) category and are considered incompatible with the entomopathogenic fungus, *B. bassiana* strain ATCC 74040 under *in vitro* conditions (Celar and Kos, 2020).

#### **Mean mycelial weight of *M. anisopliae***

The results showed that there were significant differences among various treatments. The highest (3.787g) mycelial weight was recorded in the treatment T<sub>5</sub> (Thiophanate methyl 70WP). Treatments T<sub>4</sub>-Sulphur 80WDG (3.651g) and T<sub>9</sub>- Metalaxyl 35WS (3.568g) were at par with Thiophanate methyl 70WP. The treatments T<sub>8</sub> (Captan 50WP) and T<sub>2</sub> (Mancozeb 75WP) reported 1.970g and 1.809g mycelial weight respectively. The lowest or no growth was recorded in T<sub>1</sub> (Carbendazim 50WP), T<sub>3</sub> (Copper oxychloride 50WP), T<sub>6</sub> (Hexaconazole 5EC) and T<sub>7</sub> (Difenconazole 25EC). At the same time, control reported 3.763g weight of mycelial mat.

The present findings are in accordance with Niassy *et al.* (2012) who reported that amongst all the pesticides tested, Carbendazim and Copper fungicides were highly toxic to *M. anisopliae* and

*B. bassiana* with lowest or nil biomass on broth media. Applications of the fungicides iprodione and tebuconazole, which are used routinely on *Brassica* crops, are compatible with *M. anisopliae* 389.93 against *D. radicum* under glasshouse conditions, even though these fungicides are inhibitory to fungal growth on SDA medium (Chandler *et al.*, 2005). The fungicides (carbendazim and mancozeb) produce complete growth inhibition (100%), whereas tridemefon and cyperconazole inhibit the growth of *N. rileyi* by (49.65% and 55.25%, respectively) *in vitro* at their recommended dosages (Patil *et al.*, 2014). The fungicides difenoconazole, propiconazole, trifloxystrobin and azoxystrobin are the most harmful products to all biological stages of *M. anisopliae* (strain CG 168) on PDA *in vitro* at recommended field dosage and they should not be applied together with this fungus in tank mixing (Silva *et al.*, 2013).

#### **Mean conidial germination of *M. anisopliae***

The results showed that there were significant differences among various treatments. The highest mean conidial germination 21.60 was recorded in the treatment T<sub>2</sub> (Thiophanate methyl 70WP) while other two treatments *viz.*, T<sub>3</sub> (Metalaxyl 35WS) and T<sub>1</sub> (Sulphur 80WDG) reported mean conidial germination 10.40 and 09.80 respectively and were at par with T<sub>2</sub> (Thiophanate methyl 70WP). At the same time, control recorded 92.08 mean conidial germination.

The present findings are in agreement with Silva *et al.* (2013) and Usha *et al.* (2014) who reported that fungicides caused more than 50 per cent reduction in conidial germination of EPF. The germination inhibition rate of autochthonous isolate of *Metarhizium* spp. increased with the concentration of fungicides (Aby *et al.*, 2015).

#### **Percent mortality of cow pea aphid (*A. craccivora*) by *M. anisopliae* suspension from different treatments**

There were significant differences among various treatments. The highest (63.00%) percent mortality was recorded in the treatment T<sub>2</sub> (Thiophanate methyl 70WP) followed by 43.00 percent mortality in treatment T<sub>3</sub> (Metalaxyl 35WS). The lowest percent mortality 41.00 percent was observed in treatment T<sub>1</sub> (Sulphur 80WDG). At the same time, control reported 83.00 percent mortality.

The present findings are in agreement with Mutimura *et al.* (2010) who stated that fungicides inhibited growth of mycelia and germination of the fungal conidia at all concentrations tested *in vitro*, and also reduced the efficacy of EPF in bioassays against arthropods. The pathogenicity of *M. anisopliae* in combination with fungicides showed highest mortality (33%) rate for adult weevils (*Cosmopolites sordidus*) exposed to conidia produced on media containing high concentrations of Methyl Thiophanate and Propiconazole; while 11 percent for Trifloxystrobin (Aby *et al.*, 2015).

#### **Effect of herbicides on *M. anisopliae***

The *in vitro* effect of herbicides on *M. anisopliae*, was similar to that recorded on insecticides and fungicides (Table 4).

#### **Mean radial growth**

##### **4<sup>th</sup> day after inoculation**

The results showed that there were significant differences among various treatments. The highest (4.7cm) diameter was recorded in the treatment T<sub>1</sub> (Pendimethylene 30EC) and the treatments T<sub>8</sub> (untreated control) (4.6cm), T<sub>7</sub> (Paraquat dichloride 24SL) (4.3cm) and T<sub>6</sub> (Glyphosate 41SL) (4.1cm) were at par with Pendimethalin. The treatment T<sub>5</sub> (2,4-D Dimethyl amine Salt 58WSC) and T<sub>2</sub> (Pyrazosulfuron ethyl) recorded radial growth 3.5cm and 2.1cm respectively while other treatments viz., T<sub>3</sub> (Oxadiargyl 80WP) and T<sub>4</sub> (Oxyflorofen) reported complete inhibition of germination and growth of *M. anisopliae*.

#### **6<sup>th</sup> day after inoculation**

The highest (5.6cm) diameter was recorded in the treatment T<sub>6</sub> (Glyphosate 41SL) (5.6cm) and the treatment T<sub>1</sub> (Pendimethylene 30EC) (5.3cm) was at par with glyphosate. The treatment T<sub>7</sub> (Paraquat dichloride), T<sub>5</sub> (2,4-D Dimethyl amine Salt 58WSC) and T<sub>2</sub> (Pyrazosulfuron ethyl) recorded radial growth 4.7cm, 3.8cm and 2.5cm respectively while other treatments viz., T<sub>3</sub> (Oxadiargyl 80WP) and T<sub>4</sub> (Oxyflorofen) reported complete inhibition of germination and growth of *M. anisopliae*. At the same time untreated control recorded radial diameter of 6.8cm.

#### **8<sup>th</sup> day after inoculation**

The highest (8.8cm) diameter was recorded in the treatment T<sub>6</sub> (Glyphosate 41SL) (8.8cm) and was superior over all the treatments. The treatment T<sub>7</sub> (Paraquat dichloride), T<sub>1</sub> (Pendimethylene 30EC), T<sub>5</sub> (2,4-D Dimethyl amine Salt 58WSC) and T<sub>2</sub> (Pyrazosulfuron ethyl) recorded radial growth 5.3cm, 5.3cm, 4.5cm and 2.8cm respectively while other treatments viz., T<sub>3</sub> (Oxadiargyl 80WP) and T<sub>4</sub> (Oxyflourfen) reported complete inhibition of germination and growth of *M. anisopliae*. At same time untreated control recorded radial diameter of 8.0cm. The results of data 10 and 12 days after inoculation were similar to the readings after 8 days of inoculation.

In general the results indicated that amongst various herbicides tested, glyphosate was found to be the most suitable for *M. anisopliae* followed by pendimethalin. The present findings are in contradictory with Silva *et al.* (2013) who reported a complete suppression of vegetative growth of *M. anisopliae* was caused by 2, 4-D as it expressed good mycelial growth during the present investigations. Most herbicides showed reduced mycelial growth in comparison with the control except for glyphosate. Similar findings were also reported by Silva *et al.* (2013) from Brazil.

#### **Mean mycelial weight of *M. anisopliae***

The highest mycelial growth was recorded in the treatment T<sub>6</sub> (Glyphosate 41SL) (3.928g) and was significantly superior over all the treatments. The second best treatment was T<sub>7</sub> (Paraquat dichloride 24SL) (3.715g), followed by T<sub>1</sub> (Pendimethalin 30EC) (3.206) and T<sub>2</sub> (Pyrazosulfuron ethyl 10WP) (2.690). The other treatments viz., T<sub>3</sub> (Oxadiargyl 80WP) and T<sub>4</sub> (Oxyfluorfen 23.5EC) and T<sub>5</sub> (2,4-D Dimethyl amine Salt 58WSC) reported complete inhibition of germination and growth of *M. anisopliae*. At same time untreated control recorded radial diameter of 3.572g mycelial weight.

The present findings were in agreement with the results of Ambethgar *et al.* (2009) who reported that results obtained in some in vitro and in vivo experiments suggest a sensitivity of EPF to

some herbicides and Celar and Kos (2020) who reported a strong inhibitory effect of some tested herbicides on mycelium growth of EPF in laboratory experiments. Herbicide 2,4-D is most harmful to *M. anisopliae* CG 168 as it completely inhibits conidial germination, vegetative growth and conidiation (Leipelt *et al.*, 2001). The herbicides lambda cyhalothrin and the herbicides glyphosate, bentazon, and imazapic+ imazapyr were found compatible with *M. anisopliae* (Silva *et al.*, 2013).

#### **Mean conidial germination of *M. anisopliae***

The highest mean conidial germination 36.00 was recorded in the treatment T<sub>1</sub> (Pendimethylene 30EC). While other treatments T<sub>5</sub> (Paraquat dichloride 24SL), T<sub>2</sub> (Pyrazosulfuron ethyl 10WP) and T<sub>3</sub> (2,4-D Dimethyl amine Salt 58WSC) reported mean conidial germination 35.25, 30.25 and 20.50 respectively and were at par with the treatment T<sub>1</sub> (Pendimethalin 30EC). The lowest mean conidial germination 09.50 was recorded T<sub>4</sub> (Glyphosate 41SL). At same time untreated control recorded 92.25 mean conidial germination per 10µl of suspension.

The present findings are in line with Silva *et al.* (2013) who reported that all herbicides had a deleterious effect on conidial yield of EPF compared to the untreated control.

#### **Percent mortality of cow pea aphid (*A. craccivora*) by *M. anisopliae* suspension from different treatments**

The highest (73.75%) per cent mortality of *A. craccivora* was recorded in the treatment T<sub>5</sub> (Paraquat dichloride 24SL) and was significantly superior over other treatments. While other treatments *viz.*, T<sub>3</sub> (2,4-D Dimethyl amine Salt 58WSC), T<sub>1</sub> (Pendimethalin 30EC) and T<sub>2</sub> (Pyrazosulfuron ethyl 10WP) reported 65.00 percent, 63.75 percent and 63.75 percent mortality respectively. The lowest (55.00%) per cent mortality was recorded in T<sub>4</sub> (Glyphosate 41SL). At same time untreated control recorded 85.00 percent mortality. Most of the herbicides classified as toxic, but those formulated with imazapyr, glyphosate and metribuzin are considered compatible (Botelho, 2011).

#### **Effect of pesticides on conidia production assessment of *M. anisopliae***

Since all the treatments showed germinated conidia and no report of ungerminated conidia was noticed; the conidia production assessment was said to be 100 percent in all the treatments.

In summary, our findings from *in vitro* compatibility were in agreement with Silva *et al.* (2013) who reported that fungicides are more detrimental to conidial germination, mycelial growth and sporulation of *M. anisopliae* than herbicides and insecticides, and they should not be applied together with EPF. The insecticides exhibited the least degree of toxicity to fungal pathogen, whereas the herbicides had the greatest impact on mycelial growth.

After collecting published results of many different experiments on EPF and pesticides, we observed that fungicides have the highest mycelium growth inhibitory effect, while insecticides and herbicides have a fungistatic effect.. Although the results did not consider the effect of many variables associated with field use of pesticides, our findings are of paramount importance to guide and advise

farmers that they can use *M. anisopliae* safely in combination with pesticides without any adverse effect on fungal virulence and germination.

### Future prospects

The present studies open an area of using EPF together with compatible insecticides as multiple mortality factors against target pests and also help in delaying expression of resistance to new insecticides.

### Conclusions

Our results indicate that the use of combination of *M. anisopliae* with lower dosage of insecticides might become an important component of IPM but at first, this approach must be tested under field conditions.

Integrating insecticides and entomopathogens has a few advantages:

- 1) Such approach will increase pest mortality as well as reduce the lethal time.
- 2) Prolong the use of a particular insecticide by reducing the total amount of insecticide used.
- 3) Minimize environmental contamination and increase human safety.
- 4) Accelerate the mode of action of fungus without compromising the fungus growth from cadavers that is crucial for inducing epizootic in pest population.
- 5) Sublethal dosage of synthetic insecticides can act as physiological stressors or behavioral modifiers, there by predisposing insects to diseases.

To decrease the dose of chemical pesticide used in agriculture, combination of chemical pesticides with biopesticides can be done so as to minimize the negative effect of former on the environment. Acetamaprid and Profenofos require further investigation. Also field evaluation of the interactions between *B. bassiana* and these pesticides should be undertaken to evaluate their effect on pest and beneficial insects. On the other hand, the high toxicity of a product *in vitro* not always means that the same will occur in the field, but is evidence that it may be possible to occur. Thus, when an IPM strategy is devised, it is important to take into account the compatibility of products sprayed on the crop, avoiding the use of most toxic, or using them during seasons when the effect over a natural control agent is minimized.

### References

- Aby, N., Seka, C.L., Traore, S., Kobenan, K., Kouakou, T.H. . Effects of some fungicides and nematicides used in banana plantation on pathological characteristics of *Metarhizium* sp., biological agent control of banana weevil *Cosmopolites sordidus* Gemar (Coleoptera: Curculionidae). *Int. J. Curr. Microbiol. App. Sci.* 2015; 4(6): 1050-65.
- Ambethgar, V., Swamiappan, M., Rabindra, R. J., Rabindran, R. (2009). Influence of some herbicides on in vitro vegetative growth of *Beauveria bassiana* (Balsamo) Vuillemin. *Resistant Pest Management Newsletter* 19: 13-16.
- Araujo, J. J., Marques, E., Oliveira, J. (2009). Potential of *Metarhizium anisopliae* and *Beauveria bassiana* isolates and neem oil to control the Aphid *Lipaphis erysimi* (Kalt.) (Hemiptera: Aphididae), *Neotrop Entomol*, 38(4):520.

- Autkar, S. S. (2004): Fungi Vs Fungi: A potential application of antagonistic properties. M.Sc. (Agril.) Theses (Unpublished) PGI, Dr. PDKV, Akola.
- Babu N, M., Usha, J. and Padmaja, V. (2013). *In-Vitro* evaluation of the entomopathogenic fungal isolates of *Metarhizium anisopliae* for compatibility with pesticides, fungicides and botanicals. *International Journal of Applied biology and Pharmaceutical Technology*, **5**(1): 102-113.
- Celar, F. and Kos, K. (2020). In vitro compatibility of *Beauveria bassiana* strain ATCC 74040 with copper fungicides. *Journal of Applied Entomology*. 144. 10.1111/jen.12743.
- Chandler D, Davidson, G. and Jacobson, R. J. (2005). Laboratory and glasshouse evaluation of fungi against the two-spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae), on tomato, *Lycopersicon esculentum*. *Biocontrol Sci Tech*. **15**: 37– 54.
- Duarte A, Menendez, J. M. and Triguero N. (1992). Estudiopreliminar sobre la compatibilidad de *Metarhizium anisopliae* con algunos plaguicidas quimicos. *Revista Baracoa* **22**: 31-39.
- Ericsson J.D., Todd Kabaluk J., Goettel M.S. & Myers J.H. (2007). Spinosad interacts synergistically with the insect pathogen *Metarhizium anisopliae* against the exotic wireworms *Agriotes lineatus* and *Agriotes obscurus* (Coleoptera: Elateridae). *J. Econ. Entomol.* 100: 31–38.
- Fei Yi, Zou, C., Qiongbo Hu., Meiyang Hu. (2012). The joint action of destruxins and botanical insecticides (rotenone, azadirachtin and paeonolum) against the cotton aphid, *Aphis gossypii* Glover. *Molecules*, **17**: 7533-7542.
- Gang H. and St Leger R. J. (2002). Field studies using a recombinant mycoinsecticide (*Metarhizium anisopliae*) reveal that it is rhizosphere competent. *Appl Environ Microbiol*, 68(12):6383-7. DOI: 10.1128/AEM.68.12.6383-6387.2002.
- Gardner, W.A. and Kinard, D.J. (1998). *In vitro* germination and growth response of two entomogenous fungi to imidacloprid. *Journal of Entomological Science* **33**: 322-324.
- Hemalatha, S., Ramaraju, K. and Jeyarani, S. (2015). Effect of insecticides on the white muscardine fungus, *Beauveria bassiana*. *Indian Journal of Entomology*, **77**(2):134-137
- James, R. R. and Elzen, G. W. (2001). Antagonism between *Beauveria bassiana* and imidacloprid when combined for *Bemisia argentifolii* (Homoptera: Aleyrodidae) control. *Journal of economical Entomology*, **94**: 357– 361.
- Jayaraman, G., (2005). Studies on *Metarhizium anisopliae* (Metchnikoff) against *Helicoverpa armigera* (Hubner)., M.Sc. (Agril.) Thesis (Unpub.) PGI, Dr. PDKV, Akola.
- Jing EeYii, ChoonFah, J. B., Jie Hung, P. K. and Kadir, J. (2016). Synergism of entomopathogenic fungus, *Metarhizium anisopliae* incorporated with fipronil against Oil palm pest subterranean termite, *Coptotermes curvignathus*. *Plant Protect. Sci.*, 52( 1): 35–44
- Johnson D. M. R. L. White R. Pereira and C. J. Geden (2020). *Beauveria bassiana* culturing and harvesting for bioassays with house flies *J. Insect Sci* 20(6). doi:[10.1093/jisesa/ieaa072](https://doi.org/10.1093/jisesa/ieaa072)

- Joshi M., Gaur N. and Pandey R. (2018). Compatibility of entomopathogenic fungi *Beauveria bassiana* and *Metarhizium anisopliae* with selective pesticides. *Journal of Entomological and Zoological Studies*, 6(4): 867-872.
- Kaakeh, W., Reid, B. L. and Bennett, G. W. (1997). Toxicity of Fipronil to German and American cockroaches. *Entomol. Exp. Appl.* **84**: 229-237.
- Khan, S., Bagwan, N. B., Fatima, S. and Iqbal, M. A. (2012). *In vitro* compatibility of two entomopathogenic fungi with selected insecticides, fungicides and plant growth regulators. *Libyan Agric Res Cen J Intl*, **3**: 36-41.
- Klingen, I. and Haukeland, S. (2006). 'The soil as a reservoir for natural enemies of pest insects and mites with emphasis on fungi and nematodes, In: An ecological and societal approach to biological control, eds. J. Eilenberg and H.M.T. Hokkanen, Dordrecht, *The Netherlands: Springer* pp. 145- 211.
- Kotwal, S., Parate, R. L., Mane, S. S. and Deshmukh, V. V. (2012). Effect of *Metarhizium anisopliae* on *Spodoptera litura* and compatibility with chemicals. *International Journal of Science, Environment and Technology*, 1(5): 499 – 505
- Leipelt, M.; Warnecke, D.; Zähringer, U.; Ott, C.; Müller, F.; Hube, B.; Heinz, E. (2001). Glucosylceramide synthases, a gene family responsible for the biosynthesis of glucosylsphingolipids in animals, plants and fungi. *Journal of Biological Chemistry*, 276: 33621–33629.
- Ling Xie, JiHee Han, Jeong Jun Kim<sup>1</sup> and Sang Yeob Lee (2016). Influence of eco-friendly agricultural materials on the germination, mycelial growth and control efficacy OF *Beauveria bassiana* against the sweet potato whitefly. *Academia Journal of Agricultural Research*, **4**(9): 598-606
- Mutumura, C. G., Laing, M. D., Miller, R. M. (2010). *In vitro* effects of flutriafol and azoxystrobin on *Beauveria bassiana* and its efficacy against *Tetranychus urticae*. *Pest Management Science*, 66(7): 773-778
- Neves, P.M.O.J., Hirose, E., Tchujo, P.T. and Moino J.A. (2001). Compatibility of entomopathogenic fungi with neonicotinoids insecticides. *Neotropical Entomology*, **30**: 263–268.
- Niassy, S., Subramanian, S., Gitonga, L. and Ekesi S. (2012). Compatibility of *Metarhizium anisopliae* isolate ICIPE 69 with agrochemicals used in French bean production. *International Journal of Pest Management*, 58(2): 131-137
- Pachamuthu, P., and Kamble, S. T. (2000). *In vivo* study on combined toxicity of *Metarhizium anisopliae* (Deuteromycotina: Hyphomycetes) strain ESC-1 with sublethal doses of chlorpyrifos, propetamphos, and cyfluthrin against German cockroach (Dictyoptera: Blattellidae). *Journal of Economic Entomology*, **93**:60-70.
- Patil, R. K., Bhagat, Y. S., Halappa, B. and Bhat, R. S. (2014). Evaluation of entomopathogenic fungus, *Nomuraea rileyi* (Farlow) samson for the control of groundnut *Spodoptera litura* (F.) and its compatibility with synthetic and botanical pesticides. *J. Biopest.* **7**:106-115



			Day	Day	Day	(g)	(10 µl)	(%)
T <sub>1</sub>	Cypermethrin 25EC	0.0075	2.93	3.43	4.16	3.292	13.00 (3.74)*	40.00 (39.23)**
T <sub>2</sub>	Indoxacarb 14.5SC	0.01	3.83	4.26	4.63	3.787	19.00 (4.47)	80.00 (63.43)
T <sub>3</sub>	Chlorantraniliprole 18.5 SC	0.005	2.43	2.9	3.2	3.187	08.00 (3.00)	51.65 (45.95)
T <sub>4</sub>	Buprofezin 25EC	0.015	6.36	6.86	9.0	3.484	29.33 (5.51)	58.30 (49.78)
T <sub>5</sub>	Profenophos 50EC	0.15	1.73	2.16	2.4	3.784	49.33 (7.09)	28.30 (32.14)
T <sub>6</sub>	Deltamethrin 2.8EC	0.002	0.40	0.56	1.26	3.105	20.00 (4.58)	66.65 (54.73)
T <sub>7</sub>	Dichlorvos 76 EC	0.07	2.06	2.26	2.33	2.386	39.66 (6.38)	76.65 (61.10)
T <sub>8</sub>	Azadirachtin 0.15EC	0.05	6.16	7.03	7.2	6.211	92.00 (9.64)	86.65 (68.97)
T <sub>9</sub>	Emamectin Benzoate 5SG	0.0016	4.3	4.53	4.9	4.052	33.79 (5.90)	40.00 (39.23)
T <sub>10</sub>	Control		5.56	5.9	6.03	3.605	13.00 (3.74)	80.00 (63.43)
			SIG					
S.E. ±			1.05	0.99	1.13	0.43	2.63	1.10
C.D. at 1%			4.28	4.03	4.58	1.75	11.06	4.63

\*Figures in t parentheses are square root transformed values

\*\*Figures in parentheses are arc sin transformed values

**Table 2: Effect of fungicides on *M. anisopliae***

Treat. No.	Treatment	Concentration (%)	Mean radial growth (cm)			Mean mycelial weight (g)	Mean Conidial germination (10 µl)	Mortality of test insect (%)
			4 <sup>th</sup> Day	6 <sup>th</sup> Day	8 <sup>th</sup> Day			
T <sub>1</sub>	Carbendazim 50WP	0.1	0	0	0	0.00	-	-
T <sub>2</sub>	Mancozeb 75WP	0.25	0	0	0	1.809	-	-
T <sub>3</sub>	Copper oxychloride 50WP	0.25	0	0	0	0.00	-	-
T <sub>4</sub>	Sulphur 80WDG	0.2	7.16	9	9	3.651	09.80 (3.29)*	41.00 (39.82)**
T <sub>5</sub>	Thiophanate methyl 70WP	0.1	1.2	1.26	1.26	3.787	21.60 (4.75)	63.00 (52.54)
T <sub>6</sub>	Hexaconazole 5EC	0.05	0.0	0.0	0.0	0.00	-	-
T <sub>7</sub>	Difenconazole 25EC	0.05	0.0	0.0	0.0	0.00	-	-
T <sub>8</sub>	Captan 50WP	0.2	0.0	0.0	0.0	1.970	-	-
T <sub>9</sub>	Metalaxyl 35WS	0.2	2.53	3.3	4.23	3.568	10.40 (3.38)	43.00 (40.98)
T <sub>10</sub>	Control		7.33	7.5	7.5	3.763	92.08 (9.65)	83.00 (65.65)
			SIG					
S.E. ±			0.92	0.60	0.59	0.22	1.50	0.60
C.D. at 1%			3.75	2.43	2.41	0.88	5.50	2.46

\*Figures in parentheses are square root transformed values

\*\*Figures in parentheses are arc sin transformed values

**Table 3: Effect of herbicides on radial growth of *M. anisopliae***

Treat. No.	Treatment	Concentration (%)	Mean radial growth (cm)			Mean mycelial weight (g)	Mean Conidial germination (10 µl)	Mortality of test insect (%)
			4 <sup>th</sup> Day	6 <sup>th</sup> Day	8 <sup>th</sup> Day			
T <sub>1</sub>	Pendimethalin 30EC	0.2	4.7	5.3	5.3	3.206	36.00 (6.08)*	63.75 (52.98)**
T <sub>2</sub>	Pyrazosulfuron ethyl 10WP	0.003	2.1	2.5	2.8	2.690	30.25 (5.59)	63.75 (52.98)
T <sub>3</sub>	Oxadiargyl 80WP	0.016	0.0	0.0	0.0	0.00	-	-
T <sub>4</sub>	Oxyflourfen 23.5EC	0.03	0.0	0.0	0.0	0.00	-	-
T <sub>5</sub>	2,4-D Dimethyl amine Salt 58 WSC	0.025	3.5	3.8	4.5	0.00	20.50 (4.64)	65.00 (53.73)
T <sub>6</sub>	Glyphosate 41SL	0.05	4.1	5.6	8.8	3.928	09.50 (3.24)	55.00 (47.87)
T <sub>7</sub>	Paraquat dichloride 24SL	0.015	4.3	4.7	5.3	3.715	35.25 (6.02)	73.75 (59.18)
T <sub>8</sub>	Control		4.6	6.8	8.0	3.572	92.25 (9.66)	85.00 (67.21)
			SIG					
S.E. ±			0.61	0.75	0.66	0.17	2.04	0.78
C.D. at 1%			2.55	3.16	2.80	0.72	8.51	3.19

\*Figures in parentheses are square root transformed values

\*\*Figures in parentheses are arc sin transformed values