

Effects of Biochar on Yield of Chilli, and Soil chemical properties

ABSTRACT

Aims: To examine the effects of different doses of biochar on the yield of chili and soil chemical properties under the Samtenling condition which represents the wet-subtropical agro-climatic zone of Bhutan.

Study design: Randomized Complete Block Design (RCBD) was used for the study with three replications and five treatments.

Place and Duration of Study: Agriculture Research and Development Centre (ARDC), Samtenling, Sarpang district located in southern Bhutan, between October 2020 and May 2021.

Methodology: Composite soil samples were collected before and after biochar addition and their chemical properties were analyzed at Soil and Plant Analytical Laboratory (SPAL), National Soil Services Centre, Thimphu. One hybrid variety (sv2319HA) of chili was used in all the treatments with five different biochar doses (No biochar addition, biochar addition @ 0.8 t acre⁻¹, 1.6 t acre⁻¹, 3.2 t acre⁻¹ and 4.9 t acre⁻¹). Growth and yield data of chili were collected and analyzed using the multivariate analysis of variance (MANOVA) at $P = .05$ using R studio team (2021), version 1.4.1103.

Results: The yield of chilli significantly differed ($P = .03$) under different doses of biochar but variations in growth were insignificant except the number of branches ($P = .03$). Overall, the yield of chili increased from 0.3 t acre⁻¹ to 0.9 t acre⁻¹ with an increase in biochar rates from 0.8 t acre⁻¹ to 3.2 t acre⁻¹. Biochar addition of 3.2 t acre⁻¹ significantly increased the yield of chili (0.95 t acre⁻¹) as compared to control (0.346 t acre⁻¹). Soil pH, % nitrogen, available P (mg kg⁻¹), available K (mg kg⁻¹), CEC, % BS, and % organic matter increased with increasing quantities of rice husk biochar.

Conclusion: The different doses of biochar addition to the soil had a significantly positive impact on the yield of chili as it also improved the soil's chemical properties. The addition of 3.2 t acre⁻¹ of biochar may be recommended as per the study since the maximum yield of chilli was recorded in soil treated with biochar @ 3.2 t acre⁻¹.

Keywords: Biochar; Chilli; Yield; Growth; Soil chemical properties

1. INTRODUCTION

Biochar is a recent but underutilized soil technology. Biochar (charcoal or agri-coal) is a carbon (C) rich product derived from the pyrolysis of organic material at relatively low temperature (<700°C) [1], having potential use as an amendment to enhance soil properties and agricultural systems productivity. It is mainly composed of stable aromatic forms of organic carbon that do not degrade to CO₂ making it a relatively simple climate change mitigation method [2]. The application of biochar to soil could play a substantial role in mitigating greenhouse gas emissions and climate change effects [3]. The highly porous nature of biochar improves the soil water retention ability as well as plant available water (PAW) [4]. Studies indicated that the application of biochar improves crop production by increasing plant nutrient uptake [5] and improving the physical and chemical properties of soil such as water holding capacity, cation exchange capacity (CEC), adsorption of plant nutrients and soil pH [2][6]. Therefore, it enhances the soil fertility [7] and sustains crop productivity [8] [9] while it also reduces nutrient leaching loss [2].

Chilli is an important ingredient of Bhutanese dietary habits, consumed either as a vegetable or used as food flavoring spice. In Bhutan, the average import and production of chilli from 2014-2018 was 1300.716 MT and 9,261 MT respectively with an average yield of 1.8 t acre⁻¹ [10]. The import restriction in recent years led to an increased need for domestic production. However, water and nutrient deficiency often limits the crop growth and development as chilli is a high nitrogen feeder and sensitive to water stress [11][12]. Studies also reported that the soil fertility status in Bhutan as low to medium and the mean pH value as acidic in nature (<5.6) which indicates the need to improve soil fertility through the application of various soil amendments [13]. The positive effects of biochar application on crop yields in comparison to unamended soil have been reported in various crops such as rice [14], maize [15], soybean [16], cassava [17] and chilli [18]. A study conducted by [19] showed that biochar application increased the yield of maize and cabbage by 6.66% (12.31 t ha⁻¹ to 13.13 t ha⁻¹) and 7.57% (1.98 t ha⁻¹ to 2.13 t ha⁻¹) respectively. Further, [20][21] observed that biochar doses of 6-9 t ha⁻¹ provided the best economic yields in vegetable crops. Therefore, this study was conducted to examine the effects of biochar on soil properties and assess the influence of biochar on growth and yield of chilli under Samtenling condition which represents wet-subtropical agro-climatic zone. In this experiment, we hypothesized that the biochar will significantly (I) increase the yield of chili, and (II) improve the soil physical and chemical properties.

2. MATERIALS AND METHODS

2.1. Study site

The study was conducted at Agriculture Research and Development Centre, Samtenling, Sarpang in 2020 and 2021. The experimental site falls under wet-subtropical agro-ecological zone and is located (26° 54'30" N and 90° 25'55"E) at 375 masl with an average annual rainfall of 2500mm – 5500mm; and maximum and minimum temperature of 35^o C and 12^o C respectively.

Biochar used in this study was produced from rice husk using a traditional farmer's kiln under oxygen limited conditions where the organic waste fuel was burnt in the inner cone shaped structure to heat the rice husk in the outer cylindrical drum for over 5 hours, until no visible smoke was emitted from the chimney. The standard biochar properties and total elemental composition produced from rice husk using this method are as given in Table 1. [22].

Table 1. Properties and Elemental composition of rice husk biochar

Wt. Loss (%)	Ash (%)	C (%)		N (%)	pH	Si (mg/kg)	Al (mg/kg)	Ca (mg/kg)	Mg (mg/kg)	Na (mg/kg)	K (mg/kg)	P (mg/kg)	S (mg/kg)	Fe (mg/kg)	Mn (mg/kg)	Zn (mg/kg)
Rice Husk Biochar	56	44	46	0.63	9.7	193748	212	1340	1683	12353	8502	2983	89	220	454	309

*Wt. Loss: Weight loss

2.2. Experimental methodology

The experiment was laid out using Randomized complete block design with three replications. Five treatments were involved which consisted of various combinations of biochar and farm yard manure (FYM). While the quantity of FYM was constant (i.e. 6 t acre⁻¹) in all experimental plots, addition of biochar in the experimental plots differed across treatments as follows: 1. No biochar addition (control; C), 2. Addition of biochar @ 0.8 t acre⁻¹ (B₁), 3. Addition of biochar @ 1.6 t acre⁻¹ (B₂), 4. Addition of biochar @ 3.2 t acre⁻¹ (B₃) and 5. Addition of biochar @ 4.8 t acre⁻¹ (B₄). The chilli variety used in this study was 'sv2319HA' which is a

high yielding hybrid variety. The seeds were sown in October in soil potting media in pluck trays under greenhouse condition and were transplanted in uniform rows 30 days after sowing in plot size of 6 m² (3m x 2m). Chilli seedlings were spaced 70 cm apart with 70 cm between rows. Uniform intercultural operations were applied to all treatments.

2.3 Data collection and analysis and statistical analysis

Composite soil samples were collected before and after biochar addition using random sampling methods from the respective plots. The samples were then analyzed at Soil and Plant Analytical Laboratory (SPAL), National Soil Services Centre, Thimphu to determine the changes in pH, OM, N, Avail P, Avail K and CEC as affected by biochar addition.

Data on growth parameters - plant height (PH), number of branches (NB), stem diameter (SD) and plant canopy (PC) were recorded at 20 days interval after transplantation upto 40 days. Yield parameters - single pod weight (SPW), number of pods/plant (NP), pod length (PL), pod diameter (PD), Pod weight/plant (PWP), and yield (Y) were estimated from three harvests.

The multivariate analysis of variance (MANOVA) for data on all parameters was carried out at $P = .05$ using R studio team (2021), version 1.4.1103. The packages used for statistical analysis were “dplyr” [23], and “agricolae” [24]. Post hoc analysis was conducted using Tukey’s HSD for observations with significant variations among the treatments.

3. RESULTS AND DISCUSSION

3.1 Effect of biochar application on growth and yield of chilli

Significant variations were observed among the treatments only in terms of yield, pod weight/plant, number of pods/plant and number of branches/plant (Table 2). However, results also indicated an overall trend of linear increase in fruit yield and plant traits of chilli with the corresponding increase in biochar addition till B₃. With further increase in biochar rate from B₃, there was a decrease in yield and plant height, plant canopy, pod weight, number of pods and pod size at B₄.

The biochar application has significantly influenced the yield of chilli. Although an increase in biochar rates led to a corresponding increase in the yield, application of 3.2 t acre⁻¹ (B₃) significantly increased the yield (0.95 t acre⁻¹) only over control (0.346 t acre⁻¹). However, the yield of chilli decreased at B₄. Pod weight/plant of chilli was significantly affected across the treatments ($P = .03$). Similar to yield, B₃ has the highest pod weight/plant (93.99g) and is significantly greater than C - Control (34.2g). The number of pods/plants also differed significantly among the treatments ($P = .04$). In relation to C (control) significantly higher number of pods were recorded in B₂ (27). The biochar application did not significantly influence plant height, stem diameter, plant canopy, single pod weight and pod size although there was an overall increasing trend till B₃.

Table 2. Effect of different doses of biochar on yield and yield attributes of Chilli

T	NB	PH (cm)	SD (cm)	PC (cm)	SPW (g)	NP	PL (cm)	PD (cm)	PWP (g)	Yield (t acre ⁻¹)
C	23.05 ^b	30.70	0.63	25.01	2.99	13.79 ^b	7.69	0.74	34.24 ^b	0.34 ^b
B ₁	23.72 ^b	33.86	0.67	27.81	3.01	24.11 ^{ab}	7.74	0.77	42.94 ^{ab}	0.43 ^{ab}
B ₂	31.55 ^{ab}	42.16	0.84	32.29	3.19	27.14 ^a	7.99	0.76	78.39 ^{ab}	0.79 ^{ab}

B₃	47.41 ^a	44.56	0.96	39.28	3.36	26.48 ^{ab}	7.66	0.76	93.99 ^a	0.95 ^a
B₄	33.36 ^{ab}	39.65	0.83	33.70	2.84	22.51 ^{ab}	7.52	0.74	57.20 ^{ab}	0.57 ^{ab}
Pvalue	.03	.07	.07	.052	.10	.04	.76	.71	.03	.03
CV (%)	25.95	14.95	16.89	15.95	6.90	21.03	5.70	3.93	33.40	33.38
SD	11.42	7.34	0.16	6.51	0.25	6.29	0.39	0.02	28.48	0.28

*Means with same letters are not significant at $P=0.05$

T. Treatment, C. Control (No biochar application), B₁. Biochar application @ 0.8 t acre⁻¹, B₂. Biochar application @ 1.6 t acre⁻¹, B₃. Biochar application @ 3.2 t acre⁻¹, B₄. Biochar application @ 4.8 t acre⁻¹, NB. Number of branches, PH. Plant height, SD. Stem diameter, PC. Plant canopy, SPW. Single pod weight, NP. Number of pods/plant, PL. Pod length, PD. Pod diameter, PWP. Pod weight/plant, CV. Coefficient of variation, SD. Standard deviation.

The findings of this study align with the review of [25][26] where increased growth of plants due to increased plant photosynthetic rate and enhanced root elongation stimulating plant growth and yield through biochar application was reported. Similarly, an increase in yield of red chilli by 66% for biochar-treated soil was found over the yield in pure acid soil although only minimal difference between biochar treated soil was found suggesting that larger quantities of biochar do not evidently affect crop yield [26]. Biochar application also significantly increased yield, dry matter content and specific gravity of potato tubers [27].

The observed increase in yield and yield components could be due to improved soil physical and chemical properties as reported in various studies: reduced bulk density (-4.5% and -6% with addition of 2.25 t ha⁻¹ and 4.5 t ha⁻¹ respectively) [28], increased water holding capacity (25 to 36% increase) with 7% biochar by weight addition [29] and 11% increase [30], liming effect (20 g kg⁻¹ biochar increased soil pH by almost 1 pH unit) [31], and enhanced nutrient availability (significant increases in N (up to 7%), organic C (up to 69%), and P, K, Mg and Ca, [31], increased soluble P and P retention by biochar [32], adsorption of salts [33] [34]. The increase in yield and yield components in the experiment suggests that biochar application contributed to the pool of nutrients available in the soil as biochar itself generally contains high densities of nutrient. [8] Reported a higher increase of 20% - 120% in crop productivity with biochar addition. Maize grain yield can be significantly increased by 98% - 150% after the application of biochar [35]. [36] reported 8.8% - 14% increase in rice yields. A study also observed that biochar application at 10% (w/w) reduced cumulative leaching of P (37.7%), NH₄⁺ (50.2%), and nitrate (NO₃⁻) which could either be due to the enhanced adsorption of these ions to the surface of biochar, the increased immobilization by the greater microbial biomass resulting from biochar addition, or both [37].

In addition, growing crops with biochar resulted in 11% increase in average yield on dry land soils [38]. [39] and [25] observed an increase in the OM content of sandy soil by (42-72%) and (18-70%) respectively. In sandy loam, biochar application translated to greater water use efficiency for plants grown under moisture limitation [40].

The chilli plants performed significantly better in the plots amended with biochar when compared to non-amended plots (Table 2). It may be attributed to the long term beneficial effects of biochar application on microbial growth upon release in soil solution as reported in a review conducted by [25]. Few studies also reported 6.6 to 31.2% higher fungal abundance upon biochar application to the soil which may be due to its suppressive effect on pathogens [41]. Corroborating earlier findings another study stated that biochar application to an asparagus field soil led to a decrease in root lesions caused by *Fusarium oxysporum*, *F. asparagi*, and *F. proliferatum* compared to the non-amended control [42].

The decrease in yield, yield components and growth in chilli with a higher dose of biochar application in the current study agrees with the findings of [43] and [44] where both reported the negative effect on microbial biomass when biochar was applied at higher concentration. Similarly, [45] also reported the presence of volatile compounds such as benzene in biochar which had detrimental effects on soil microbial biomass. [46] also noted that high concentrations of biochar could have a negative impact on soil microbes through the suppression of biological N fixation. Similarly, [26] also reported that larger quantities of biochar do not substantially affect crop growth in red chilli grown in heavy acid soil.

3.1.1. Correlation for quantitative traits of chilli

The correlation analysis for plant traits (PH, NB, SD, PC) and fruit traits (SPW, NP, PL, PD and PWP) of chilli revealed high (>0.7) significant ($P < .01$) positive correlation of PWP (fruit yield) with PH, NB, SD and PC (Table 3). [47] Also reported significant and positive correlation for fruit yield per plant with fruit weight and number of branches in chilli. [48] Reported similar results where number of branches, stem diameter and canopy width had positively significant phenotypic and genotypic correlations with fruit yield. PWP was positively correlated with SPW, NP, PL and PD although it was non-significant ($P > .05$). [489] also reported a positive association of fruit yield with fruit length and fruit diameter in a similar correlation study carried out in hot pepper.

Table 3. Pearson's correlation for quantitative traits of chilli

	PH (cm)	NB	SD (cm)	PC (cm)	SPW (g)	NP	PL (cm)	PD (cm)	PWP (kg)
PH (cm)	1.00**								
NB	0.77*	1.00**							
SD	0.92**	0.87**	1.00**						
PC (cm)	0.83**	0.84**	0.92**	1.00**					
SPW (g)	0.31	0.62	0.43	0.37	1.00**				
NP	0.36	0.30	0.38	0.39	0.45	1.00**			
PL (cm)	0.33	0.40	0.38	0.15	0.57	0.15	1.00**		
PD (cm)	0.35	0.47	0.46	0.52	0.50	0.34	0.48	1.00**	
PWP	0.84**	0.84**	0.90**	0.79**	0.61	0.59	0.48	0.46	1.00**

*Significant at $P = .05$

**Significant at $P = .01$

PH. Plant height, NP. Number of pods/plant, NB. Number of branches, PL. Pod length, SD. Stem diameter, PD. Pod diameter, PC. Plant canopy, PWP, Pod weight/plant, SPW: Single pod weight.

While PH was significantly and positively correlated with NB, SD, PC and PWP, NP had weak positive correlation (<0.2) with PL. [50] also found positive correlation between PH and SD, and negative correlation for NP and PL in a similar correlation analysis conducted in chilli. NB had high positive (>0.7) significant correlation with PC which is plausible since higher number of branches could result in a larger canopy width. Correlation analysis helps determine the traits on which selection should be based for plant breeding purposes [51]. Hence, as per the results in the study, strong positive correlations of plant traits with fruit yield (PWP) can be considered for indirect selection in breeding programs.

3.2. Effect of biochar on soil chemical properties

The initial soil condition of the study area had an average pH of 4.3 (strongly acidic) and sandy loam texture. The soil had very low N, P and K content with an average value of 0.20% and 23.08 mg/kg and 7.4 mg/kg respectively. Average Cation exchange capacity, soil carbon and organic matter were also recorded with 16.3me/100g, 1.8% and 3.1% respectively. All the values indicated that the soil had poor fertility and only a small amount of nutrients were available to the plants. We recorded changes in soil chemical properties caused by the application of biochar into the soil. The addition of biochar significantly influenced the soil pH. Changes in avail N, P, K, C, OM and CEC were also recorded, although no significant differences were recorded statistically (table 4).

The increase in pH was significantly different between control and treatments ($P = .00$) with the maximum increase in B_1 (1.28). However, progressively increasing the amount of biochar from B_1 to B_4 did not substantially lead to a corresponding increase in soil pH. Although soil available P, K and CEC also increased with biochar addition, the increase was not significantly different among the treatments ($p > 0.05$). The maximum increase in P, K and CEC was observed in B_3 , B_4 and B_1 respectively.

Available N decreased with biochar application although the extent of decrease was not significant between the treatments ($P > 0.05$). The maximum and minimum decline were observed in B_2 (-0.10 mg) and B_1 (-0.02 mg). Soil carbon content decreased with application of rice husk biochar except for B_2 (0.23 %). However, the decrease in C was not significant across the treatments ($p = 0.52$). C (-0.17%), B_1 (-0.05 %) and B_3 (-0.51 %) indicated decline in O.M. content while B_2 (0.40 %) and B_4 (0.11 %) showed an increase in O.M.

Table 4. Between-treatments comparison of mean differences observed in soil properties before and after biochar addition

Treatment	pH ^{2*} -pH ^{1*}	N ² -N ¹ (%)	P ² -P ¹ (mg/kg)	K ² -K ¹ (mg/kg)	C ² -C ¹ (%)	OM ² -OM ¹ (%)	CEC ² -CEC ¹ (me/100g)
C	0.10 ^a	-0.05	10.99	16.50	-0.10	-0.17	-3.66
B_1	1.28 ^b	-0.02	27.85	11.53	-0.03	-0.05	11.62
B_2	0.85 ^b	-0.10	20.43	18.55	0.23	0.40	3.20
B_3	1.13 ^b	-0.07	29.59	11.85	-0.30	-0.51	11.44
B_4	1.11 ^b	-0.06	8.49	18.84	-0.06	0.11	2.32
P value	.00	0.52	0.51	0.96	0.52	0.52	0.22
C.V.	55.70	0.00	75.10	45.80	113.40	113.40	49.60

Means with the same letters are not significantly different at $P = .05$.

²Value after biochar addition

¹Value before biochar addition

Biochar addition induced overall changes in the soil physical and chemical properties. The increase in pH due to biochar addition was observed in various studies [26][52]. This increase in soil pH may be due to the

high pH value of added biochar which can increase soil base saturation, decrease the level of exchangeable aluminum, and consume soil protons [53]. The increase in pH may also be explained by the alkalinity of added biochar which can act as a liming agent in an acidic soil [54][55]. Biochar reduces soil acidity by 31.9% through liming effect [56] [57] [58] [59]. Biochar addition alters nutrient cycles by directly adding nutrients (N, P, K, Ca, Mg) in biochar to the soil [60] and by indirectly influencing the exchange of nutrients due to increased reactive surfaces [61] [62]. Several mechanisms have been proposed for the increase of soil nutrient availability including 1) the initial addition of soluble nutrients from biochar [63] and the mineralization of the labile fraction of biochar [64]; 2) reduction of nutrient leaching due to biochar's physicochemical properties [65]; and 3) reduced N losses by ammonia volatilization and N₂O from denitrification [66] [67].

Biochar application has been found to improve soil phosphorus availability to plants under arid and semiarid agro-ecosystems since biochar alters soil available P by acting as a P source itself, altering P solubility through changes in soil pH [68] or by promoting P solubilizing bacteria [69]. Biochar has a significant amount of soluble P salts generated during the charring of organic materials [70]. According to [71] biochar's liming impact on acid soil causes Al and Fe to precipitate as Fe(OH)₃ and Al(OH)₃, increasing phosphorus availability in the soil system; P availability is greatest in the pH range 5.5-6.0.

The application of biochar increased % organic matter, available K (mg/kg) and CEC which agrees with the study conducted by [26] who reported that biochar application improves acid soil properties (pH, % OM, and % OC) to the levels needed to support plant growth. The possible reason for increased K and CEC might be due to the interaction and reaction of biochar with soil in the short term, such as adsorption and desorption, dissolution, precipitation and redox reactions [72]. It could also be due to the presence of strong carboxylic and phenolic functional groups on the surface of the biochar particles [73] [25] or due to the presence of cation exchange sites on its surface. The increase in K may be due to biochar's porous structure, large surface area, and negative surface charge which can increase the soil's CEC and allow for the retention of nutrients, including K [74].

Biochar has also been shown to increase cation exchange capacity (CEC) by 1.9% [74] which increases the potential for sorption of many organic and inorganic substances, including essential plant nutrients [75] [76]. Applying biochar into soil increases CEC over time due to biochar surface oxidation and the abundance of negatively charged surface functional groups [77][78]. A 2-year field experiment in degraded uplands of East Java, Indonesia (tropical) reported an increase in CEC on biochar application which was attributed to high surface negative charge resulting from oxidation of carboxylic and phenolic groups of biochar [17]. [79] indicated that the biochar treatments significantly increased CEC by 4 to 30 % compared to the controls. Similarly, [80] also reported a 21 % increase in CEC in soil following application of biochar produced from rice husk and sawdust. The Organic matter content of the soil increased with increasing doses of biochar which could be because biochar also undergoes biodegradation, ages and weathers, further increasing CEC [76] although it is considered stable in the soil system [81].

The current study found a decrease in Available N and corresponding increase in pH. [82] stated that the decline in soil nitrogen is caused by the increase in soil pH promoting the transformation of ammonium nitrogen into nitrate nitrogen leading to reduction of available soil nitrogen. Few studies also indicated that the biochar composed of 0.55 – 1.13 % N suggesting that the biochar application is not increasing the amount of nitrogen available in the soil [83]. The rates tested in this study were closely matched by some of those in the meta-analysis (1.5, 3, 5, 8 Mg ha⁻¹). These rates did not show significant changes due to biochar application which suggests that very small fractions of biochar N are in soluble or readily mineralized forms.




Figure 1: Values of soil pH, Avail N, Avail P, Avail K, Soil carbon, OM and CEC before and after application of biochar.

C.Control (No biochar application), B₁. Biochar application @ 0.8 t acre⁻¹, B₂. Biochar application @ 1.6 t acre⁻¹, B₃. Biochar application @ 3.2 t acre⁻¹, B₄. Biochar application @ 4.9 t acre⁻¹, pH 1. Soil pH before biochar addition, pH 2. Soil pH after biochar addition, N 1. Available nitrogen before biochar addition, N 2. Available nitrogen after biochar addition, C 1. Soil carbon before biochar addition, C 2. Soil carbon after biochar addition, O.M. 1. Soil organic matter before biochar addition, O.M. 2. Soil organic matter after biochar addition, P 1. Available phosphorus before biochar addition, P 2. Available phosphorus after biochar addition, K 1. Available potassium before biochar addition, K 2. Available potassium after biochar addition, CEC 1. Cation exchange capacity before biochar addition, CEC 2. Cation exchange capacity after biochar addition.

4. CONCLUSION

The subtropical region of Bhutan experiences annual rainfall ranging from 2500-5000 mm making the soil resource very vulnerable to nutrient leaching, erosion losses, and soil degradation. To reduce land degradation and nutrient leaching losses we must adopt new management options to improve soil productivity and ensure sustainable use of land. The application of biochar proved to improve the soil's

chemical properties as well as the performance of chili through the findings of this study. The different doses of biochar studied in this experiment favored the improvement of soil chemical properties such as soil pH, P, K, CEC, O.M, and C. The yield of chili also increased in soils treated with biochar with the maximum yield of 0.975 t acre⁻¹ obtained with the use of 3.2 t acre⁻¹ of biochar. To improve the soil properties and production of chili, biochar dose @3.2 t acre⁻¹ for acidic soil of sub-tropical regions is recommended as per the findings of the study. However, further research pertaining to biochar addition in different doses and its impact on crop yields is necessary to further validate the findings.

References:

1. Lehmann J., Joseph S. Biochar for environmental management: science and technology.2009.
2. Sohi S, Lopez-Capel E, Krull E, Bol R. Biochar, climate change and soil: A review to guide future research.2009.CSIROhttps://www.researchgate.net/publication/228656328_Biochar_Climate_Change_and_Soil_A_Review_to_Guide_Future_Research
3. Woolf, D., Amonette, J. E., Street-Perrott, F. A., Lehmann, J., & Joseph, S. Sustainable biochar to mitigate global climate change. *Nature Communications*. 2010;1(1). doi:10.1038/ncomms1053
4. Mishra, A. & Shinogi, Y. Effects of biochar on soil physio-chemical propertie. Proceedings of International Exchange and Innovation Conference on Engineering & Sciences (IEICES).2018;4:102-105 <https://doi.org/10.15017/1961296>
5. Major, J., M. Rondon, D. Molina, S. J. Riha and J. Lehmann (2010b). Maize yield and nutrition during 4 years after biochar application to a Colombian savanna oxisol. *Plant and Soil* 333(1-2): 117-128.
6. Lehmann, J. & Joseph, S. Biochar for environmental management: An introduction. *Biochar for Environmental Management*, 2015; 33-46. Doi: <https://doi.org/10.4324/9780203762264>
7. Kloss S., Zehetner F., Wimmer B., Buecker J., Rempt F., Soja G. (2014): Biochar application to temperate soils: Effects on soil fertility and crop growth under greenhouse conditions. *Journal of Plant Nutrition and Soil Science*, 177: 3–15
8. Lehmann J, Rondon M. Bio-char soil management on highly-weathered soils in the humid tropics. In N. T. Uphoff et al. (Eds), *Biological approaches to sustainable soil systems*. Crc/Taylor & Francis, Boca Raton. 2006. Link: <https://bit.ly/2HU6C2C>
9. Asai H, Samson BK, Stephan HM, Songyikhangsuthor K, Homma K, Kiyono Y. et al. Biochar amendment techniques for upland rice production in Northern Laos. 1. Soil physical properties, leaf SPAD and grain yield. 2009;<https://tohoku.pure.elsevier.com/en/publications/biochar-amendment-techniques-for-upland-rice-production-in-northe>
10. Bhutan RNR statistics, 2018. <https://www.doa.gov.bt/wp-content/uploads/2020/06/RNR-statistics-2018.pdf>
11. Walters, A.; Hughes, C. Chili Pepper Information Sheet for Afghanistan; Univ. California Regents: Davis, CA, USA, 2004; Available online: http://afghanag.ucdavis.edu/a_horticulture/row-crops/peppers/IS_Chili_Peppers.pdf (accessed on 20 August 2016).
12. Walters, S., & Jha, A. Sustaining Chili Pepper Production in Afghanistan through Better Irrigation Practices and Management. *Agriculture*. 2016; 6(4):62. MDPI AG. Retrieved from <http://dx.doi.org/10.3390/agriculture6040062>
13. Chhetri, S., Gurung, T.R., Uden, T. Gurung, M. & Jamtsho, K.L. "Assessment of soil fertility status using soil nutrient index in three land use systems in Bhutan" *Bhutanese Journal of Agriculture*. 2020;2(1):60-

[72https://www.bja.gov.bt/assessment-of-soil-fertility-status-using-soil-nutrient-index-in-three-landuse-systems-in-bhutan/](https://www.bja.gov.bt/assessment-of-soil-fertility-status-using-soil-nutrient-index-in-three-landuse-systems-in-bhutan/)

14. Masulili, A., Utomo, W.H. and Syechfani. Rice husk biochar for rice-based cropping system in acid soil and the characteristics of rice husk biochar and its influence on the properties of acid sulphate soils and rice growth in West Kalimantan, Indonesia," J. Agric. Sci. (Canada). 2010; 3:25–33.<https://doi.org/10.5539/jas.v2n1p39>
15. Utomo, W.H., Sukartono, Kusuma, Z. and Nugroho, W.H. "Soil fertility status, nutrient uptake, and maize (*Zea mays* L.) Yield following biochar application on sandy soils of Lombok, Indonesia," J. Trop. Agric. 2011; 49:47–52. <http://jtropag.kau.in/index.php/ojs2/article/view/236/236>
16. Tagoe, S.O., Horiuchi, T. & Matsui, T. Effects of carbonized and dried chicken manures on the growth, yield, and N content of soybean. *Plant* .2008; 06:211–220.<https://doi.org/10.1007/s11104-008-9573-9>
17. Islami, T. Guritno, B., Basuki, N. And Suryanto, A. "Biochar for sustaining productivity of cassava based cropping systems in the degraded lands of East Java, Indonesia," Journal of Tropical Agriculture. 2011; 49:40-46. <http://jtropag.kau.in/index.php/ojs2/article/view/235>
18. Wisnubroto, E.I., Utomo, W.H. and Indrayatie, E.R. "Residual Effect of Biochar on Growth and Yield of Red Chili (*Capsicum Annum* L.)," Journal of Advanced Agricultural Technologies. 2017;4(1):28-31.Doi: 10.18178/joaat.4.1.28-31
19. A Akolgo, G., Kemausuor, F., Awafo, E.A., & Amankwah, E. Biochar as a Soil Amendment Tool: Effects on Soil Properties and Yield of Maize and Cabbage in Brong-Ahafo Region, Ghana. *Open Journal of Soil Science*. 2020;10(3):91-108. <http://dx.doi.org/10.4236/ojss.2020.103005>
20. Situmeang, Y.P. & Suarta, M. The Effect of Biochar Bamboo on Growth and Results of Kangkung (*Ipomoea Reptans* P.). *Journal of Biological and Chemical Research (JBCR)*. 2018; 35 (2);463-468. <https://core.ac.uk/download/pdf/160644344.pdf>
21. Situmeang Y P, Suarta M, Irianto I K and Andriani A A S P R. Biochar Bamboo Application on Growth and Yield of Red Amaranth (*Amaranthus tricolor* L.) *IOP Conference Series: Materials Science and Engineering*. 2018; 434<https://iopscience.iop.org/article/10.1088/1757-899X/434/1/012231>
22. Prakongkep N., Gilkes R.J., Wiryakitnateekul W., Duangchan A. & Darunsontaya. The Effects of Pyrolysis Conditions on the Chemical and Physical Properties of Rice Husk Biochar: *International Journal of Material Science (IJMSC)*. 2013;3(3) https://www.researchgate.net/publication/283347475_The_Effects_of_Pyrolysis_Conditions_on_the_Chemical_and_Physical_Properties_of_Rice_Husk_Biochar
23. Hadley Wickham, Romain François, Lionel Henry and Kirill Müller (2022). *dplyr: A Grammar of Data Manipulation*. <https://dplyr.tidyverse.org>, <https://github.com/tidyverse/dplyr>.
24. Felipe de Mendiburu and Muhammad Yaseen(2020). *agricolae: Statistical Procedures for Agricultural Research*. R package version 1.4. 0 , <https://myaseen208.github.io/agricolae/><https://cran.r-project.org/package=agricolae>
25. Alkharabsheh, H.M., Seleiman, M.F. Battaglia, M.L. Shami,A., Jalal, R.S., Alhammad, B.A. et al. Biochar and Its Broad Impacts in Soil Quality and Fertility, Nutrient Leaching and Crop Productivity: A Review *Agronomy*. 2021; 11, 993. <https://doi.org/10.3390/agronomy11050993>.
26. Hanpattanakit P., Vanitchung S., Saeng-Ngam S., Pearaksa P. Effect of Biochar on Red Chili Growth and Production in Heavy Acid Soil, *Chemical Engineering Transactions*. 2021;83: 283-288 DOI:10.3303/CET2183048
27. Mollick et al., 2020. Effect of Biochar on Yield and Quality of Potato (*Solanum tuberosum*) Tuber. *International Journal of Bio-resource and Stress Management* 2020, 11(5), 445-450. [HTTPS://DOI.ORG/10.23910/1.2020.2140](https://doi.org/10.23910/1.2020.2140)
28. Chen HX, Du ZL, Guo W, Zhang QZ. Effects of biochar amendment on cropland soil bulk density, cation exchange capacity, and particulate organic matter content in the North China plain. *Ying Yong Sheng Tai Xue Bao*. 2011;22(11):2930-2934
29. Kinney, T.J., C.A. Masiello, B. Dugan, W.C. Hockaday, M.R. Dean, K. Zygourakis, et al. 2012. Hydrologic properties of biochars produced at different temperatures. *Biomass and Bioenergy* 41: 34-43. doi:<http://dx.doi.org/10.1016/j.biombioe.2012.01.033>

30. Karhu, K., T. Mattila, I. Bergström and K. Regina. 2011. Biochar addition to agricultural soil increased CH₄ uptake and water holding capacity. Results from a short-term pilot field study. *Agriculture, Ecosystems & Environment* 140: 309-313. doi:<http://dx.doi.org/10.1016/j.agee.2010.12.005>.
31. Laird, D., N. P. Rogovska, M. Garcia-Perez, H. P. Collins, J. D. Streubel and M. Smith. Pyrolysis and biochar. Sustainable alternative fuel feedstocks 11 opportunities, challenges and roadmaps for six U.S. regions. R. Braun, D. Karlen and D. Johnson. Atlanta, GA, Soil and Water Conservation Society – Proceedings of the Sustainable Feedstocks for Advance Biofuels Workshop. 2010.
32. Mukherjee, A. and Zimmerman, A.R. (2013) Organic Carbon and Nutrient Release from a Range of Laboratory-Produced Biochars and Biochar-Soil Mixtures. *Geoderma*, 193, 122-130. <https://doi.org/10.1016/j.geoderma.2012.10.002>
33. Park, J.H., Choppala, G.K., Bolan, N.S., Chung, J.W. and Chuasavathi, T. (2011) Biochar Reduces the Bioavailability and Phytotoxicity of Heavy Metals. *Plant and Soil*, 348, 439-451. <https://doi.org/10.1007/s11104-011-0948-y>
34. Lashari, M.S., Liu, Y.M., Li, L.Q., Pan, W.N., Fu, J.Y., Pan, G.X., Zheng, J.F., Zheng, J.W., Zhang, X.H. and Yu, X.Y. Effects of Amendment of Biochar-Manure Compost in Conjunction with Pyrolygneous Solution on Soil Quality and Wheat Yield of a Salt-Stressed Cropland from Central China Great Plain. *Field Crops Research*, 2013;144, 113-118. <https://doi.org/10.1016/j.fcr.2012.11.015>
35. Uzoma, K.C., Inoue, M., Andry, H., Fujimaki, H., Zahoor, A. and Nishihara, E. Effect of Cow Manure Biochar on Maize Productivity under Sandy Soil Condition. *Soil Use and Management*, 2011; 27, 205-212. <https://doi.org/10.1111/j.1475-2743.2011.00340.x>
36. Zhang, H.H., Lin, K.D., Wang, H.L. and Gan, J. Effect of Pinus radiata Derived Biochars on Soil Sorption and Desorption of Phenanthrene. *Environmental pollution*. 2010;158, 2821-2825. <https://doi.org/10.1016/j.envpol.2010.06.025>
37. Rubin, R.L.; Anderson, T.R.; Ballantine, K.A. Biochar simultaneously reduces nutrient leaching and greenhouse gas emissions in restored wetland soils.2020;11:1–11 <https://pubag.nal.usda.gov/catalog/7219949>
38. Liu, X., Zhang, A., Ji, C., Joseph, S., Bian, R., Li, L. et al. Bio char's effect on crop productivity and the dependence on EXPERIMENTAL conditions—a meta-analysis of LITERATURE data. *Plant and Soil*, 2013; 373(1-2): 583–594. <https://doi.org/10.1007/s11104-013-1806-x>
39. El-Naggar, A.; El-Naggar, A.H.; Shaheen, S.M.; Sarkar, B.; Chang, S.X.; Tsang, D.C.W.; Rinklebe, J.; Ok, Y.S. Biochar composition-dependent impacts on soil nutrient release, carbon mineralization, and potential environmental risk: A review. *J. Environ. Manag.* **2019**; 241: 458–467. [[Google Scholar](#)] [[CrossRef](#)]
40. Batool, A., Taj, S., Rashid, A., Khalid, A., Qadeer, S., Saleem, A. R. et al. Potential of soil amendments (Biochar and Gypsum) in increasing water use efficiency of *Abelmoschus esculentus* L. Moench [Abstract]. *Frontiers in Plant Science*, 2015; 6. <https://doi.org/10.3389/fpls.2015.00733>
41. Matsubara Y, Hasegawa N, Fukui H. Incidence of Fusarium root rot in asparagus seedlings infected with arbuscular mycorrhizal fungus as affected by several soil amendments. *Journal of the Japanese Society for Horticultural Science*. 2002;71:370-374 <https://doi.org/10.2503/jjshs.71.370>
42. Elmer, Wade & Pignatello, Joseph. Effect of Biochar Amendments on Mycorrhizal Associations and Fusarium Crown and Root Rot of Asparagus in Replant Soils. *Plant Disease*.2011; 95:960-966. DOI: 10.1094/PDIS-10-10-0741
43. Zhang, Q.-z., Dijkstra, F. A., Liu, X.-r., Wang, Y.-d., Huang, J. & Lu, N. Effects of biochar on soil microbial biomass after four years of consecutive application in the North China Plain. *PLoS ONE* 2014;9(7):e102062 <https://doi.org/10.1371/journal.pone.0102062>
44. Rondon, M. A., Lehmann, J., Ramírez, J. & Hurtado, M. Biological nitrogen fixation by common beans (*Phaseolus vulgaris* L.) increases with bio-char additions. *Biology and Fertility of Soils*. 2007; 43(6): 699-708. <http://dx.doi.org/10.1007/s00374-006-0152-z#citeas>

45. Girvan MS, Campbell CD, Killham K, Prosser JI, Glover LA.. Bacterial diversity promotes community stability and functional resilience after perturbation. *Environ Microbiol.*2005; 7(3): 301-313. DOI:10.1111/j.1462-2920.2005.00695.x
46. Horel, Á., Gelybó, G., Potyó, I., Pokovai, K. & Bakacsi, Z. Soil nutrient dynamics and nitrogen fixation rate changes over plant growth in temperate soil. *Agronomy.* 2019; 9(4):179; doi: 110.3390/agronomy9040179.
47. Bijalwan, P. & Mishra, A.C. Correlation and path coefficient analysis in chili (*Capsicum annum* L.) for yield and yield attributing traits. *International journal of science and research (IJSR).* 2016; 5(3):1589-1592. https://www.ijsr.net/search_index_results_paperid.php?id=NOV162097
48. Shumbulo A, Nigussie M, Alamerew S (2017) Correlation and Path Coefficient Analysis of Hot Pepper (*Capsicum annum* L.) Genotypes for Yield and its Components in Ethiopia. *Adv Crop Sci Tech* 5: 277. doi: 10.4172/2329-8863.1000277
49. Belay, F. & Tsehaye, Y. Variability, Association and Path Coefficient Analysis of Green Pod Yield and Yield Components of Hot pepper (*Capsicum annum* L.) Landraces at Mereb Lehke, Northern Ethiopia. *Journal of Plant Breeding and Crop Science.* 2019;12(1):58-69, DOI:10.5897/JPBCS2019.0840
50. Silva, A.R.D, Rego, E.R.D, Pessoa, A.M.D.S & Rego, M.M.D. Correlation network analysis between phenotypic and genotypic traits of chilli pepper” *Pesquisa Agropecuária Brasileira.* 2016; 51 (4):372-377. <https://doi.org/10.1590/S0100-204X2016000400010>
51. Odunayo Joseph Olawuyi^{1*}, Segun Gbolagade Jonathan², Fikayo Ezekiel Babatunde³, Busayo Joshua Babalola², Ola Oluwa Simon Yaya⁴, James Oludare Agbolade⁵. et al. Accession x Treatment Interaction, Variability and Correlation Studies of Pepper (*Capsicum* spp.) under the Influence of Arbuscular Mycorrhiza Fungus (*Glomus clarum*) and Cow Dung. *American Journal of Plant Sciences.* 2014; 5: 683-690.
52. Nurhidayati N., Mariati N. Utilization of maize cob biochar and rice husk charcoal as soil amendments for improving acid soil fertility and productivity, *Journal of Degraded and Mining Lands Management.* 2014; 2(1): 223-230.
53. Zhang, C.; Lin, Y.; Tian, X.; Xu, Q.; Chen, Z.; Lin, W. Tobacco bacterial wilt suppression with biochar soil addition associates to improved soil physiochemical properties and increased rhizosphere bacteria abundance. *Appl. Soil Ecol.*2017; 112: 90–96 <https://www.cabdirect.org/cabdirect/abstract/20173070195>
54. Major J, Rondon M, Molina D, Riha SJ, Lehmann J. Maize yield and nutrition during 4 years after biochar application to a Colombian Savanna oxisol. *Plant and soil.* 2010; 333: 117-128. Link: <https://bit.ly/328NoNR>
55. Bedassa M. Soil acid Management using Biochar: Review. *Int J Agric Sc Food Technol.* 2020; 6(2): 211-217. DOI:[10.17352/2455-815X.000076](https://doi.org/10.17352/2455-815X.000076)
56. Van Zwieten, L., Kimber, S., Morris, S., Chan, K.Y., Downie, A., Rust, J., Joseph, S. and Cowie, A. Effects of Biochar from Slow Pyrolysis of Papermill Waste on Agronomic Performance and Soil Fertility. *Plant and Soil.* 2010; 327, 235-246. <https://doi.org/10.1007/s11104-009-0050-x>
57. Oguntunde, P.G., Fosu, M., Ajayi, A.E. and van de Giesen, N. (2004) Effects of Charcoal Production on Maize Yield, Chemical Properties and Texture of Soil. *Biology and Fertility of Soils*, 39, 295-299. <https://doi.org/10.1007/s00374-003-0707-1>
58. Chan, K.Y., Van Zwieten, L., Meszaros, I., Downie, A. and Joseph, S. (2007) Agronomic Values of Greenwaste Biochar as a Soil Amendment. *Australian Journal of Soil Research*, 45, 629-634. <https://doi.org/10.1071/SR07109>
59. [48] Rogovska, N., Laird, D.A., Rathke, S.J. and Karlen, D.L. (2014) Biochar Impact on Midwestern Mollisols and Maize Nutrient Availability. *Geoderma*, 230, 340-347. <https://doi.org/10.1016/j.geoderma.2014.04.009>
60. Gaskin, J.W., Steiner, C., Harris, K., Das, K.C. and Bibens, B. Effect of Low-Temperature Pyrolysis Conditions on Biochar for Agricultural Use. *Transactions of the Asabe*, 2008; 51, 2061-2069. <https://doi.org/10.13031/2013.25409>

61. Gundale, M.J. and DeLuca, T.H. Charcoal Effects on Soil Solution Chemistry and Growth of *Koeleria Macrantha* in the Ponderosa Pine/Douglas-Fir Ecosystem. *Biology and Fertility of Soils*, 2007; 43, 303-311. <https://doi.org/10.1007/s00374-006-0106-5>
62. Silber, A., Levkovitch, I. and Graber, E.R. pH-Dependent Mineral Release and Surface Properties of Cornstraw Biochar: Agronomic Implications. *Environmental Science & Technology*. 2010; 44, 9318-9323. <https://doi.org/10.1021/es101283d>
63. Sohi, S.P., Krull, E., Lopez-Capel, E. and Bol, R. A Review of Biochar and Its D. F. Hui DOI: 10.4236/as.2021.123014 224 *Agricultural Sciences Use and Function in Soil. Advances in Agronomy*. 2010; 105, 47-82. [https://doi.org/10.1016/S0065-2113\(10\)05002-9](https://doi.org/10.1016/S0065-2113(10)05002-9)
64. Lehmann, J., Rillig, M.C., Thies, J., Masiello, C.A., Hockaday, W.C. and Crowley, D. Biochar Effects on Soil Biota—A Review. *Soil Biology & Biochemistry*. 2011; 43, 1812-1836. <https://doi.org/10.1016/j.soilbio.2011.04.022>
65. Liang, B., Lehmann, J., Solomon, D., Sohi, S., Thies, J.E., Skjemstad, J.O., Luizao, F.J., Engelhard, M.H., Neves, E.G. and Wirick, S. Stability of Biomass-Derived Black Carbon in Soils. *Geochimica et Cosmochimica Acta*, 72. 2008; 6069-6078. <https://doi.org/10.1016/j.gca.2008.09.028>
66. Ding, Z.H., Wan, Y.S., Hu, X., Wang, S.S., Zimmerman, A.R. and Gao, B. Sorption of Lead and Methylene Blue onto Hickory Biochars from Different Pyrolysis Temperatures: Importance of Physicochemical Properties. *Journal of Industrial and Engineering Chemistry*, 37. 2016; 261-267. <https://doi.org/10.1016/j.jiec.2016.03.035>
67. Cayuela, M.L., Sanchez-Monedero, M.A., Roig, A., Hanley, K., Enders, A. and Lehmann, J. Biochar and Denitrification in Soils: When, How Much and Why D. F. Hui DOI: 10.4236/as.2021.123014 226 *Agricultural Sciences Does Biochar Reduce N₂O Emissions? Scientific Reports*, 3. 2013; Article No. 1732. <https://doi.org/10.1038/srep01732>
68. Muhammad A, Muhammad I, Muhammad R, Kawsar A, Kamran S, Izhar ul H, Shah F . Biochar improves phosphorus use efficiency of organic-inorganic fertilizers, maize-wheat productivity and soil quality in a low fertility alkaline soil. *Field Crop Res*. 2017; 214:25–37. <https://doi.org/10.1016/j.fcr.2017.08.01>
69. Gao S, Thomas HDL. Influence of biochar on soil nutrient transformations, nutrient leaching, and crop yield. *Adv Plants Agric Res*. 2016; 4(5):348-362. DOI:[10.15406/apar.2016.04.00150](https://doi.org/10.15406/apar.2016.04.00150)
70. Deluca TH, Mackenzie MD, Gundale MJ. Biochar effects on soil nutrient transformations. In Lehmann J. and Joseph S. (Eds.). *Biochar for environmental management: Science and Technology*. 2009; 251-270.
71. Griffith B. *Essential role of phosphorus in plants*. 2009.
72. Joseph, S. D., Campsarbertain, M., Lin, Y., Munroe, P., Chia, C. H., Hook, J., et al. An investigation into the reactions of biochar in soil. *Soil Res*. 48. 2010; 501–515. doi: 10.1614/IPSM-D-12-00058.1
73. Tian, S.; Tan, Z.; Kasiulienė, A.; Ai, P. Transformation mechanism of nutrient elements in the process of biochar preparation for returning biochar to soil. *Chin. J. Chem. Eng*. 2017; 25:477–486. <https://doi.org/10.1016/j.cjche.2016.09.009>
74. Liang, Biqing & Lehmann, Johannes & Solomon, Dawit & Kinyangi, James & Grossman, Julie & B, O'Neill & JO, Skjemstad & Thies, Janice & Luizão, Flávio & J, Petersen & Neves, Eduardo. Black Carbon Increases Cation Exchange Capacity in Soils. *Soil Science Society of America Journal*. 2006; 70:1719-1730. https://www.researchgate.net/publication/200736465_Black_Carbon_Increases_Cation_Exchange_Capacity_in_Soils
75. Beesley, L., E. Moreno-Jimenez, J.L. Gomez-Eyles, E. Harris, B. Robinson and T. Sizmur. A review of biochar's potential role in the remediation, revegetation and restoration of contaminated soils. *Environmental Pollution* 2011; 159: 3269-3282. doi:<http://dx.doi.org/10.1016/j.envpol.2011.07.023>

76. Liang, B., J. Lehmann, D. Solomon, J. Kinyangi, J. Grossman & B. O'Neill. Black Carbon increases cation exchange capacity in soils. *Soil Science Society of America Journal* 70: 2006; 1719-1730. doi:10.2136/sssaj2005.0383.
77. Cheng C-H, Lehmann J, Engelhard MH Natural oxidation of black carbon in soils: changes in molecular form and surface charge along a climosequence. *Geochim Cosmochim Acta*. 2008; 72:1598–1610
78. Diatta, A.A., Fike, J.H., Battaglia, M.L., Galbraith, J.M. & Baig, M.B. "Effects of biochar on soil fertility and crop productivity in arid regions: a review" *Arabian Journal of Geosciences*. 2020;13(14) DOI: [10.1007/s12517-020-05586-2](https://doi.org/10.1007/s12517-020-05586-2)
79. Laird, D., Fleming, P., Wang, B., Horton, R., & Karlen, D. Biochar impact on nutrient leaching from a Midwestern agricultural soil. *Geoderma*. 2010; 158(3-4): 436-442. doi:10.1016/j.geoderma.2010.05.01
80. Ndor, E.; Jayeoba, O.J.; Asadu, C.L.A. Effect of biochar soil amendment on soil properties and yield of sesame varieties in Lafia, Nigeria. *J. Exp. Agric. Int.* 2015;9:1–8 <http://dx.doi.org/10.9734/AJEA/2015/19637>
81. Silva, I.C.B., Basílio, J.J.N., Fernandes, L.A., Colen, F., Sampaio, R.A. and Frazão, L.A. Biochar from Different Residues on Soil Properties and Common Bean Production. *Scientia Agricola*, 2007; 74: 378-382. <http://dx.doi.org/10.1590/1678-992x-2016-0242>
82. Chen C.R., Phillips I.R., Condon L.M., Goloran J., Xu Z.H., Chan K.Y. Impacts of green waste biochar on ammonia volatilization from bauxite processing residue sand. *Plant and Soil*. 2013; 367: 301–312. <https://www.jstor.org/stable/42952895>
83. Edmunds, Charles Warren, "The Effects of Biochar Amendment to Soil on Bioenergy Crop Yield and Biomass Composition." Master's Thesis, University of Tennessee, 2012. https://trace.tennessee.edu/utk_gradthes/1150-