

Original Research Article

Studies on Relative Economics of Apple Nursery Production under Different Weed Management Practices in Kashmir Valley, India

ABSTRACT

Aims: The purpose of this study was to assess the relative economics of various weed management practices in apple nursery in Kashmir valley.

Study design: A Randomized Complete Block Design with seven treatments and three replications was employed as the experimental design.

Place and Duration of Study: The experiment was laid out at the Experimental Farm, Division of Fruit Science, Faculty of Horticulture, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar Campus, Srinagar, Jammu and Kashmir, India during the year 2020.

Methodology: Seven weed management treatments: manual weeding, pendimethalin @ 1 kg a.i. ha⁻¹, pendimethalin @ 1 kg a.i. ha⁻¹ + manual weeding, paddy straw mulch-6 cm thick, black polyethylene mulch-200 micron, weed-free and weedy check were tested in apple nursery (cv. Silver Spur grafted on M9-T337). The input costs and output in terms of monetary values were calculated under different treatments at prevailing market rates. The net return and benefit-cost (B:C) ratio was worked out based on input and output values under each treatment.

Results: The treatment paddy straw mulch showed the highest input cost (₹9543725.90) while black polyethylene mulch resulted the maximum output (17113252.80). The maximum net returns of ₹78,02,726 from grafted apple nursery plants were recorded under black polyethylene mulch with a benefit-cost ratio of 1.83 followed by pendimethalin @ 1 kg a.i. ha⁻¹ + manual weeding with net returns of ₹71,60,541 and benefit-cost ratio of 1.77.

Conclusion: Black polyethylene mulch resulted highest net return and B:C ratio, hence it found to be most profitable weed management practice in apple nursery. Pendimethalin @ 1 kg a.i. ha⁻¹ + manual weeding can also be considered as alternative weed management option.

Keywords: Apple Nursery; Benefit-cost ratio; Mulch; Pendimethalin; Weed management

1. INTRODUCTION

The cultivated apple (*Malus x domestica* Borkh.; Family Rosaceae) is world's leading temperate fruits. Globally, apple is grown over an area of 4.717 mha with annual production of 87.236 [1]. In India, it is predominantly cultivated in Jammu & Kashmir and Himachal Pradesh [2]. In Kashmir valley, most of the apple orchards are established in traditional system of planting with 5-6 m spacing. The high-density planting system (HDP) is increasingly being envisioned as an alternative production method with the potential to improve productivity, increase yield efficiency, reduce input costs, minimise hazards, and

maximise returns. Although, apple trees are generally propagated through seeds or by vegetative means but commercial propagation for high density planting are carried out by vegetative means i.e. budding and grafting on clonal rootstocks viz. M-9 and MM-106. The ultimate success in apple cultivation depends on the quality of the nursery plants used for establishing the orchards, it is imperative that the nursery plants should not only be true-to-type, but they should also be healthy, disease free and of good quality. The nursery plants require adequate care and management for one or two years before their transplantation to permanent orchard sites.

Apple nursery plants are very often prone to weed infestation particularly in their early growth period. Weeds cause significant losses by competing with the primary crop for water and nutrients, as well as providing a possible breeding habitat for numerous insects, pests, and diseases due to the shallow root systems of nursery plants. Due to high competition for water, light, nutrients and interference with other operational factors among plants and weeds; inadequate weeds management lead to poor plant growth and development [3]. Weed management in fruit nurseries is normally achieved by a variety of methods around the world, either mechanically through specific cultivation practices or with the application of herbicides; however, the traditional hand weeding approach is the most common in India, particularly in the Kashmir.

In crop production systems, mulching plays an important role in the conservation of soil moisture during dry periods [4] in addition to providing benefits such as weed suppression [5], reducing water runoff and soil erosion, improving water and fertilizer use efficiency [6] and improve the aesthetic value of landscapes and economic value of crops [7]. Organic mulches are particularly appealing because they provide a means of improving not only the physical properties of soil but also its nutrient content [8-10]. Organic mulches include straw, coconut coir, grasses, and cover crop cuttings, all of which provide nutrients to the soil as they decompose. Organic mulches are environmentally friendly and that can be easily applied to orchards and nurseries. Among inorganic mulches, black polyethylene mulch is widely used in various crop production systems as it has greater efficiency in weed control and [11]. Because of their operational benefits, inorganic plastic mulch films are commonly utilized in specialty crop production systems [12]. It also results in greater temperature regulation of soil [13].

Nevertheless, manual weed control is not only laborious but is also highly expensive. Thus, of late, manual and mechanical weed control methods are gradually being replaced by other alternatives such as the use of mulches and herbicides as these are easier, cheaper and less time consuming. Thus meaningful analysis of economics of various weed management strategies is essential so as to analyse whether the returns exceed the cost or not. Therefore present study was carried out with the objective find out the relative economics of different weed management treatments in apple nursery production in Kashmir valley.

2. MATERIAL AND METHODS

2.1 Experimental site

The experiment was conducted in the Experimental Field of the Division of Fruit Science, Sher-e-Kashmir University of Agricultural Science & Technology of Kashmir (SKUAST-K), Shalimar Campus, Srinagar, Jammu and Kashmir (India) during the year 2020. The experimental site is located at 34.1° North latitude and 74.9° East longitude and 1587 m above the mean sea level.

2.2 Plant Materials

Apple cv. Silver Spur grafted on M-9 T337 rootstock was used as plant materials for assessing the weed management trial in apple nursery in present study. Apple cv. Silver Spur is a compact to medium tree vigour with excellent spur density, medium and large size conical fruits with pronounced lobes, deep red skin with stripes, mature in the middle of the season recommended for cultivation in Jammu & Kashmir, Himachal Pradesh, Arunachal Pradesh, Uttarakhand [14]. The M-9 T337 is a dwarfing type clonal rootstock introduced in Kashmir from Holland and being used for production of grafted planting materials of apple for high density planting. Morphologically M9 T337 rootstock is of weak vigour with short internodes, reddish brown in colour (sunny side), strong undulated leaf margin, weak prudence on lower side of leaf, long petiole length with small stipule [15].

2.3 Weed control treatments

The treatments included: manual weeding (T_1), pendimethalin (T_2), pendimethalin + manual weeding (T_3), paddy straw mulch - 6cm thick (T_4), black polyethylene mulch (T_5), weed free (T_6) and weedy check (T_7). The paddy straw mulch (6 cm thick) and black polyethylene mulch (200 micron) were applied around the plants in each treatment plot to completely cover the soil with the mulch. On March 15, pendimethalin @ 1 kg a.i. Ha^{-1}) was applied as a pre-emergence herbicide in respective treatment plots.

2.4 Experimental Design

The experiment was laid out with seven weed management treatments in Randomized complete block design where each treatment was replicated thrice.

2.5 Data collection

The Cost concepts used to work out the nursery production economics were the variable cost, fixed cost and total cost of cultivation. Cost 'A' includes the costs on account of hired human labour, manures and fertilizers, rootstock, scion wood, machinery charges, irrigation charges, plant protection charges, land revenue, depreciation and repairs, interest on working capital etc. Cost 'B' includes rental value of land and interest on fixed capital cost which is added to the Cost 'A'. Cost 'C' is the total cost of production, which included all the item's costs.

2.5.1 Cost of production

Cost of production was computed on the basis of costs of all inputs for production of grafted apple nursery under different treatments. It includes fixed cost, variable costs, depreciations, rental value of land, interest on working capital.

2.5.2 Gross return

Gross return from survived grafted plant materials was estimated on the basis of the sale price of grafted plant materials (₹200 per plant). For estimating net return, total cost of production was subtracted from gross return as follows:

$$\text{Net return} = \text{Gross return} - \text{Cost of production}$$

2.5.3 Benefit Cost Ratio:

Benefit: cost ratio (B:C) is defined as the amount received in the shape of profit on the costs of one rupee.

The BCR was computed by using following formula:

$$\text{Benefit: cost (B: C) ratio} = \frac{\text{Gross return}}{\text{Cost of production}}$$

2.6 Data analysis

The data recorded on various parameters were statistically analyzed at a 5% significance level according to Panse and Sukhatme's standard method [16].

3. RESULTS AND DISCUSSION

The fixed input cost irrespective of the treatments incurred presented in Table 1. It was ₹ 56307.20 on account of rental value of land, depreciation on implements, and interests on fixed capital.

Table 1: Fixed cost for production of grafted apple nursery for all the weed management treatments (per ha basis)

S.No.	Particulars	Cost (₹)
1.	Rental value of land	50000
2.	Depreciation on implements	3120
	Sub-total	53120
3.	Interest on fixed capital (6 %)	3187.20
	Total	56307.20

Common variable cost (other than treatment and its application costs) irrespective of treatments was calculated ₹8645895 (Table 2). It includes hired human labour, machinery charges, manures and fertilizers, rootstock, scion wood, irrigation charges, plant protection charges, land revenue, depreciation and repairs, interest on working capital etc. Input cost due to treatments varied from highest in weed free conditions to the lowest in weedy check (Table 2). The rental value of land was found to be the maximum part of the common fixed cost. Because of the limited land availability for agricultural use in the Kashmir valley and apple cultivation being most remunerative crop, the rental value of lands also comparatively on higher side.

Data in Table 3 reveals that the highest cost of production (₹ 9543725.90) was incurred under paddy straw mulch, followed by weed free (₹9392675.90), and black polyethylene mulch (₹9310525.90). Treatment pendimethalin @ 1 kg a.i. ha⁻¹ + manual weeding exhibited higher cost of cultivation (₹9261381.64) compared to manual weeding and pendimethalin @ 1 kg a.i. ha⁻¹ as the cost of production estimated under manual weeding and pendimethalin @ 1 kg a.i. ha⁻¹ was ₹9259115.90 and ₹9223221.64, respectively. The lowest cost of production (₹9220955.90) was estimated in weedy check. Higher cost of production under paddy straw mulched is mainly attributed to the facts that the unit cost of the paddy straw as mulching material is comparatively high than that of other mulch materials. In Kashmir valley, paddy straw also used as animal feed thereby the cost of the paddy straw is higher.

The highest gross return (₹17113252.80) was estimated under black polyethylene mulch, followed by paddy straw mulch (₹16639143.92), weed free (₹16591921.92). Pendimethalin @ 1 kg a.i. ha⁻¹ + manual weeding treatment exhibited higher gross return (₹16421922.72)

Table 2: Variable cost for production of grafted apple nursery cv. Silver Spur on M9-T337 under different weed management treatments (per ha basis)

Treatment	Common variable input cost (₹)	Cost due to weed management treatments (₹)			Interest working capital (6%) (₹)	Total variable cost (₹)
		Labour	Herbicide	Mulch		
T ₁ : Manual weeding	8645895	36000	-	-	520913.70	9202808.70
T ₂ : Pendimethalin @ 1 kg a.i. ha ⁻¹	8645895	-	2137.49	-	518881.95	9166914.44
T ₃ : Pendimethalin @ 1 kg a.i. ha ⁻¹ + manual weeding	8645895	36000	2137.49	-	521041.95	9205074.44
T ₄ : Paddy straw mulch (6 cm thick)	8645895	4500	-	300000	537023.70	9487418.70
T ₅ : Black polyethylene mulch (200 micron)	8645895	4500	-	80000	523823.70	9254218.70
T ₆ : Weed free	8645895	162000	-	-	528473.70	9336368.70
T ₇ : Weedy check	8645895	-	-	-	518753.70	9164648.70

Table 3: Economics of grafted apple nursery cv. Silver Spur on M9-T337 rootstock under different weed management practices (per ha basis)

Treatment	Cost of production (₹)	Gross return(₹)	Net return (₹)	B:C ratio
T ₁ : Manual weeding	9259115.90	15898702.96	6639587.06	1.72
T ₂ : Pendimethalin @ 1 kg a.i. ha ⁻¹	9223221.64	11112281.04	1889059.40	1.20
T ₃ : Pendimethalin @ 1 kg a.i. ha ⁻¹ + manual weeding	9261381.64	16421922.72	7160541.08	1.77
T ₄ : Paddy straw mulch (6 cm thick)	9543725.90	16639143.92	7095418.02	1.74
T ₅ : Black polyethylene mulch (200 micron)	9310525.90	17113252.80	7802726.90	1.83
T ₆ : Weed free	9392675.90	16591921.92	7199246.02	1.77
T ₇ : Weedy check	9220955.90	9051512.96	(-)169442.94	0.98

compared to manual weeding and pendimethalin @ 1 kg a.i. ha⁻¹ as the gross return estimated under manual weeding and pendimethalin @1 kg a.i. ha⁻¹ was ₹15898702.96 and ₹11112281.04, respectively (Table 3). The lowest cost of production (₹9051512.96) was estimated in weedy check. The lowest cost of production (₹9051512.96) was estimated in weedy check. The highest gross return under black polyethylene mulch is attributed to the facts that black polyethylene mulch results in greater proportion of salable plants compared to other treatments.

Black polyethylene mulch resulted the highest net return (₹7802726.90) was estimated under followed by weed free (₹7199246.02) and pendimethalin @ 1 kg a.i. ha⁻¹ + manual weeding (₹7160541.08) (Table 3). Treatment paddy straw mulch exhibited higher net return (₹7199246.02) compared to manual weeding and pendimethalin @1 kg a.i. ha⁻¹ as the net return estimated under manual weeding and pendimethalin @1 kg a.i. ha⁻¹ was ₹6639587.06 and ₹ 1889059.40, respectively. There was loss of ₹169442.94 under weedy check as net return was less than the total cost of production. As the net return is obtained by deducting cost of production from gross return, the proportionate outcome was higher in black polyethylene much compared to other treatments.

It is evident from data in Table 3, that the highest B:C ratio (1.83) was recorded under black polyethylene mulch, by weed free (1.77) and Pendimethalin @ 1 kg a.i. ha⁻¹ + manual weeding (1.77). Treatment paddy straw mulch exhibited higher B:C ratio (1.74) compared to manual weeding and pendimethalin @ 1 kg a.i. ha⁻¹ as the gross return estimated under manual weeding and pendimethalin @ 1 kg a.i. ha⁻¹ was 1.72 and 1.20, respectively. The lowest B:C ratio (0.98) was estimated in weedy check.

In present study, black polyethylene mulch emerged the most beneficial treatment in terms of monetary benefits as it yielded highest gross return because of higher graft survival that exerted to highest net return and thus constituted the highest Benefit: cost ratio compared to other treatments. Meena [17] advocated the use grass mulch followed by glyphosate sprays at 0.8 kg ha⁻¹ to be cost-effective strategy for weed management in peach. Hegazi [18] advocated that plastic mulching control of weeds and most importantly to save water and increase the net income from grapevines in comparison of mulching, straw mulching, and herbicides. Sharma and Kathiravan [19] recorded pine needle as the best mulch material for plum as it provided maximum income compared to others mulches viz. transparent polythene, black polythene, bicoloured polythene, field grass, pine needles and un-mulched control. In Aonla (*Emblica officinalis* Gaertn.), Iqbal *et al.* [20] while studying the economics of different mulching materials viz., black polythene, white polythene, paddy straw, saw dust, sarkanda, dry grass and control (unmulched) on aonla, they recorded higher cost of cultivation due to black polythene mulch but net return and B:C ratio found to be maximum in black polythene mulch (2.04:1) and minimum in control (1.69:1). Rakesh *et al.* [21] recorded grass mulching as most cost-effective treatment in plum (cv. Red Beut) in comparison of herbicides, mulching and hand weeding treatments.

4. CONCLUSION

The treatment paddy straw mulch incurred the highest input cost (₹ 9543725.90) while black polyethylene mulch resulted in maximum output (₹17113252.80). The maximum net returns of ₹78,02,726 from grafted apple nursery plants was recorded under black polyethylene mulch with a benefit: cost ratio of 1.83; followed by pendimethalin @ 1 kg a.i. ha⁻¹ + manual weeding with net returns of ₹71,60,541 and benefit cost ratio of 1.77. Hence, black polyethylene mulch found to be most profitable weed management practice in apple nursery in Kashmir valley. Pendimethalin @ 1 kg a.i. ha⁻¹ + manual weeding can also be considered as alternative weed management option.

ACKNOWLEDGEMENTS

Authors are highly thankful to Head, Division of Fruit Science, Sher-e-Kashmir University of Agricultural Sciences and technology of Kashmir, Shalimar Campus, Srinagar, Jammu and Kashmir, India for providing necessary facilities for carrying out the this research.

AUTHORS' CONTRIBUTIONS

Authors AUH, AA, AK FUK, SAM and BAA designed the study. Authors AUH and AA managed the literature searches. Authors AUH, AA and AK managed the experimental field. Authors AUH, FMM and FF wrote the first draft of the manuscript. All authors read and approved the final manuscript.

CONSENT

Not Applicable.

ETHICAL APPROVAL

Not Applicable.

COMPETING INTERESTS:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. FAOSTAT. Area and Production of apple, 2019. Food and Agriculture Organization, Rome, Italy. Available online: <http://www.fao.org/faostat/en/#data/QI>.
2. NHB, Indian Horticulture Database. National Horticulture Board, Ministry of Agriculture, Gurugram, India; 2019. Available online: <http://www.nhb.gov.in/>
3. Leela D. Present status and future scenario of weed control in horticultural crops. In: Proceedings of Golden Jubilee Symposium on Horticultural Research - Changing *Scenario*, Bangalore, India; 1993.
4. Hira GS, Jalota SK, Arora, VK. Efficient management of water resources for sustainable cropping in Punjab. Technical Bulletin, Department of Soils, Punjab Agricultural University, India; 2004.
5. Kaur K, Kaundal GS. Efficacy of herbicides, mulching and sod cover on control of weeds in plum orchards. Indian J. Weed Sci. 2009;41:110-112.
6. Sharma R, Bhardwaj. Effect of mulching on soil and water conservation - A review. Agric. Rev. 2017;38(4):311-315. DOI: <https://doi.org/10.18805/ag.R-1732>.
7. Iqbal R, Raza MAS, Valipour MB, Saleem MF, Zaheer MS, Ahmad A, Toleikiené M, Haider I, Aslam MU, Nazar MA. Potential agricultural and environmental benefits of mulches-a review. Bull. Natl. Res. Cent. 2020;44:75. DOI: <https://doi.org/10.1186/s42269-020-00290-3>.

8. Dahiya R, Ingwersen J Streck T. The effect of mulching and tillage on the water and temperature regimes of a loess soil: experimental findings and modelling. *Soil Tillage Res.* 2007;96:52-63. DOI: <https://doi.org/10.1016/j.still.2007.02.004>.
9. Chalker-Scott L. Impact of mulches on landscape plants and the environment - a review. *J. Environ. Hort.* 2007;25:239-249. DOI: <https://doi.org/10.24266/0738-2898-25.4.239>.
10. Jordan A, Zawala LM, Gill J. Effects of mulching on soil physical properties and runoff under semi-arid conditions in southern Spain. *Catena* 2010;81(1):77-85. DOI: <https://doi.org/10.1016/j.catena.2010.01.007>.
11. Moreno MM, Moreno A. Effect of different biodegradable and polythene mulches on soil properties and production in a tomato crop. *Scientia Hort.* 2008;116:256-63. DOI: <https://doi.org/10.1016/j.scienta.2008.01.007>
12. Espi E, Salmeron A, Fontecha A, Garcia Y, Real AI. Plasticfilms for agricultural applications. *J. Plastic Film Sheeting.* 2006;22:85-102. DOI: <https://doi.org/10.1177/8756087906064220>.
13. Kader MA, Senge M, Mojid MA, Ito K. Recent advances in mulching materials and methods for modifying soil environment. *Soil Tillage Res.* 2017;168:155-166. DOI: <https://doi.org/10.1016/j.still.2017.01.001>.
14. Singh G. Checklist of commercial varieties of fruits. Department of Agriculture and Co-operation Ministry of Agriculture, India: New Delhi; 2012.
15. Ullah SS, Bhat ZA, Bhat KM, Shikari AB, Khan I, Khan FA. Morphological characterization of clonal rootstocks of apple. *Intl. J. Chem. Studies.* 2018;6(4):1584-1588.
16. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers.* Indian Council of Agricultural Research Publication, India: New Delhi; 1985.
17. Meena MK, Sharma DD, Meena OP. Effect of different weed management practices on weed population, yield potential and nutrient status of peach cv. July Elberta. *Res. Crops.* 2015;16(3):519-525. DOI: <https://doi.org/10.5958/2348-7542.2015.00073.X>.
18. Hegazi A. Plastic mulching for weed control and water economy in vineyards. *Acta Hort.* 2000;536:245-250. DOI: <https://doi.org/10.17660/ActaHortic.2000.536.28>.
19. Sharma JC, Kathiravan G. Effect of mulches on soil hydrothermal regimes and growth of plum in mid hill region of Himachal Pradesh. *Indian J. Hort.* 2009;66(4):465-471.
20. Iqbal M, Bakshi P, Wali VK, Hamid N. Economics of different mulching materials on aonla (*Emblica officinalis* gaertn.) under rainfed conditions of Jammu. *Intl. Res. J. Agric. Eco. Stat.* 2017;8(2):418-421. <https://doi.org/10.15740/HAS/IRJAES/8.2/418-421>.
21. Rakesh K, Sharma DD, Gopal S, Bhawna K, Kuchay MA. Effect of different weed management practices on soil properties and soil and leaf nutrient content of plum (*Prunus salicina* L) cv Red Beaut. *Intl. J. Farm Sci.* 2020;10(3-4):93-97. DOI: <https://doi.org/10.5958/2250-0499.2020.00059.2>.