

### **A review paper on agro-food waste and food by-product valorization into value added products for application in the food industry: Opportunities and challenges for Cameroon bioeconomy.**

#### **ABSTRACT**

Agricultural production, agro-industrial food processing, distribution and consumption generate high amount of varied food by-products and waste which place a heavy burden on the environment and cause losses to the food industry. Food waste is commonly disposed of in landfills or incinerated, causing many environmental, social, and economic issues. However, many of these by-products and wastes have been reported to have at times a nutritional or functional properties higher than the final product, making them potential raw materials for application in the agro-food industry. This along with the recent sustainable development goals of food security, environmental protection, and energy efficiency are the key drivers for food waste valorization. Valorization of food waste within the bio-economy approach offers an economical and environmental opportunity, which can reduce the problems of its conventional disposal. Traditionally, in Africa, especially in Cameroon, food by-products and waste have been valorized into a range of product for application in food and food preparation, including, food additives and spices, food emulsifiers and stabilizers, food salts and nutraceuticals. Traditional Waste valorization methods could achieve sustainable development in technologically underdeveloped countries, methods that could not only improve agro-food waste management but could also lead to the production of industrially important biochemical, food ingredients and food products, in essence, valuable added products from waste. In addition, the processing and conversion of these agro-food by-products and waste generated in the poor regions of the world for the production and formulation of novel foods and biochemicals will directly benefit the local communities by reducing environmental pollution and increasing income in the food industry. The aim of this review is to provide an insight on the latest trends in food waste valorization using traditional methods in an Africa country such as Cameroon. This paper presents the variety and volume of food waste within the food chain and types of food waste feedstock's that valorized into various products using traditional methods. Furthermore, a series of examples of key food waste valorization schemes and value added products as case studies to demonstrate the advancement in traditional bioconversions are described, bringing out the opportunities and challenges for the Cameroon bioeconomy.

**Key Words:** Food waste, pollution, feed-stock, valorization, Value-added products, Bioeconomy

## 1. INTRODUCTION

The ever increasing demand for food coupled with higher environmental standards is reshaping agricultural activities toward ecologically sustainable and efficient systems (Ominski *et al.*, 2021). While the existing literature has mainly focused on increasing food production, the magnitude of waste and by-products are too large to be ignored (Irfanoglu *et al.*, 2014, FAO, 2019b). Agricultural production and agro-industrial processing generate high amounts of food by-products and waste, which have been reported to have at times a nutritional or functional content higher than the final product (Ayala *et al.*, 2011). Roughly, one-third of the edible parts of global food produced for human consumption is estimated to be lost or wasted (FAO, 2015), and these losses have been valued at 1 trillion USD (FAO, 2013). Fruit and food waste is also generated by damage during transportation, storage, and processing. To manage the nutritional problems of today's society, we require more composite nutritional sources. Food wastes and byproducts are of paramount importance here due to the presence of sufficient quantities of beneficial nutrients and bioactive compounds (Torres-Leon *et al.*, 2018). Many waste valorization methods could achieve sustainable development, methods that could not only improve agro-food waste management but could also lead to the production of industrially important bio-chemicals, food ingredients and food products, in essence, valuable end products from waste.

Food by products are generally secondary products derived from primary agro-food production processes, and they represent an interesting and cheaper source of potential functional ingredients (Faustino *et al.*, 2019). On the other hand, food waste is “unwanted or unusable material, substances, or by-products” that are “eliminated or discarded as no longer useful or required after the completion of a process” (Markus *et al.*, 2019). Food by-products and waste remain a global dilemma, with negative environmental, social, and economic consequences (Ominski *et al.*, 2021) associated with estimated annual losses of approximately one-third of all edible food (1.3 billion metric tonnes, MMT) across the supply chain (FAO, 2011a). Many of these biomaterials are not utilized and end up in municipal landfills where they create serious environmental problems due to microbial decomposition and leachate production. In some cases, the byproducts are burned to remove fungi and parasites. From the economic point of view, the adverse impact is due to the costs related to the handling of solid waste in land fills. Moreover, the management of large amounts of different degradable materials poses a challenge (Novoa-Muñoz *et al.*, 2008; Pateiro-Moure *et al.*, 2009). Social impacts may be attributed to an ethical and moral dimension within the general concept of global food security since 805 million people across the globe suffer from hunger (FAO, 2014). Most of the studies on food waste and by-products in sub-Saharan Africa, particularly in Cameroon concerned with estimating the extent of waste and by-product generation, without, however, worries of the technological importance and potential on development of bio-economy.

Traditionally, food waste is usually incinerated or disposed of in landfills leading to subsequent air/water pollution, and soil/food contamination. To decrease these problems, the European Union (EU) is promoting the reduction of food wastes, and the search for new end-uses for food by-products (Wunder *et al.*, 2018). Following these recommendations, food waste from several agro-food industries (vegetables, fruits, beverages, sugar, meat, aquaculture and marine products, seafood, etc.) represent an interesting and cheaper source of potentially functional or bioactive compounds. In this context, the biorefinery concept has emerged as the development of integrated and combined extraction or recovery processes for the conversion of biomass into many added-value products (Mazzutti, 2020).

Food waste is abundant in sub-Saharan Africa, especially in Cameroon, and previous studies on waste management in Cameroon have focused on waste collection and disposal practices and their environmental implications (Vermande and Ngnikam, 1994; Ngnikam, 2000) with little consideration on the valorization into value added products and biochemical. In Cameroon for example, food crops such as maize (*Zea mays*), rice (*Oryza sativa*), cassava (*Manihot esculenta Crantz*) and some fruits like oranges, lemons and pineapple are widely cultivated for subsistence alongside vegetables such as amaranth (*Amaranthus cruentus*) (Njukwe *et al.*, 2014; Tata *et al.*, 2016). Numerous reports (Lipinski *et al.*, 2013; Foresight, 2011; FAO, 2011a; Stuart, 2009) have underlined the significance of food by-products and waste and the need to reduce them to improve food security and sustainability of food systems.

The available disposal options in under-developed technological poor countries like Cameroon still largely remain slash and burn or land fillings, producing large quantities of greenhouse gases (GHG). Incineration and composting also produce greenhouse gases, and wastewater from anaerobic digestion causes eutrophication and acidification of local ecosystems (Salemdeeb and Al-Tabbaa, 2015; Whiting and Azapagic, 2014). The processing and conversion of agro-food by-products and waste generated in the poor regions of the world for the production and formulation of novel foods and biochemicals will directly benefit the local communities. In Cameroon, for example, the use of Plantain and Banana byproducts has been suggested as the main raw material for the production of traditional food salts, commonly called Nikkihused in yellow achu-soup preparation due to emulsification properties (Ngwasiriet *et al.*, 2021).

The valorization of residues and byproducts generated in the poor regions of the world for the formulation of novel foods will directly benefit the local communities (Luque and Clark, 2013). The valorization of agro-food byproducts and waste into different high-value products has been reviewed by numerous studies, (Maina *et al.*, 2017; Mirabella *et al.* 2014; Lin *et al.*, 2013; Tuck *et al.*, 2012) while others (Xiong *et al.*, 2019; Serna-Loaiza *et al.*, 2019; Fritsch *et al.*, 2017; Pleissner *et al.*, 2016) investigated and described the chemical processes involved in the conversion. However, these reviews focused on valorization using advanced technology, affordable to developed and industrialized countries, with little focus on the contribution of agro-food waste valorization for food security in undeveloped African countries. In this review paper we will therefore try to frame the key issues associated with traditional valorization of agro-food waste into value added food products and biochemicals within the context of the emerging bio-economy and circular-economy, suggesting that valorizing these agro-food wastes can significantly contribute to solving malnutrition and hunger in undeveloped African countries, especially in Cameroon. The following examples have been chosen as being a contrasting set of materials for which much information is available and for which clear improvements and opportunities are envisaged.

## **2. AGRICULTURAL BY-PRODUCTS AND WASTE GENERATION IN CAMEROON**

Cameroon is located in the Gulf of Guinea between Gabon, Guinea Republic and Nigeria with 26,55 million inhabitants as of 2020. The importance of agriculture, fishery and livestock and their contribution to the economy of Cameroon is significant as it is highly dependent on it, which provides 70 percent of its active population employment opportunity. Due to the high

agricultural activities, both plantation and non-plantation waste biomass are abundant, which are highly under-utilized. In Cameroon for example, plantation agriculture under the CDC (the second largest employer after the government), Pamol and, SOCAPALM, among others, are legacies of European administration. Due to its abundant natural resources, Cameroon stands as a major global producer of goods like cocoa, coffee, bananas, palm products, tobacco, rubber and cotton. Other important products include: cereals (maize, millet and sorghum, paddy, etc.), roots and tubers (cassava, cocoyam, taro, potato, yam, etc.), oilseeds such as groundnut, cottonseed, etc., fruits and vegetables, including citrus, pineapple, tropical fruits, legumes and pulses, spices and condiments, leafy vegetables and mushrooms, plants and ornamental flowers, etc (Benoit Daviron *et al.*, 2004).

The agricultural sector is one of the main sectors generating the largest quantities of waste and by-products, if allowed to accumulate indiscriminately will constitute nuisance to global health and threat to food security (European Commission, 2015, Bracco *et al.*, 2018). In Cameroon for example, food crops such as maize (*Zea mays*), rice (*Oryza sativa*) and cassava (*Manihot esculenta* Crantz) are widely cultivated for subsistence alongside vegetables such as amaranth (*Amaranthus cruentus*) (Njukwe *et al.* 2014; Tata *et al.* 2016). These food crops generate wastes such as corncob, rice husk and groundnut husk, for example, Kouame *et al.* (2013) reported a 10% and 20% rates of postharvest loss of amaranths and black nightshades, respectively. These waste are not used as fuel because of their low heating value and the volume of smoke produced (Kung *et al.* 2015; Kumer *et al.* 2015). Low bulk density and slow decomposition also limit their use in agriculture as a soil amendment (Steiner *et al.* 2010; Enders *et al.* 2012) whereas they contain appreciable quantities nutrients, bioactive compounds and phytochemicals which could offer both nutrition and health benefits.

Cameroon, as in the case of most sub-Saharan African countries, is currently facing one of its most serious food crises. (World Food Program, 2017). The number of food-insecure households in Cameroon is estimated at around 16% (about 3.9 million people), of which 1% are in a situation of severe food insecurity. There are many reasons for this. While food unavailability and inaccessibility are incriminated, food wastage is rampant. Estimated annual food quantitative losses and waste in the supply chain represent approximately 40 to 50% of the world's fresh products—30% for cereals and 20% for oilseeds, meat, and milk products. (Gustavsson *et al.* 2011) Food losses and waste refer to the decrease in quantity or quality of food along the food supply chain (FAO, 2019). In the case of perishable foods such as fruits, waste also occurs along the post-harvest chain at the retail level due to a wide variety of factors. Microorganisms and residues of pesticides and other toxic chemicals or organic inputs that are used to improve crop production can also accelerate the process of post-harvest losses and lead to population health problems (Goletti, F. and Wolff, C. 1999, Blanke *et al.* 2015). It is recognized that toxic exposures. may worsen the micronutrient status, e.g., by increasing the nutritional requirements; vice versa, imbalanced diets and micronutrients deficiencies may increase the vulnerability to the effects of toxic substances and alter body defense systems (Frazzoli *et al.* 2020). Estimates of postharvest losses of fruits and vegetables vary considerably in developed and developing countries (Kader 2005). In some African countries, about 30% of products are lost, and this figure can rise to 50% for very perishable foods, such as fruits and vegetables. 48.5% postharvest loss was reported in Nigeria (Kughur *et al.* 2015). 45.9% postharvest banana loss reported in Ethiopia, of which about 15.7%, 22.1%, and 8.1% were incurred at the farm, wholesale, and retailer levels, respectively (Zenebe *et al.* 2015). This rate of post-harvest losses of fruits and vegetables varies from 20 to 40% in Bangladesh (Wills *et al.*

2004). In Cameroon, 10% and 20% rates of postharvest loss of amaranths and black nightshades was reported, respectively (Kouame *et al.* 2013). Most of these studies were concerned with estimating the extent of losses without, however, worries of the causes and consequences on the health of populations.

Though Cameroon coffee production fluctuated substantially in recent years, the total production in 2020 was 36,207 tonnes (ICO, 2021). One of the main concerns in relation to the sustainability of the coffee production chain involves the recovery of a large amount of its generated by-products (Kasuya *et al.*, 2015; Oliveira and Franca, 2015). The residues and the by-products of coffee processing constitute around 60% of the wet weight of the fresh fruit, represent a relevant source of pollution and environmental threat (Rathinavelu *et al.*, 2005; Franca and Oliveira, 2009; Hughes *et al.*, 2014). Among the solid residues, the peel is the first to be generated. It represents 39% of fresh weight or 29% of fruit's dry matter, while the parchment represents around 12% of the fruit (Matos, 2015). The production statistics from Cameroon clearly indicates availability of large amount of coffee peels and parchment, which are highly underutilized.

The production of plantain and banana, which is an essential component in the diet of West and Central African populations (Folefack *et al.* 2017), is an environmental challenge due to the large volume of waste and by-products generated. Cameroon, which is the fourth largest plantain and banana producer in West and Central Africa, does not escape this situation (Folefack *et al.*, 2017) with about 1.6 million tonnes produced annually (Lescot, 2017). Plantain and Banana peel waste are the solid waste derived from the processing of plantain and banana and, it constitutes about 40% of fresh fruit weight. Bananas contain 60% pulp and 40% peel, 7.25 kg of peel produced from a banana box of 18.14 kg (Sharma *et al.*, 2016). However, the shell contains carbon-rich organic compounds such as cellulose (7.6–9.6%), hemicellulose (6.4–9.4%), pectin (10–21%), lignin (6–12%), chlorophyll pigments and some other low molecular weight compounds. If not treated properly, these wastes create an annoying odor due to the natural decomposition and produce gases that contribute to the greenhouse effect (Tibolla *et al.*, 2018).

### **3. CURRENT USES AND VALORIZATION OF AGRO-FOOD WASTE AND Food BY-PRODUCTS IN FOOD PREPARATION IN AFRICA AND CAMEROON**

Most of studies on waste in Cameroon concerned with estimating the extent of food waste, by-products and losses and their disposal without, however, worries of sustainable valorization of these waste into value added products. It is important that these wastes are identified and properly valorized in order to reduce losses of valuable nutrients and potential raw materials, and protect the dwellers in the community as well as the environment because waste is directly linked to human development, both technologically and socially (Tobias *et al.*, 2014).

#### **3.1. Value added food products from plantain and banana wastes and by-products**

Plantain (*Musaparadisiaca*) is a large herbaceous perennial crop belonging to the family Musaceae. The family Musaceae also includes bananas (*Musa sapientum* and *Musa cavandish*). Plantain (*Musa* spp) is one of the world's most important fruit. This plant widely cultivated in tropical countries for its valuable applications in food industry. Bananas and plantains have been said to be the 2<sup>th</sup> largest fruit crop of the world (FAO 2014). The peel of banana represents about 30-40% of the total weight of fresh banana and has been underutilized. The world production is estimated to be 139 million tons, in which tropical Africa alone produces about 17 million tons of bananas annually. It has become a basic food crop for over 70 million people in Africa. Over 50 species of *Musa* are in existence, in which the main groups of edible bananas or plantains are

derived from *Musa acuminata* and *Musa balbisiana* (Onyegbado et al, 2004). Plantains and bananas belong to the same family and have the same method of growth and development, but can be differentiated from one another by form of stem, colour and size, leaf colour, fruit (finger) shape and size, composition of nutrients in fruits (food and peel), and the composition of plantains is mostly carbohydrates while in bananas, it is sugar (Ogidi *et al.*, 2018). The parts of the banana and plantain plants which are generated as byproducts after cultivation and processing include the peels, leaves, sheaths, pseudostem, pith and male bud (Mohapatra *et al.*, 2010).

### **3.1.1 Traditional food salts (Nikkih) from Banana and plantain peel**

Plantain and banana peels are valorized using traditional **technology for** the production of a traditional food salt or potash, commonly called “nikkih” in many regions of Cameroon, especially the West and North West Regions. This traditional food salt (nikkih) is the crude brownish or blackish extract produced traditionally by leaching the ashes of combusted agro-food waste with water to obtain a potassium-carbonate-rich crude bio-extract (Oyegbado et al. 2004; Adewuyi et al. 2008). Nikkih can either be obtained from peels boiled at 90°C before drying, or directly dried raw peels which are combusted to produce ash that is leached with water to obtain the crude blackish or brownish extract (Ngwasiri *et al.*, 2021). Their chemical composition shows that they are a mixture of salts and thus, made of cations and anions, the major cation being generally sodium or potassium whereas the major anions are generally carbonates, bicarbonates, sulfates, and chlorides (Saidou et al., 2015; Osano et al., 2013). These bio-base functional plant extracts are fast replacing the common lake salt called ‘kangwa’, as they are regarded as cheaper, safer, less-toxic and readily available from food waste biomass and their production from waste biomass contributes to environmental protection.

Nikkih is now used in the preparation of a variety of foods due to its functional properties since it serves as emulsifier, tenderizer, thickener, seasoning, potentiating adjunct and preservative (Ajiboye et al., 2013). The functionalities of nikkih have been attributed to the alkalinity of the aqueous solution (Onwuka et al., 2003). With respect to its functionality, the ability to reduce cooking time has been studied (Bergeson et al., 2018; Doumta et al., 2012). It is an important household ingredient which is traditionally used in many food preparation processes in Cameroon, and also in food industries. It is used in the preparation of yellow “achu” soup, and gives vegetables a good texture and appearance during their preparation. In the dairy industry, when cream is separated from whole milk during the production of butter, lime water is often added to the cream to reduce acidity prior to pasteurization.

### **3.1.2 Flour from banana and plantain pseudostem**

The banana pseudostem is particularly rich in cellulose, hemicellulose, protein, fat and dietary fibres along with other nutritive elements. One of the most popular ways of exploiting the underutilized plant resources has been through the preparation of composite flours, such as Banana and plantain pseudostem flour. Banana and plantain pseudostem flour is a product obtained by peeling the epidermis of the stem, after which the peeled pseudostem is washed, cut, boiled (for 10 minutes), sliced, dried (at 60°C for 24 h), ground in a blender and sieved to obtain a fine powder (Aziz *et al.*, 2011). Flour made from banana pseudostem is rich in fiber, macro minerals like potassium, sodium, calcium, magnesium and phosphorus, and it can be used in the enrichment of food products such as dairy and bakery products (Thorat and Bobade, 2018).

Studies showed that by the partial replacement of wheat flour with banana pseudostem flour in bakery products, the dietary fiber content was increased (Tiroutchelvameet *et al.*, 2019). Composite flours are mixtures of several flours obtained from cereals, legumes or other plant

resources with or without the presence of wheat flours. Composite flours have gained much importance owing to its enriched nutritional profile and better digestibility. Moreover, the usage of composite flours can help overcome the production deficit of wheat in several tropical countries.

### **3.1.3 Banana and plantain leaf extract**

Banana and plantain leaves have been traditionally used for packaging of certain foods, but new techniques for the valorization of these byproducts are being studied (Fiallos-Cárdenas *et al.*, 2021), such as making use of the leaf extracts. Banana leaf extract is produced by passing the washed leaves three times through a mill called “trapiche”, after which the juice is collected and stored at low temperatures for further use (Fiallos-Cárdenas *et al.*, 2021). It has been shown to be potent in the prevention of enzymatic browning and quality deterioration of freshly cut guava slices during storage (Shomodder *et al.*, 2021). This implies that banana leaf extract can potentially be used in the prevention of enzymatic browning in other food products as well. Studies have also shown that banana leaves present anti-diabetic properties such as a decrease in glucose levels and increase in glycogen and plasma insulin levels when administered to hyperglycemic rats (Kappel *et al.*, 2013), and can therefore serve as a good raw material in the development of functional foods for diabetes patients. Rutin, a pharmacologically active phytochemical that decreases glycemia, increases insulin secretion and inhibits  $\alpha$ -glucosidase is the major component responsible for the antidiabetic activity of banana leaves (Kappel *et al.*, 2013).

### **3.1.4 Banana and plantain inflorescences**

Banana inflorescences have been used in some parts of the world to make pie fillings, salads, increase the yield of meat-based meals, and when converted into flour via drying and grinding, these inflorescences can be used in food enrichment and functional food development due to their low caloric content, high fiber, and high potassium content (Fingolo *et al.*, 2012). They have also been shown to have antioxidant, antidiabetic, antimicrobial, anti-inflammatory, anti-cancer and cardio-protective properties (Lau *et al.*, 2020). Extracts of banana inflorescences have been used in the production of both a beverage and flour which were rich in alkaloids, saponins, glycosides, tannins, flavonoids, and steroids, showed antioxidant activity, and helped to increase breast milk production in lactating mothers (Amornlerdpison *et al.*, 2020).

## **3.2. Value added products from pineapple waste and by-products**

Pineapple is one of the most produced fruits, of which Costa Rica, Philippines, Brazil, Thailand and India are the primary pineapple producers (Mohd *et al.*, 2020). Cultivated in tropical and subtropical zones, pineapple belongs to the Bromeliaceae family and has a unique shape with wide leaves and fruits that differentiates them from other monocotyledons plants (Wali *et al.*, 2019). Besides, pineapple has 47.8 mg of vitamin C and 13 mg of calcium per 100 g of pineapples. There has been a growing demand for pineapple fruits and products for the past few years. Pineapple (*Ananas comosus*) is a tropical fruit with attractive sensorial (mechanical properties, flavor, acidity/sweetness ratio, color) and nutritional (vitamin A, B and C, minerals, fibers, and antioxidants) characteristics. During pineapple processing, transportation and storage, about 80% of the parts such as the crown, peels, leaves, core and stems, are discarded and end up as waste (Hamzah *et al.*, 2021). During processing, large amounts of byproducts, consisting mainly of peel and pomace are generated, representing approximately 30% – 35% of the pineapple fruit, which is usually discarded as low-value byproducts (Varzakas *et al.*, 2016). Apart from processing, pineapple wastes are also generated from poor handling and storage of fresh fruit, or lack of good and reliable transportation systems (Rabiu *et al.*, 2018). Pineapple's

composition, including high sugar content, trace elements (potassium, calcium and magnesium) and polyphenolic compound, has raised the interest in converting these wastes to valuable products. Pineapple can be utilized in the production of various valuable products including nanocrystals, bromelain enzyme, bioactive compound, wine, vinegar, biopolymer, biopackaging, organic acid, adsorbent, biofuel and biogas. Previously, pineapple waste has also been utilised to form phenolic antioxidants, anti-dyeing agents and animal feed (Hossain *et al.*, 2016, Upadhyay *et al.*, 2013).

Increasing demand for pineapple contributes to an enormous pineapple production, and as a consequence, a large amount of waste is generated. About 80% of pineapple parts, including the crown, peels, leaves, core and stems, are discarded during pineapple processing, transportations and storage and ends as waste (Roda *et al.*, 2019, Zainal *et al.*, 2020). It is essential to know the composition and characteristics of pineapple waste before converting them to valuable products. The identification of major polyphenolic compounds from pineapple waste amounts to 31.76 mg of gallic acid, 58.51 mg of catechin, 50 mg of epicatechin and 19.5 mg of ferulic acid per 100 g of dry extracts (Li *et al.*, 2014). Recent research by Dahunsi (Dahunsi *et al.*, 2019) observed that pineapple waste contains 19.4% lignin, 32.4% cellulose and 23.2% hemicellulose.

### **3.2.1 Wine and Vinegar from pineapple peels**

High sugar content in pineapple waste, especially peels, makes them notable as a source for wine and vinegar production. Therefore, it increases the possibilities of utilising pineapple waste for alcohol and acetic acid to produce vinegar (Roda *et al.*, 2019). Beverages such as wine and beer can be made through the alcohol fermentation process and require an additional step such as oxidation to produce vinegar (Roda *et al.*, 2017). Production of vinegar from pineapple peel using three different acetic acid bacteria strains showed that optimum acetic acid yield (6.15 g/L) was found at 72 h fermentation time using propionic bacterium acid ipropionici acetic acid bacterial strain. Umaru. (Umaru *et al.*, 2015) used the juice from pineapple waste to produce wine with 10.8% alcohol content by *Saccharomyces cerevisiae*. Then, it underwent oxidation to acetic acid (vinegar) and contained a total acidity of 3%. In contrast, lower alcohol content (7%) was observed from wine prepared using pineapple's peels and core (Roda *et al.*, 2017). However, 5% acetic acid was obtained despite the low alcohol content, which is higher in the previous study. The difference was due to the different fermentation methods used by the authors. Correspondingly, there is lower alcohol content in wine produced from pineapple wine produced by *Saccharomyces cerevisiae* (6.60%) and *Saccharomyces bayanus* (6.75%) (Ekechukwu *et al.*, 2021) compared to the alcohol content in (Umaru *et al.*, 2015). In another research, (Roda *et al.*, 2017) reported that physical and enzymatic combination before alcohol fermentation were required to produce good quality wines. By varying the *Saccharomyces cerevisiae* strain and temperature, a substantial difference in the wine's fruity character was detected. Other than that, the sensory evaluation of pineapple organic side-stream syrup revealed its potential when combined in bakery products (Tortoe *et al.*, 2013). Although the studies proved that pineapple residues could be utilized as food enhancers and beverages, further studies should explore more the pineapple residue's utilisation potential in the food and beverage industries, in terms of its production by focusing on the quality and purity.

### **3.2.2 Cellulose Nanocrystals from pineapple leaf waste**

The composition of pineapple leaf waste has been shown to be 13.05% lignin, 21.02% hemicellulose and 41.15% cellulose (Hamzah *et al.*, 2021). As a result, pineapple leaves have been used in the development of nanofibers with desirable properties which can be applied in the

food packaging sector (Wahyuningsih *et al.*, 2016), and being plant fibers they can serve as a good potential alternative to synthetic fibers from petroleum-based non-renewable resources (Gaba *et al.*, 2021). Pineapple leaves have also been used for the extraction of ethanol, which is rich in phytochemicals such as *p*-coumaric acid, 1-*o*-*p*-coumaroylglycerol, caffeic acid and 1-*o*-caffeoylglycerol, and when administered to diabetic rats it inhibited the increase in blood glucose and postprandial triglycerides (Upadhyay *et al.*, 2010). Pineapple leaf waste was shown by dos Santos *et al.* (2013) to be a suitable raw material for the production of cellulose nanocrystals, which can be used as a source of dietary fiber in functional foods, and for the production of food thickeners, stabilizers and flavor carriers (Mua *et al.*, 2019). Cellulose nanocrystals (CNC) derived from the abundance of cellulose in the biomass are one of the most favourable materials for nanocomposites. Currently, CNC serve as a reinforcing agent in the nanocomposites field. CNC have a large surface area, high mechanical strength, are non-toxic, hydrophilic, biocompatible and biodegradable ( Raquel *et al.*, 2018, Dos *et al.*, 2013).

### **3.2.3 Bromelain Enzyme from pineapple peel waste**

Pineapple peels are used to process fruit juices both locally and industrially, acting as flavor enhancers in juice making. Pineapple peel drying has been adopted in Cameroon as a strategy to extend the shelf life for the pineapples peels. Waste from pineapple processing could provide a range of value added ingredients for the food industry, including the enzyme bromelain. Bromelain is a proteolytic enzyme that is usually extracted from the stems or juice of pineapple. Bromelain helps the digestion process and is essential in various applications. The main protease that exists in the bromelain enzyme is identified as stem bromelain (EC 3.4.22.32) and fruit bromelain (EC3.4.22.33) (Nor *et al.*, 2015). It has been used commercially in the food industry where it is known for meat tenderizing, brewing, baking and for the production of protein hydrolysates, among other things (Ketnawa *et al.*, 2012).

Pineapple peels have also been used in the alkali extraction of ferulic acid, which is a widely used phenolic antioxidant in the food and cosmetic industries (Upadhyay *et al.*, 2010). These peels which account for 35 to 50% of total pineapple fresh weight (Chongkhong and Tongurai, 2019) have been shown to be a potential raw material for the production of biofuels such as ethanol and hydrogen (Choonut *et al.*, 2014). Work done by Cornelia and Kristyanti (2021) showed that pineapple peels can be used in the production of cider, which is rich in phenolic compounds and has potential antioxidant activity. Gallic acid, catechin, epicatechin and ferulic acid were identified as the main polyphenolics in pineapple peels (Li *et al.*, 2014). Vinegar can also be produced from pineapple peels via fermentation with acetic acid bacteria (Chalchisa and Dereje, 2021). The addition of pineapple peel flour to wheat flour in the production of biscuits led to an increase in the proximate composition and the calcium, potassium, sodium and copper contents of the biscuits (Adeoye *et al.*, 2017), indicating that pineapple peel flour can be used to enrich food products.

### **3.2.4 Bioactive Compound**

Pineapples and their wastes serve as a good source of antioxidants. The formation of free radicals caused by the oxidation of biological molecules can be prevented by an antioxidant, for example, a phenolic compound which has antioxidant, antimicrobial and anticancer properties (Van *et al.*, 2013). Sepúlveda *et al.*, 2018 reported that the extraction of pineapple waste using the auto hydrolysis process resulted in obtaining polyphenols with high antioxidant activity. The study also revealed that pineapple waste is a good source of antioxidants compared to readily available antioxidants in the market, and all pineapple waste extracts showed higher antioxidant activity.

### **3.3. Value added products from cereal waste and byproducts**

The term “cereals” refers to members of the Gramineae family and determines nine species: wheat (*Triticum* spp.), rye (*Secale* spp.), barley (*Hordeum* spp.), oat (*Avena* spp.), rice (*Oryza* spp.), millet (*Pennisetum* spp.), corn (*Zea* spp.), sorghum (*Sorghum* spp.), and triticale (x *Triticosecale* Wittmack)—which is a hybrid of wheat and rye. Cereals are cultivated for the edible components of their grain or the kernel. Strictly speaking, a cereal is a caryopsis which is composed of the fruit coat (pericarp) and a seed. The fruit coat adheres tightly to the seed coat surrounding the remainder of the seed consisting of germ and endosperm. The aleurone layer lies next to the pericarp. This layer is rich in protein and minerals. The endosperm is the large central portion of the kernel made up mostly of starch, and the germ/embryo is the small structure at the lower end of the kernel (Delcour and Hosney, 2010). Cereal grains are usually milled to remove the fibrous bran, which is one of the major by-products of cereal processing.

Although the micronutrients are generally present in higher concentrations in the outer part of the grain, it is often undervalued and used as animal feed (Hemery *et al.*, 2011). Cereal bran is a nutritional storehouse of the grains. The Chemical composition of cereal bran is highly complex and the multiple beneficial effects of cereal bran could be exploited by incorporating it into the daily diet. Besides regular nutrients like proteins, vitamins, minerals, and fats, cereal bran is a good source of functional food ingredients in the form of bioactive compounds such as dietary fiber, phytosterols, polyphenols and phenolic acids which may provide a wide spectrum of biological activities and other health benefits as seen among populations consuming diets based on cereal grains (Patel, 2012).

#### **3.3.1 Products obtained from rice bran**

Rice bran is the most attractive byproduct generated during rice processing because even though it accounts for only 9% of the rice weight, it contains about 65% of the nutrients of the whole rice grain and is rich in proteins, oil, fiber, micronutrients such as vitamins, and minerals, such as aluminum, calcium, chlorine, iron, magnesium, and manganese (Garofalo *et al.*, 2020). It can be used in the production of rice bran oil and as a dietary fiber source in bakery products since it has anti-oxidant and anti-inflammatory properties (Friedman, 2013; Min *et al.*, 2011). The production of rice bran oil is one of the most common uses of rice bran due to the fact that rice bran oil is very rich in  $\gamma$ -oryzanol, tocopherols, tocotrienols, and phytosterols, a powerful antioxidant mixture of bioactive molecules (Garofalo *et al.*, 2020; Lavanya *et al.*, 2017). Rice bran oil can be used as an alternative to bakery shortening, and has shown improvement in the baking quality when added to baked products (Gul *et al.*, 2015). The presence of Gamma-oryzanol in rice bran makes it a very attractive potential food ingredient because this compound has been shown to decrease animal serum-cholesterol levels and anti-inflammatory activities while inhibiting cholesterol oxidation in vitro (Esa *et al.*, 2013). Due to its high fiber content, rice bran can give texture, gelling, thickening, emulsifying and stabilizing properties to certain foods (Gul *et al.*, 2015), which is very useful in the food industry. Rice bran wax has been used in the formation of corn germ oil oleogels, and this formulation showed potential to act as a replacer of commercial shortening in the baking industry (Zhao *et al.*, 2019).

Other than oil, rice bran also has a 10-15% protein content, and these proteins have been found to be of high quality and application in food and pharmaceutical industries due to their unique hypoallergenicity and anticancer effects (Esa *et al.*, 2013). The partial replacement of gelatinized corn flour with rice bran flour in the production of corn flakes and tortilla chips led to an increase in the protein content of these products (Özdestan *et al.*, 2014), making rice bran a

potential beneficial ingredient for the enrichment of bakery products. A protein formulation based on rice bran can be used to target and overcome protein related nourishment disorders (Gul *et al.*, 2015) in some of the underdeveloped countries of the world.

### **3.3.2. Products obtained from corn bran**

Corn bran is mainly made of insoluble fiber, cellulose, hemicellulose, and xylooligosaccharides making it indigestible by enzymes in humans and monogastric animals, but it can be degraded by colon bacterial communities (de Almeida *et al.*, 2021). It can be used as a natural food additive in cellulosic fiber gel (commercially available as Z-Trim), cellulose fiber gum and in corn fiber oil. Submerged fermentation of corn bran with *Monascus purpureus* led to the production of pigments belonging to a group known as azafilones which can be used as colour additives in the food industry (de Almeida *et al.*, 2021). Purple corn bran can be used for the recovery of anthocyanins, which are water-soluble pigments that can provide attractive colors to foods and have potential beneficial health effects such as antioxidant, anti-inflammatory, anticancer, antiobesity, and anti-diabetic properties (Chen *et al.*, 2018). Corn bran hydrolysates have been used in the production of pullulan, which can be used in food stabilization, coating and the production of packaging materials (Haghighatpanah *et al.*, 2021). It can be concentrated and dried to produce crude bio-based fiber gums which can serve as emulsifiers in the food industry (Yadav *et al.*, 2015). Corn bran has also been used in the production of ferulic acid which has many beneficial functions such as antioxidant, antimicrobial, anti-inflammatory, anticancer and free radical scavenging activities (Zhao *et al.*, 2015).

### **3.4. Value added products from palm kernel waste and byproducts**

The oil palm (*Elaeis guineensis* Jacq) is a native plant of the humid tropics of West Africa. Cultivation originated where oil palm trees were inter-planted in traditional agricultural production systems along with other annual and perennial crops. After the extraction of oil from the palm kernel, several by-products are generated such as the empty fruit bunches, palm pressed fibers and shells which are mainly composed of lignin, cellulose, hemicellulose and other carbonaceous materials (Tan, 2006).

#### **3.4.1 Products from palm kernel cake**

Palm kernel oil is obtained from the seed (the kernel or endosperm) which contains about 50 per cent oil. When the oil has been extracted, the residue known as “palm kernel cake” (PKC) is rich in carbohydrates (48%) and proteins (19%), and the ash contains large amounts of potassium. A portion of these wastes is used as feed supplements for livestock (Oluwaseun *et al.*, 2006).

According to Sahin and Elhussein (2018), palm kernel cake is rich in various phytochemicals such as caffeic acid, vanillic acid, d glucuronic acid, ferulic acid, glutaric acid, protocatechuic acid, quinic acid p-coumaric acid, p hydroxybenzoic acid, salicylic acid, shikimic acid, sinapic acid and syringic acid, which have a great role to play in extending the shelf life of several products as well as providing added-value properties with their antioxidant and antimicrobial properties.

Phenolic extracts of palm kernel cake can be used as antioxidants in the food industry, due to the fact that their experimental addition to sunflower oil showed an increase in its induction time by more than 50%. (Tsouko *et al.*, 2018). Hydrolysis of palm kernel cake protein using different proteases led to the production of useful protein hydrolysates or bioactive peptides which showed radical scavenging activity, with the protein hydrolyzed by papain resulting in the production of the hydrolysate with the highest antioxidant activity (Zarei *et al.*, 2012). This further supports the potential of palm kernel cake in the production of antioxidants

for use in the food industry.

In a study carried out by Belo *et al.* (2018) to evaluate the potential of polysaccharides extracted from palm kernel cake for use as prebiotics, it was found that these soluble polysaccharides showed high resistance to hydrolysis when subjected to artificial human gastric juice and promoted the proliferation of *Lb. plantarum* and *Lb. rhamnosus* with a decrease in the pH of the medium and the production of organic acids. This implies that palm kernel cake polysaccharides can be exploited as probiotics.

Through solid-state lacto-fermentation, low molecular weight peptides were generated from palm kernel cake and when added to bread, these peptides increased its shelf life by inhibiting fungal growth since they showed strong antifungal activity against *Aspergillus flavus*, *Aspergillus niger*, *Fusarium* sp., and *Penicillium* sp. (Asri *et al.*, 2020). This shows that peptides extracted from palm kernel cake have potential to be used in the food industry to extend the shelf life of bakery products and other food products thereby promoting food safety, security, and sustainability. Via solid state fermentation, palm kernel cake has also been used in the production of tannase, the enzyme which catalyzes the hydrolysis of tannic acid by breaking its ester and depside bonds releasing glucose and gallic acid. This enzyme is has various uses in the food industry such as the processing of tea, production of gallic acid, treatment of tannery effluents, stabilization of malt polyphenols, clarification of beer and fruit juices, and for the prevention of phenol-induced madeirization in wine and fruit juices (Sabu *et al.*, 2005).

#### **3.4.2. Products from palm pressed fibre**

Palm pressed fiber is a form of recovered fiber from pressed palm fruits which is rich in carotenoids, vitamin E (tocopherol and tocotrienols), and sterols, and has been used for the recovery of palm pressed fiber oil which contains significant quantities of carotenoids (4000–6000 ppm), vitamin E (2400–3500 ppm), sterols (4500–8500 ppm), and coenzyme Q10 quantities greater than those found in crude palm oil (Neoh *et al.*, 2011). As a result, palm pressed fiber oil has antioxidant and anti-inflammatory properties (Teh *et al.*, 2019) which can be applied in the development of functional foods. The presence of phosphorus in palm pressed fiber oil further enhances its antioxidant properties, and this phosphorus can be extracted for use as a food additive (Majid *et al.*, 2012).

#### **3.4.3. Products from palm kernel shells**

Palm kernel shell is a natural fiber which is rich in lignin, cellulose and hemicellulose, and can be used in the production of fiber-reinforced plastics, thereby contributing to the food packaging sector (Baffour-Awuah *et al.*, 2020). It can also be used as an adsorbent for the treatment of heavy metal contaminated water (Baby *et al.*, 2019), making it a useful material for food wastewater treatment processes.

#### **3.4.4. Products from empty palm kernel fruit bunches**

Palm kernel empty fruit bunches are rich in cellulose and nanocellulose. and they have the lowest phenolic content among all the side streams derived from the palm oil production process (Tsouko *et al.*, 2018). However, their phenolic extracts still showed antioxidant activity. Cellulose and nanocellulose have been extracted from empty fruit bunches (Sundalian *et al.*, 2021), and these compounds can be used as food additives or in the development of food packaging materials. Xylan can also be extracted from these fruit bunches and used as a thickening agent, or converted to xylo-oligosaccharides which are prebiotics (Sundalian *et al.*, 2021). These empty fruit bunches can also be converted into valuable compounds such as ethanol, biovanillin, p-hydroxybenzoate and lignin-containing cellulosic nano fibrils (Rame, 2018; Sudiyani, 2013).

### 3.5. Value added products from mango wastes and byproducts

Mango is one of the most important tropical fruits in the world and currently ranked 5th in total world production among the major fruit crops (Wyman., 2001, Berardini *et al.*, 2005, Abdeldaiem *et al.*, 2012). The total amount of mango production in the world was around 35 million tonnes by the year 2009. The amount of mango production in Africa during 2009 is 13.6 million tonnes (Ajila *et al.*, 2013). 1-2% of the production is processed to make products such as jelly powders, nectars, jams, juices, fruit bars, flakes concentrates, and mango chips but the majority of mango production is consumed fresh (Ajila *et al.*, 2010b, Abdeldaiem *et al.*, 2012, Ajila *et al.*, 2013). Mango processing industries dispose of seeds (seed coat and kernel), cull fruits: fresh fruits unsuitable for human consumption, and peel as a by-product.

The waste generated from the mango processing industry, derived mainly from the epicarp and endocarp has been estimated at 75000 MT (Dorta *et al.*, 2012), and is on the rise due to growth in mango fruit production and processing industry. However, there is virtually no commercial utilization of mango seed kernel which in most cases is discarded as waste in the fruit processing industry.

Mango seed is a good source of carbohydrates (58-80%), protein (6-13%) with good 135 profiles of essential amino acids and lipids (6-16%) rich in oleic and stearic acids (Siaka, 2014). The seed has a high protein content with the presence 139 of all essential amino acids at higher levels than those referenced by the FAO as good quality 140 protein. The seed also has high lipid content, these have typical characteristics of a vegetable 141 butter (Muchiri *et al.*, 2012).

Carotenoids play a potentially important role in human health by acting as biological antioxidants, protecting cells and tissues from the damaging effects of free radicals and singlet oxygen and are used as natural colorants in the food industry (Oreopoulou and Tzia, 2007). The carotenoid content was found to be 4–8 times higher in ripe mango peels compared to raw fruit peels.

The dietary fiber content in mango peels of different varieties has been estimated. The total dietary fiber content in dry peel varied from 45 to 78% (Palafox-Carlos *et al.*, 2010). The soluble dietary fiber content in both raw and ripe mangos peels are more than 35% of total dietary fiber. Insoluble dietary fiber relates to both water absorption and intestinal regulation whereas soluble dietary fiber associates with cholesterol in blood and diminishes its intestinal absorption. Mango seed kernel oil has been reported to be a good source of polyunsaturated fatty acids such as oleic and linoleic acids which have health benefits (Kittiphoom *et al.*, 2013). The potent antimicrobial activity demonstrated by the mango kernel extracts could be attributed to the presence of specific phytochemicals such as flavonoids, terpenes, tannins, and coumarins. High antimicrobial and antifungal activity against *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Escherichia coli*, and *Candida albicans* has been reported in kernel powder of South African mango variety (Ahmed *et al.*, 2005).

#### 3.5.1. Starch extraction:

Starch, the principle carbohydrate constituent of most plant materials, merits a detailed investigation to better understand its biochemical and functional characteristics as well as its variations. Extensive research has been conducted on the structure and functional properties of the main starches of commerce, such as wheat, corn, potato, and rice, due to their ready availability and their extensive utilisation in food and non-food applications. The result revealed that mango by-products are a potential starch source due to their high starch content (more than 50%). Using aqueous extraction technique one can extract starch from those sources (Tesfaye *et*

*al.*,2015, Tamrat *et al.*,2016). The extracted starch could be used in textile industries for yarn sizing, for paint, paper, pharmaceuticals, leather, and Bioplastics industries.

### **3.5.2. Enzymes production:**

From the physicochemical properties of mango by-products, the result revealed that this waste could be used in the production of carboxymethyl cellulose, pectinases, and cellulase enzyme. Carboxymethylcellulose and cellulase enzyme could be used in food industry as a viscosity modifier or thickener, and to stabilise emulsions in various products including ice cream, as toothpaste, laxatives, diet pills, water-based paints, detergents, lubricants, textile sizing, and various paper products. As mango by-products are a rich source of pectin, (Malviya *et al.*, 2010, Koubala *et al.*, 2013) so degrading pectin using any microorganisms could be used for fermentative production of pectinases. Pectinase enzymes could be used in processes involving the degradation of plant materials, especially in wine production.

### **3.5.3. Lactic acid production:**

Lactic acid is used as a food preservative, an ingredient in processed foods and is used as a decontaminant during meat processing curing agent, and flavouring agent. It is also used as a starting material in the production of polylactic acid polymer. Lactic acid is produced commercially by fermentation of carbohydrates such as glucose, sucrose, or lactose, or by chemical synthesis but the production of this substance is very costly (Ajila *et al.*, 2010b). The use of agro-industrial wastes would provide an alternative way to produce lactic acid from less costly raw materials. Seeing that the chemical composition of mango by-products they could be used as a source of lactic acid (Chen *et al.*, 2004). The production of lactic acid takes place through a two-step procedure; pretreatment followed by acid hydrolysis of the by-product followed by microbial fermentation. The production of lactic acid from mango by-products seems to have a practical advantage because the method is economically viable since the cost of the raw material (mango by-product) has low cost.

## **3.6. Value added products from citrus fruit wastes and byproducts**

Citrus fruits (CF) are a family of widely consumed fruits worldwide. Citrus plants belonging to the family of Rutaceae which include fruits such as orange. Mandarin, lime, lemon, orange and grape fruit appear as a well known promising source of multiple beneficial nutrients to humans.

In contrast with other types of fruits, citrus fruits have a small edible portion. Hence, large amounts of waste material, such as peels and seeds, are discarded during juice and food processing. Residues of citrus juice production are a source of several compounds, principally by water, soluble sugars, fiber, organic acids, amino acids and proteins, minerals, oils and lipids, and also contains flavonoids and vitamins.

Citrus peel is one of the most underutilized and most geographically diverse biowaste residues on the planet. After juice extraction the residual peel accounts for 50 wt% of the fruit, presenting an environmental problem. (Marn *et al.*, 2007). There is a real challenge to utilize this resource, with 15.6 million metric tonnes of waste produced from 31.2 million metric tonnes of processed citrus fruit annually. (Djilas *et al.*, 2009). Waste orange peel (WOP) is composed of 20% dry matter (sugars, cellulose, hemicellulose, pectin, and d-limonene) and 80% water. (Bampidis *et al.*, 2006). However, research carried out on citrus waste valorization has nearly always focused on production of a single component, such as d-limonene, (Espina *et al.*, 2011) pectin or bioethanol. (Widmer *et al.*, 2010).

By-products from Citrus fruits, including exhausted peel, seeds, pressed pulp, secondary juice (obtained by pressing the residual pulp after the primary juice extraction) and leaves, are a source of polyphenols (i.e., flavonoids and phenolic acids), sugars (i.e., glucose, fructose and sucrose), dietary fibers (i.e., pectin and cellulose), proteins, lipids (i.e., linolenic, oleic, palmitic and stearic acids), organic acids (i.e., citric, malic and oxalic acids) carotenoids (i.e., carotene and lutein), vitamins (i.e., vitamin C and vitamin B complex) and monoterpenes (i.e., limonene and linalool) (Mahato *et al.*, 2018). Molecular composition of each by-product may vary depending on the type of cultivar, the cultivation method, the harvesting time and the degree of ripeness of the fruit.

CF peel represents almost 50% of the wet fruit mass after juice extraction (Sharma *et al.*, 2017), and is particularly high in fragrant compounds, dietary fibers, pectin, natural pigments as well as polyphenols (Rafiq *et al.*, 2018). CF peel is mainly employed for the extraction of essential oils (EOs), which are contained in the oil sacs of both peels and cuticles, although they can also be isolated from seeds or leaves in much lesser quantities. Chemical composition of EOs consists of monoterpenes and sesquiterpenes compounds (i.e., hydrocarbons with two or three isoprene units in their structure) and oxygenated derivatives (i.e., alcohols, ketones, aldehydes and esters). Limonene is the major constituent of EOs extracted from Citrus by-products, whereas  $\beta$ -pinene, sabinene and  $\beta$ -ocimene are characteristic of the EOs from Citrus leaves (Chi *et al.*, 2018). CF EOs have long been used as flavorings in preparation of food, cosmetic and pharmaceutical products and, more recently, have been re-evaluated for their health beneficial properties (Dosoky *et al.*, 2018, Bruni *et al.*, 2019).

Exhausted CF peels are a source of pectin and dietary fibers, which are also present in juice and pulp (Dimopoulou *et al.*, 2019). Pectin is a complex polysaccharide composed of D-galacturonic acid units linked together by  $\alpha$ -1,4 glycosidic bonds, partially esterified with methanol or acetic acid. It commonly exists in complex or insoluble forms, from white to light brown color and, being a naturally gelling agent, it is used as a thickener, emulsifier, texturizer and stabilizer in the preparation of confectionery, jams and jellies, as well as biodegradable products. Dietary fibers are non-starch polysaccharides including at least ten carbohydrate units, not easily digested nor absorbed in the intestine, and they can exist in soluble (i.e., gum, pectin and a part of cellulose) and insoluble forms (i.e., cellulose, hemicellulose and lignin) (Dimopoulou *et al.*, 2019).

Secondary CF juices are a great source of carotenoids and flavonoids that are present also in peels. Carotenoids are pigments biosynthesized in different fruits and vegetables and can be subdivided into two classes: oxygenated carotenoids (or xanthophylls), such as lutein and violaxanthin, and hydrocarbon carotenoids (or carotenes), such as  $\beta$ -carotene and lycopene. (Saini *et al.*, 2018, Sharma *et al.*, 2021). They are precursors of vitamin A, which is involved in epithelial tissues growth, strengthening of the immune system and promotes proper functioning of vision (Widjaja *et al.*, 2018).

Flavonoids are a wide class of secondary metabolites, synthesized by plants to protect against ultraviolet radiation or pathogenic injuries. They are subdivided into six groups, which are flavones, flavanones, flavonols, isoflavones, anthocyanidins and flavanols (Barreca *et al.*, 2017, Durazzo *et al.*, 2019, Barreca *et al.*, 2020, Abbate *et al.*, 2021). Citrus flavonoids have been extensively studied for their anti-cancer (Cirmi *et al.*, 2016), anti-inflammatory (Musumeci *et al.*, 2020) and neuroprotective activities (Cirmi *et al.*, 2016). In particular, naringin, hesperidin, hesperetin, neohesperidin, narirutin and rutin were quantified as the main flavanones of "satsuma mandarin" juice processing waste (Kim *et al.*, 2016).

Another interesting application of citrus fruits by-products is as cattle feed, especially for ruminants. The most important by-products that can be used are fresh or dried pulp, citrus silage, citrus meal and fines, citrus molasses, citrus peel liquor, and citrus activated sludge. Bampidis and Robinson, (2006) characterized physical and nutrient composition, digestibility, fermentation and effects on ruminants (weight and lactating production) of these feeds. Authors stated that these wastes could be effectively used as feedstuff in rations that support growth and lactation in ruminants.

### **3.7. Value added products from potato peels**

Due to increasing urban populations and rising incomes of people living in developing country, the demand for processed food has dramatically increased (Maria *et al.*, 2010). Potato chips are the most common food processed potato product which produce a large amount of potato peels as by-product. Potato peel is a zero value by-product, which occurs in huge amounts after processing. Depending on the peeling method used, it ranges from 15 to 40% of the first product mass. In food processing industries, the most common cause of environmental pollution is associated with organic waste decomposition which occurs when bacterial and other biological forms use the compounds as a source of food. To avoid such problems, the use of potato waste as an antioxidant in food systems due to its high phenolic content has been explored (Sepelev and Galoburda, 2015).

Potato is the world's fourth-largest food crop, following maize, wheat and rice. Its production (225 to 285 million tons) occurs in 10% of cultivated area (Pandey *et al.*, 2008). It is a significant food crop in developing countries, and is often grown at a significant scale in more than 130 countries around the world. Cameroon accounts for 0.1% of the world's total production of potato and 0.8% of Africa's total production (220,000 tons of 29 million tons) (FAOSTAT 2019). Nowadays, types of processed potato products have increased from time to time to meet the tremendous interest of consumers. Food processing factories produce many volumes of wastes as a byproduct (Charmley *et al.*, 2006; Schieber & Aranda, 2009). Potato popularly known as "the king of vegetables" because it grows in more than 100 countries. In developing countries, potato production has increased at an average rate of five percent per year. Currently the total share of developing countries to global potato production rose from 20% to 52%. This is a remarkable achievement, considering the past two decades (FAO, 2008). Potato waste contains valuable chemical components like phenols which are suitable to apply in food preservation and pharmaceutical industries (Grunert, 2018). Food processing industries generate phenolic- rich vegetable by- products, and this has been an area of research investigations as a sources of antioxidants and antimicrobial for food preservation (Pezeshk *et al.*, 2015). The entire tissue of fruits and vegetables is rich in bioactive compounds or phenols but the by- products have higher contents of antioxidant (Sonia *et al.*, 2016). Potato peel is one of the most important waste products with sufficient amount of phenolic compound so this could be used as a replacement for the current synthetic antioxidant and antimicrobial. The dominant phenolic compounds of potato peel extracts are chlorogenic and gallic acids, which are natural antioxidants that prevent oxidation of vegetable oil, and they have been shown to inhibit soybean oil oxidation reactions by minimizing peroxide, totox, and p-anisidine indices (Amado *et al.*, 2014; Mohdal *et al.*, 2010).

Synthetic food preservatives could be used alone or in combination with natural preservatives both synthetic and natural antioxidants been used in food industry; however, application of synthetics preservatives has potential carcinogenic effects but use of natural preservatives alone has a better advantage for human health with low side effect. As a result, attention has being

given to vegetable waste with rich source phenols (Sonia *et al.*, 2016; Tiwari *et al.*, 2009). Phenolic compound is found ubiquitously in plants and is of noticeable interest due to their antioxidant and antimicrobial properties (Pezeshk *et al.*, 2015).

### **3.8. Value added products from coffee wastes and byproducts**

Nowadays, coffee plants are cultivated in more than 70 countries. Coffee berries are obtained from two usually grown species: *Coffea canephora* (Robusta), the most widely cultivated variety, especially in Central Africa, Southeast Asia and Brazil and *Coffea arabica* (Arabica), cultivated in Latin America, Eastern Africa and Asia. About 60% of coffee beans worldwide production is arabica and the rest of 40% is Robusta (Batista *et al.*, 2016).

Coffee berries are picked when ripped, then are processed and dried, becoming coffee beans. Roasting of coffee beans constitutes a very important stage in coffee obtaining process, because it influences physically and chemically properties of beans and determine their sensorial quality, especially flavour and colour. Roasted beans are grinded and brewed with near-boiling water, in order to obtain the coffee beverage. Coffee is one of the most commercialized commodities worldwide, after petroleum. In the same time, it is the second most popular beverage, next after water (Mussatto *et al.*, 2011b; Giroto *et al.*, 2018). Thus, there is a great worldwide interest for its production and commercialization.

As world total coffee consumption increases every year, an important problem becomes to be the major quantities of organic waste resulted from coffee industry: by-products from beans processing and roasting (>50% of the fruit mass) and spent coffee grounds (SCG), from beverage preparation (Campos-Vega *et al.*, 2015a). SCG is the solid residue which remains after roasted coffee beans are grinded and brewed, both in coffee shops chains and in industry, for obtaining instant coffee and represents the most abundant coffee by-product (45%) (Murthy and Naidu, 2012).

SCG chemical composition Spent coffee grounds contain important quantities of organic compounds (phenolics, lipids, proteins, lignin, cellulose, hemicellulose and other polysaccharides), which determine its importance as a real source of valuable products (Kourmentza *et al.*, 2018). Polysaccharides fraction covers about 50% of SCG total mass, of which about 50% are galactomannans, 25% arabinogalactans and 25% cellulose (Oosterveld *et al.*, 2003). Presence of mannose, galactose, glucose and arabinose, polymerized into hemicellulose and cellulose (Ballesteros *et al.*, 2014; Mussatto *et al.*, 2011a) and high content of galactomannans are highlighted in SCG, lignin being also present in a significant amount (Pujol *et al.*, 2013). Dietary fiber represent about 43% of total SCG dry weight (42% insoluble, 1% soluble fibre respectively), which are approved to be used as raw material to develop functional foods. The fibre from SCG includes, among others, resistant starch, oligosaccharides and manno-oligosaccharides (Campos-Vega *et al.*, 2015b; Vázquez-Sánchez *et al.*, 2018; Tian *et al.*, 2017). There are also present some bioactive secondary metabolites, such as diterpenes, sterols, chlorogenic acids, flavonoids and caffeine (Massaya *et al.*, 2019). Caffeine is a major biological active compound of spent coffee grounds (Cruz *et al.*, 2012b). Till present, several important applications have been experimental developed for spent coffee grounds valorisation, especially as biofuels, composts and animal feed, a functional ingredient for food products with real health benefits, bio-composite materials, decontaminants of waste waters.

In addition to SCG application in animal feed, currently there are studies which support the usage of this by-product as a nutraceutical or food functional additive, improving both nutrition and health (del Castillo *et al.*, 2014). SCG constitutes a valuable source of phenolic compounds and melanoidins, which can be further included as functional ingredients in human diet (Borelli

et al., 2004; Mussatto *et al.*, 2011c; Zuorro and Lavecchia, 2012; Xu *et al.*, 2015). Thus, it was demonstrated the SCG antioxidant, antihypertensive and antimicrobial activities in intestine microbiota (Campos-Vega *et al.*, 2015a), with an important role in preventing diseases related to free radicals (Wang *et al.*, 2011). Phytochemicals from SCG can be digested, absorbed and fermented in colon, exerting healthy effects by influencing the metabolic activity of the microbiota (López-Barrera *et al.*, 2016). SCG phenolic extracts can also be used as anti-inflammatory additives (Lopez-Barrera *et al.*, 2016) and dermatological antimelanogenesis agents (Huang *et al.*, 2016). Regarding extracting of natural antioxidants and caffeine from SCG, there were proposed different methods, such as solid-liquid extraction using aqueous alcohol solution (methanol, ethanol and isopropanol) (Mata *et al.*, 2018) or by pressurized liquid extraction (PLE) method with water and ethanol (Shang *et al.*, 2017).

Other potential use of SCG is obtaining of valuable bio-sugars, such as oligosaccharides, manno-oligosaccharides and mannose, after its delignification and defatting, process which proved large-scale feasibility (Nguyen *et al.*, 2019). Peshev *et al.* (2018) revealed utilization of SCG for obtaining water extracts with sufficiently high caffeine concentration. Appliance of nano filtration to these extracts, by using a suitable membrane, conducted to valuable products, as permeate and retentate fractions. Permeate can be further used for soft and energy drinks, while retentate for coffee drink or as functional food ingredient.

### **3.9. Value added products from *Aframomum meleguta* (alligator pepper) wastes and byproducts**

Alligator pepper (*Aframomum meleguta*) otherwise known as grains of paradise is a perennial deciduous herb of the Zingiberaceae family of plants native to the swampy areas on the West African coasts. Others in that family include *A. danielli*, *A. citratum* and *A. exscapum*. Widely known for its hot, spicy, and aromatic seeds, it is also called as mbongo spice, Afrika kakulesi, or Guinea pepper in some parts of Africa. (Desai *et al.*, 2013).

Alligator pepper seeds have been known for ages among spices and ingredients for different African traditional and socio-cultural applications in many parts of Africa and some parts of Asia. (Anupam *et al.*, 2018). It is widely documented as an antidote for some ailments in human health such as body pains, diarrhea, sore throat, catarrh and rheumatism. Based on current research, different parts of the plant possess specific secondary metabolites such as flavonoids, phenolic compounds, alkaloids, tannins, terpenoids, saponins, and cardiac glycosides that have healing potential and medicinal and therapeutic purposes. In addition, Ethanol extracts from the seeds indicated antibacterial and antiseptic properties found useful for wound-care and antidote for certain infections (Okwu 2004). The active compounds found in alligator pepper are in the class of naturally occurring food preservatives. In some traditional African meetings and events like naming ceremonies, marriages and funeral, it is snacked upon with bitter kola and kola nuts in customary rites. Apart from its medicinal applications, Sunil *et al.* (2018), documented that alligator pepper seeds contain few calories of energy and substantial amount of iron, magnesium, and calcium (Yashin *et al.*, 2017). Okunade *et al.* (2019), reported that it contains 7.5, 4.78, 2.84, and 13.01 g/100 g dry matter of fat, crude fibre, ash, and protein, respectively. It is therefore no surprise that alligator pepper also has culinary applications as ingredient in assorted dishes such as pepper soup and barbecues. Its essential oil has been documented as having pharmaceutical and industrial applications in flavors and perfumery. The pulp and peels of this wild fruit can be used to produce pectin which can be used in the food industry as a gelling agent.

## **4. CHALLENGES IN AGRO-FOOD WASTE AND BY-PRODUCT VALORIZATION IN CAMEROON**

The industrial recovery of valuable compounds from food by-products is a story that started few years ago. Nowadays, at least 50 companies around the world recover valuable compounds from food waste and sell them as clean-label ingredients for processed food products (e.g., natural preservatives to maintain shelf-life requirements, functional compounds), without impacting flavor or texture (Galanakis *et al.*, 2016). These commercially available applications inspired the scientific community to intensify its efforts for the valorization of all kinds of food by-products for recovery purposes. However, the industrialization of such processes and the implementation of the “Universal Recovery Strategy” is not an easy task. It typically includes laboratory research, transfer to pilot plan and full-scale production, protection of intellectual properties, development of definite applications and commercialization problems. For example, waste emerges seasonally and often in large quantities, whereas it is prone to microbial spoilage. Proper management of collection processes, cooling of the material and addition of preservatives can provide solutions.

Other problems such as the broad content of wasted by-products may be modified by adding a pretreatment step. Besides, transition of batch to continuous processes is usually accompanied with extension of mixing and heating time, heavier handling, increased air incorporation and higher degree of scrutiny. Subsequently, process cost is increased, as industrially recovered compounds are used in food formulations in higher concentrations compared to laboratory-

recovered compounds. Therefore, solving these issues are necessary in order to ensure the sustainability of the process, the economic benefit and the perpetual establishment of the derived products in the market (Galanakis *et al.*, 2016).

A commercially feasible product can be generated only if a certain degree of flexibility and alternative solutions can be adapted in the developing methodology, e.g., less extraction steps are cheaper and scale up easier. However, they generate cruder products with lower concentrations of target compounds; thus it is recommended to use nonthermal technologies, green solvents and safer materials (possessing GRAS-status) to improve efficiency and clarification of the final product. The development of precise applications for well-defined markets is also very important (Galanakis, 2012). For example, natural antioxidants like polyphenols to be used in fresh, mom-approved and kid-friendly foods with clean ingredient lines and a minimum of three-month refrigerated shelf-life. Besides, modern new products should aim at the fulfillment of consumer needs and the realization of consumer value rather than at the development of products or enabling technologies per se. This means that the developed products should meet the high expectations of the consumers in an increasingly competitive market, e.g., development of “green” “organic” and “Protected Designation of Origin” products. Unlike the needs to delight the consumer and minimize environmental impact, developers should also ensure that the final product and process meet particular specifications, e.g., provide clean-label ingredients without impacting flavor or texture. Currently, the manufacturer’s label typically provides only limited information about the origin and composition of the used extract in the final product formulation. A clearer label of the products containing recovered compounds would enable nutritionists and/or pharmacists to be more confident when recommending these products.

In addition, legal authorities regulate the way in which companies can advertise health benefits. However, those regulations (driven by the need of protecting consumers from dubious claims) create implications for the food industry. This is mainly because the demonstration of health benefits is very costly (constituting a high cost especially to small and medium-sized companies) and the risk of claims rejection by the corresponding authorities is too high. So far, only a small number of compounds and products have been approved, whereas the overwhelming majority of health claims has been declined by regulatory bodies (Galanakis *et al.*, 2015c). The legislation challenges regulating health beneficial dietary products are lying to the nature of the products, which have the characteristics of both food and biologically active ingredients. For instance, EFSA of EU has approved health claims for only a small number of compounds (e.g., hydroxytyrosol in olive oil) (Galanakis *et al.*, 2016). This means that if you recover hydroxytyrosol from OMW in order to fortify foods (e.g., bakery, meat products or even oils), it is not allowed to add the health claim on the label.

Considering all the above innovation challenges, it can be concluded that food waste recovery may be an interesting but difficult to implement opportunity for food companies (compared to food waste valorization for nonfood applications). Decision upon recovery or valorization strategy should always be taken account in relation to the substrate and industry’s goals.

## 5. CONCLUSION

Large amounts of diversified agro-food waste and by-products are produced in Africa general and in Cameroon in particular, which courses enourmous environmental and economic problems. Biorefineries can have the capacity of reducing the environmental and economic burdens of

agro-food and, at the same time, permit the production of value added biochemicals and products through application of appropriate process technologies. This work reviews agro-food waste and by-product production and provides inside on the quantity and quality of the waste as a renewable bioresources. It further highlights the current traditional methods of agro-food waste valorisation for the production of environmentally friendly value added products within the concept of bioeconomy or circular economy. The paper presents the exciting future and challenges of circular technology development in Cameroon, which largely delay on developing the current traditional process technologies for agro-food waste valorisation to ensure more sustainable production of quality value added products. To address such challenges, proposals on implementing bioeconomy concepts in the food industry to ensure valorization of food waste as a route to innovation and wealth creation are indicated. More research and development should be carried on traditional process technology for agro-food waste valorisation, which will create the pathway to scale-ups and industrialisation.

## REFERENCES

- Abbate, F., Maugeri, A., Laurà, R., Levanti, M., Navarra, M., Cirmi, S. and Germanà, A. (2021). Zebrafish as a Useful Model to Study Oxidative Stress-Linked Disorders: Focus on Flavonoids. *Antioxidants*. 10, 668. [CrossRef] [PubMed]
- Abdeldaiem, M. H. and Hoda, G. M. A. (2012). Use of irradiated mango (*Mangifera indica*) peels powder as a potential source of dietary fibre and antioxidant in a beef burger. *Journal of Applied Sciences Research* 8: 3677-3687.
- Adeoye, B. K., Alao, A. I. and Famurewa, J. A. V. (2017). Quality Evaluation Biscuits Produced from Wheat and Pineapple Peel Flour. *Applied Tropical Agriculture*, 22(2), 210-217.
- Adewuyi, G. O., Obi-Egbedi, N. O. and Babayemi, J.O. (2008). Evaluation of ten different African wood species for potash production. *Int. J. Physical Sciences*. 3: 63-68.
- Ahmed, I. S., Tohami, S. M., Almagboul, Z. A. and Verpoorte, R. (2012) Characterization of anti-microbial compounds. *University of Africa, Journal of Sciences*. Volume 2, 77-91
- Ajiboye, T. O., Komolafe, Y. O., Yakubu, M. T. and Ogunbode, S. M. (2013). Effects of iron on the redox status of cellular system of male rats. *Toxicology and Industrial Health*. 3(3): 1-9.
- Ajila, C. M. and Prasada Rao, U. J. S. (2013). Mango peels dietary fibre: Composition and associated bound phenolics. *Journal of Functional Foods* 5: 444-450.
- Ajila, C.M., Aalami, M., Leelavathi, K. and Prasada Rao, U. J. S. (2010b). Mango peel powder: A potential source of antioxidant and dietary fiber in macaroni preparations. *Innov. Food Sci. Emerg.* 11: 219-224.
- Alancay M. M., Lobo, M. O., Quinzio, C. M. and Iturriaga, L. B., (2017). Extraction and physicochemical characterization of pectin from tomato processing waste. *J. Food. Meas. Charact.*, 11, 2119–2130. <https://doi.org/10.1007/s11694-017-9596-0>
- Al-Wandawi, H., Abdul Rehman, M. H. and Al shaikhly, K. A. (1985). Tomato processing wastes as essential raw materials source. *J. Agric. Food Chem.* 33, 804–807. [CrossRef]
- Amado, I. R., Franco, D., Sánchez, M., Zapata, C., & Vázquez, J. A. (2014). Optimisation of antioxidant extraction from *Solanum tuberosum* potato peel waste by surface response methodology. *Food Chemistry*, 165, 290-299. Doi: <https://doi.org/10.1016/j.foodchem.2014.05.103>

- Amornlerdpison, D., Choommongkol, V., Narkprasom, K. and Yimyam, S. (2020). Bioactive Compounds and Antioxidant Properties of Banana Inflorescence in a Beverage for Maternal Breastfeeding. *Appl. Sci.*, 11, 343.
- Anupam, K. R., Sachan, S. K., Kiran, K. and Deepti, S. (2018). Medicinal uses of spices used in our traditional culture: Worldwide. *Indian Journal Pharmaceutical Sciences*, 40: 104-105.
- Asri, N. M., Muhialdin, B. J., Zarei, M., Saari, N. (2020). Low molecular weight peptides generated from palm kernel cake via solid state lacto-fermentation extend the shelf life of bread. *LWT - Food Science and Technology*, 134, 110206. Doi: <https://doi.org/10.1016/j.lwt.2020.110206>
- Atef, M., and Mahdi Ojagh, S. (2017). Health benefits and food applications of bioactive compounds from fish byproducts: A review. *Journal of Functional Foods*, 35, 673–681. <https://doi.org/10.1016/j.jff.2017.06.034>
- Ayala, J., Vega, V., Rosas, C., Palafox, H., Villa, J., Siddiqui, W., et al. (2011). Agro-industrial potential of exotic fruit byproducts as a source of food additives. *Food Res. Int.*, 44, 1866–1874. doi: 10.1016/j.foodres.2011.02.021
- Aziz, N. A. A., Ho, L. H., Azahari, B., Bhat, R., Cheng, L. H. and Ibrahim, M. N. M. (2011). Chemical and functional properties of the native banana (*Musa acuminata*×*balbisiana* Colla cv. Awak) pseudo-stem and pseudo-stem tender core flours. *Food Chemistry*, 128(3), 748-753. doi: 10.1016/j.foodchem.2011.03.
- Baby, R., Saifullah, B. and Hussein, M. Z. (2019). Palm Kernel Shell as an effective adsorbent for the treatment of heavy metal contaminated water. *Scientific Reports*, 9, 18955. Doi: <https://doi.org/10.1038/s41598-019-55099-6>
- Baffour-Awuah, E., Akinlabi, S. A., Jen, T. C., Hassan, S., Okokpujie, I. P. and Ishola, F. (2020). Characteristics of Palm Kernel Shell and Palm Kernel Shell Polymer Composites: A Review. *IOP Conf. Series: Materials Science and Engineering*, 1107, 012090. doi:10.1088/1757-899X/1107/1/012090
- Bampidis, V. A. and Robinson, P. H. (2006). *Anim. Feed Sci. Technol.* 128, 175– 217.
- Banerjee, S., Ranganathan, V., Patti, A. and Arora, A. (2018). Valorisation of pineapple wastes for food and therapeutic applications. *Trends Food Sci. Technol.* 82, 60–70. [CrossRef]
- Barreca, D., Gattuso, G., Bellocco, E., Calderaro, A., Trombetta, D., Smeriglio, A., Lagana, G., Daglia, M., Meneghini, S. and Nabavi, S. M. (2017). Flavanones: Citrus phytochemical with health-promoting properties. *BioFactors*. 43, 495–506. [CrossRef] [PubMed]
- Barreca, D., Mandalari, G., Calderaro, A., Smeriglio, A., Trombetta, D., Felice, M. R. and Gattuso, G. (2020). Citrus Flavones: An Update on Sources, Biological Functions, and Health Promoting Properties. *Plants*. 9, 288. [CrossRef] [PubMed]
- Batista, L. R., Chalfoun de Souza, S. M., Silvae Batista, C. F. and Schwan, R. F. (2016). *Encyclopedia of Food and Health. Coffee: Types and Production*, pp. 244
- Bello, B., Mustafa, S., Tan, J. S., Ibrahim, T. A. T., Yew Joon Tam, Y. J., Ariff, A. B., Manap, M. Y. and Abbasiliasi, S. (2018). Evaluation of the effect of soluble polysaccharides of palm kernel cake as a potential prebiotic on the growth of probiotics. *Biotech.*, 8, 346-360. Doi: <https://doi.org/10.1007/s13205-018-1362-4>
- Benoit Daviron. (2004). *Manuel d'élaboration des Politiques Agricoles. Construction d'argumentaires pour l'intervention publique en Afrique de l'Ouest et du Centre.* Editions du Gret.
- Berardini, N., Knodler, M., Schieber, A. and Carle, R. (2005). Utilisation of mango peels as a source of pectin and polyphenolics. *Innov. Food Sci. Emerg.* 6: 442-452.

- Bergeson, T. L., Opio, C. and MacMillan, P. D. (2018). Crop ash filtrate influence on cooking time and sensory preferences for dried black beans (*Phaseolus vulgaris* L.). *Afr J Food Sci.* 10:132–42.
- Blanke, M. (2015). Challenges of Reducing Fresh Produce Waste in Europe—From Farm to Fork. *Agriculture*, 5, 389–399. [CrossRef]
- Borrelli, R.C., Esposito, F., Napolitano, A., Ritieni, A. and Fogliano, V. (2004). Characterization of a new potential functional ingredient: coffee silverskin. *Journal of Agricultural and Food Chemistry* 52, 1338–1343
- Bracco S, Calicioglu O, Juan MGS. And Flammini A. (2018). Assessing the contribution of bioeconomy to the total economy: A review of national frameworks. *Sustainability*. 10:1698. DOI: 10.3390/su10061698
- Briones-Labarca, V., Giovagnoli-Vicuña, C. and Cañas-Sarazúa, R. (2019). Optimization of extraction yield, flavonoids and lycopene from tomato pulp by high hydrostatic pressure-assisted extraction. *Food Chem.* 278, 751–759. <https://doi.org/10.1016/j.foodchem.2018.11.106>
- Bruni, R., Barreca, D., Protti, M., Brighenti, V., Righetti, L., Anceschi, L., Mercolini, L., Benvenuti, S., Gattuso, G. and Pellati, F. (2019). Botanical Sources, Chemistry, Analysis, and Biological Activity of Furanocoumarins of Pharmaceutical Interest. *Molecules*. 24, 2163. [CrossRef] [PubMed]
- Campos-Vega, R., Vázquez-Sánchez, K., López-Barrera, D., Loarca-Piña, G., MendozaDíaz, S. and Oomah, B. D. (2015a). Simulated gastrointestinal digestion and in vitro colonic fermentation of spent coffee (*Coffea arabica* L.): Bioaccessibility and intestinal permeability. *Food Research International* 77, 156–161
- Caseiro, M., Ascenso, A., Costa, A., Creagh-Flynn, J., Johnson, M. and Simões, S. (2020). Lycopene in human health. *Lwt-Food Sci. Technol.* 127, 109323. <https://doi.org/10.1016/j.lwt.2020.109323>
- Chalchisa, T. and Dereje, B. (2021). From waste to food: utilization of pineapple peels for vinegar production. *MOJ Food Processing & Technology*, 9(1), 1-5.
- Charmley, E., Nelson, D. and Zvomuya, F. (2006). Nutrient cyclin in the vegetable processing industry: Utilization of potato by-products. *Canadian Journal of Soil Science*, 86, 621–629.
- Chen, L., Yang, M., Mou, H. and Kong, Q. (2004). Ultrasound-assisted extraction and characterization of anthocyanins from purple corn bran. *Journal of Food Processing and Preservation*, 13377. Doi: <https://doi.org/10.1111/jfpp.13377>
- Chi, P. T. L., Van Hung, P., Le Thanh, H. and Phi, N. T. L. (2018). Valorization of citrus leaves: Chemical composition, antioxidant and antibacterial activities of essential oils. *Waste Biomass Valorization*. 11, 4849–4857. [CrossRef]
- Chongkhong, S. and Tongurai, C. (2019). Optimization of soluble sugar production from pineapple peel by microwave-assisted water pretreatment. *Songklanakarin J. Sci. Technol.*, 41 (1), 237-245.
- Choonut, A., Saejong, M. and Sangkharak, K. (2014). The Production of Ethanol and Hydrogen from Pineapple Peel by *Saccharomyces cerevisiae* and *Enterobacter aerogenes*. *Energy Procedia*, 52, 242 – 249.
- Cifarelli A., Cigognini, I. M., Bolzoni, L. and Montanari, A. (2019). Physical– chemical characteristics of cutin separated from tomato waste for the preparation of bio-lacquers. *Adv. Sci. Eng.*, 11, 33–45. <https://doi.org/10.32732/ase.2019.11.1.33>

- Cigognini, I., Montanari, A., Carrerars, R. D. L. T. and Montserrat, G. C. B. (2015). Extraction method of a polyester polymer or cutin from the wasted tomato peels and polyester polimer so extracted (World Intellectual Property Organization Patent N. WO2015028299A1). <https://patents.google.com/patent/WO2015028299A1/en>
- Cirmi, S., Ferlazzo, N., Lombardo, G.E., Ventura-Spagnolo, E., Gangemi, S., Calapai, G. and Navarra, M. (2016). Neurodegenerative diseases: Might citrus flavonoids play a protective role? *Molecules*. 21, 1312. [CrossRef]
- Cornelia, M. and Kristyanti, T. (2021). Utilization of Pineapple's (*Ananas comosus* L. Merr) Peel Waste as Raw Material in Cider Making. In *Proceedings of the 3rd International Conference of Computer, Environment, Agriculture, Social Science, Health Science, Engineering and Technology (ICEST 2018)*, 258-263.
- Cristian, T., Ramírez-Guzman, N., Londoño-Hernandez, L., Martinez-Medina, G. A., Díaz-Herrera, R., Navarro-Macias, V., Alvarez-Pérez O. B., Picazo, B., Villarreal, M., Ascacio-Valdes, J. and Aguilar, C. N. (2018). Food waste and byproducts: An opportunity to minimize malnutrition and hunger in developing countries. *Front. Sustain. Food Syst.*, 2(52), 1-17.
- Cruz, R., Cardoso, M. M., Fernandes, L., Oliveira, M., Mendes, E., Baptista, P. and Casal, S. (2012b). Espresso coffee residues: A valuable source of unextracted compounds. *Journal of Agricultural and Food Chemistry* 60(32), 7777–7784
- Dahunsi, S. O. (2019). Liquefaction of pineapple peel: Pretreatment and process optimization. *Energy*, 185, 1017–1031. [CrossRef]
- De Almeida, A. B., Santos, N. H., de Lima, T. M., Santana, R. V., Filho, J. G. O., Peres, D. S. and Egea, M. B. (2021). Pigment bioproduction by *Monascus purpureus* using corn bran, a byproduct of the corn industry. *Biocatalysis and Agricultural Biotechnology*, 32, 101931. Doi: <https://doi.org/10.1016/j.bcab.2021.101931>
- Del Castillo, M.D., Martinez-Saez, N. and Ullate, M. (2014). Food formulation comprising spent coffee grounds. WO2014128320
- Del Valle, M., Cámara, M. and Torija, M.-E. (2006). Chemical characterization of tomato pomace. *J. Sci. Food Agr.*, 86, 1232–1236. <https://doi.org/10.1002/jsfa.2474>
- Delcour, J. A. and Hoseney, R. (2010). Principles of cereal science and technology. AACC International press, USA.
- Desai, N. N., Modi, V. M. and Patel, D. B. (2013). Analysis of physical and chemical properties of dill seed. *African Journal of Biotechnology*, 6 (21), 3-6.
- Dimopoulou, M., Alba, K., Campbell, G. and Kontogiorgos, V. (2019). Pectin recovery and characterization from lemon juice waste streams. *J. Sci. Food Agric.* 99, 6191–6198. [CrossRef]
- Djilas, S., Canadanovic-Brunet, J. and Cetkovic, G. (2009). *Chem. Ind. Chem. Eng. Q.* 15, 191–202.
- Domínguez, E., Heredia-Guerrero, J. A. and Heredia, A. (2011). The biophysical design of plant cuticles: an overview. *New Phytol.*, 189, 938–949. <https://doi.org/10.1111/j.1469-8137.2010.03553.x>
- Dorta, E., Lobo, G. and Gonzalez, M. (2012). Using drying treatments to stabilise mango peel and seed: Effect on antioxidant activity. *Lebensmittel-Wissenschaft und-Technologie*, 45, 261. Doi: 10.1016/j.lwt.2011.08.016.

- Dos Santos, R. M., Neto, W. P. F., Silvério, H. A., Martins, D. F., Dantas, N. O. and Pasquini, D. (2013). Cellulose nanocrystals from pineapple leaf, a new approach for the reuse of this agro-waste. *Industrial Crops and Products*, 50, 707-714.
- Dosoky, N. S. and Setzer, W. N. (2018). Biological Activities and Safety of Citrus spp. Essential Oils. *Int. J.Mol. Sci.* 19, 1966. [CrossRef]
- Doumta, C. F. and Tchiégang, C. (2012). Standardisation process to soften African Locust Beans Seeds (*Parkia biglobosa* Benth) by a traditional method. *Food Glob Sci Books*.6:33–7.
- Durazzo, A., Lucarini, M., Souto, E.B., Cicala, C., Caiazzo, E., Izzo, A.A., Novellino, E. and Santini, A. (2019). Polyphenols: A concise overview on the chemistry, occurrence, and human health. *Phytother. Res. PTR.* 33, 2221–2243. [CrossRef] [PubMed]
- Ekechukwu, C. C., Ikele, C. C., Okafor, C. A., and Igwebuike, I.C. (2021). Comparative assessment of three yeast samples for wine production from pineapple. *Multidiscip. Sci. J.* 2021, 3, e2021003. [CrossRef]
- Enders, A., Hanley, K., Whitman, T., Joseph, S. and Lehmann, J. (2012). Characterization of biochars to evaluate recalcitrance and agronomic performance. *Biores. Technol.*, 114, 644-653. Doi: <https://doi.org/10.1016/j.biortech.2012.03.022>
- Esa, N. M., Ling, T. B. and Peng, L. S. (2013). By-products of Rice Processing: An Overview of Health Benefits and Applications. *J Rice Res*, 1, 107. doi: 10.4172/jrr.1000107
- Espina, L., Somolinos, M., Loran, S., Conchello, P., Garcia, D. and Pagan, R. (2011). *Food Control*. 22, 896–902.
- European Commission. (2015). EIP-AGRI Workshop Opportunities for Agriculture and Forestry in the Circular Economy. Workshop Report 28-29 October 2015. Brussels, Belgium. Available from: [https://ec.europa.eu/eip/agriculture/sites/agri-eip/files/eip-agri\\_ws\\_circular\\_economy\\_final\\_report\\_2015\\_en.pdf](https://ec.europa.eu/eip/agriculture/sites/agri-eip/files/eip-agri_ws_circular_economy_final_report_2015_en.pdf) [Accessed: 13 August 2020]
- FAOSTAT (2019). Data: Production. Crop. Available online: <http://www.fao.org/faostat/en/#data/QC>.
- Faustino, M., Veiga, M., Sousa, P., Costa, E., Silva, S. and Pintado, M. (2019). Agro-Food Byproducts as a New Source of Natural Food Additives. *Molecules*, 24(6), 1056. Doi: 10.3390/molecules24061056
- Fiallos-Cárdenas, M., Ramírez, A. D., Pérez-Martínez, S., Romero, B. H., Ordoñez-Viñan, M., Ruiz-Barzola, O. and Reinoso, M. A. (2021). Bacterial Nanocellulose Derived from Banana Leaf Extract: Yield and Variation Factors. *Resources*, 10(12), 121. doi: <https://doi.org/10.3390/resources10120121>
- Fingolo, C. E., Braga, J. M. A., Ana C.M. Vieira, A. C. M., Moura, M. R. L. and Kaplan, M. A. C. (2012). The natural impact of banana inflorescences (*Musa acuminata*) on human nutrition. *Anais da Academia Brasileira de Ciências*, 84(4), 891-898.
- Folefack, D. P., FongangFouepe, G. H., Adamou, K. M., Ebongue, J. P., Bikoi, A. and Noupadja, P. (2017). Analysis of the plantain supply system of markets in the city of Douala. *J. Econ. Sustain. Dev.* 8 (6). <http://www.iiste.org> (ISSN 2222-1700 (Paper) ISSN 2222-2855 (Online)).
- FAO. (2011a). Global food losses and food waste – extent, causes and prevention, by J. Gustavsson, C.Cederberg, U. Sonesson, R. van Otterdijk & A. Meybeck. Rome (<http://www.fao.org/docrep/014/mb060e/mb060e00.pdf>)
- FAO. (2008). The global potato economy, (International Year of Potato, 2008). Rome, Italy: FAO

- FAO. (2013). Food Wastage Footprint: Impacts on Natural Resources : Summary Report. <http://site.ebrary.com/id/10815985>. Google Scholar
- FAO. (2014). Panorama de la Seguridad Alimentaria y Nutricional en América Latina y el Caribe. FAO. Available online at: <http://www.fao.org/docrep/018/i3068s/i3068s.pdf> (Accessed July 18, 2018).
- FAO. (2015). Global Initiative on Food Loss and Waste Reduction, 8.
- FAO. (2019). The State of Food and Agriculture. Moving Forward on Food Loss and Waste Reduction; Food and Agriculture Organization of the United Nations: Rome, Italy. 182p.
- FAO. (2019b). The State of Food Security and Nutrition in the World: Safeguarding against Economic Slowdowns and Downturns.
- Foresight (2011). Foresight Project on Global Food and Farming Futures. Synthesis Report C7: Reducing Waste. The Government Office for Science, London, United Kingdom.
- Franca, S. and Oliveira, L. S. (2009). Alternative uses for coffee husks - a solid waste from green coffee production In: Chemical, Biological and Environmental Engineering. Singapore: World Scientific Publishing Co. Pte. pp. 21-4.
- Frazzoli, C. and Mantovani, A. (2020). Toxicological risk factors in the burden of malnutrition: The case of nutrition (and risk) transition in sub-Saharan Africa. *Food Chem. Toxicol.* 146, 111789. [CrossRef] [PubMed]
- Friedman, M. (2013). Rice bran, rice bran oils, and rice hulls: composition, food and industrial uses, and bioactivities in humans, animals, and cells. *J Agric Food Chem*, 61, 10626-10641. Doi: <https://doi.org/10.1021/jf403635v>
- Fritsch, C., Staebler, A., Happel, A., Márquez, M. A. C., Aguiló-Aguayo, I., Abadias, M., Gallur, M., Cigognini, I. M., Montanari, A., López, M. J., Suárez-Estrella, F., Brunton, N., Luengo, E., Sisti, L., Ferri, M. and Belotti, G. (2017). Processing, Valorization and Application of Bio-Waste Derived Compounds from Potato, Tomato, Olive and Cereals: A Review. *Sustainability*, 9, 1492, doi: 10.3390/su9081492
- Gaba, E. W., Asimeng, B. O., Kaufmann, E. E., Katu, S. K., Foster, E. J. and Tiburu, E. K. (2021). Mechanical and Structural Characterization of Pineapple Leaf Fiber. *Fibers*, 9, 1-11.
- Galanakis, C. M. (2012). Recovery of high added-value components from food wastes: conventional, emerging *technologies and commercialized applications*. *Trends Food Sci. Technol.*, 26, 68-87.
- Galanakis, C. M. (2016). Separation of functional macromolecules and micromolecules: from ultrafiltration to the border of nanofiltration. *Trends Food Sci. Technol.*, 42, 44-63.
- Galanakis, C. M., Tornberg, E. and Gekas, V. (2015c). Dietary fiber suspensions from olive mill wastewater as potential fat replacements in meatballs. *LWT-Food Sci. Technol.*, 43, 1018-1025.
- Garofalo, S. F., Tommasi, T. and Fino, D. (2020). A short review of green extraction technologies for rice bran oil. *Biomass Conversion and Biorefinery*. Doi: <https://doi.org/10.1007/s13399-020-00846-3>
- George, B., Kaur, C., Khurdiya, D. S. and Kapper, H.C. (2014). Antioxidants in tomato (*Lycopersicon esculentum*) as a function of genotype. *Food Chem.* 84, 45–51. [CrossRef]
- Giroto, F., Pivato, A., Cossu, R., Nkeng, G. E. and Lavagnolo, M. C., (2018). The broad spectrum of possibilities for spent coffee grounds valorisation. *Journal of Material Cycles and Waste Management* 20, 695–701

Goletti, F. and Wolff, C. (1999). *The Impact of Postharvest Research. Markets and Structural Studies Division; International Food Policy Research Institute: Washington, DC, USA.* 109p

Górecka, D., Wawrzyniak, A., Jędrusek-Golińska, A., Dziedzic, K., Hamułka, J., Kowalczewski, P.Ł. and Walkowiak, J. (2020). Lycopene in tomatoes and tomato products. *Open Chem.* 18, 752–756. <https://doi.org/10.1515/chem-2020-0050>.

Grassino, A. N., Halambek, J., Djaković, S., Rimac Brnčić, S., Dent, M. and Grabarić, Z. (2016). Utilization of tomato peel waste from canning factory as a potential source for pectin production and application as tin corrosion inhibitor. *Food Hydrocol.*, 52, 265–274. <https://doi.org/10.1016/j.foodhyd.2015.06.020>

Grunert, K. G. (2018). Food quality and safety: Consumer perception and demand. *European Review of Agricultural Economics*, 32, 369–391.

Gul, K., Yousuf, B., Singh, A. K., Singh, P. and Wani, A. A. (2015). Rice bran: Nutritional values and its emerging potential for development of functional food: A review. *Bioactive Carbohydrates and Dietary Fibre*, 6, 24-30. Doi: <http://dx.doi.org/10.1016/j.bcdf.2015.06.002>

Gustavsson, J., Cederberg, C., Sonesson, U., van Otterdijk, R. and Meybeck, A. (2011). *Global Food Losses and Food Waste—Extent, Causes and Prevention*; Food and Agriculture Organization of the United Nations (FAO): Rome, Italy. p. 37.

Haghighatpanah, N., Khodaiyan, F., Kennedy, J. F. and Hosseini, S. S. (2021). Optimization and characterization of pullulan obtained from corn bran hydrolysates by *Aerobasidiom pullulan* KY767024. *Biocatalysis and Agricultural Biotechnology*, 33, 101959. Doi: <https://doi.org/10.1016/j.bcab.2021.101959>

Hamzah, A. F. A., Hamzah, M. H., Man, H. C., Jamali, N. S., Siajam, S. I. and Ismail, M. H. (2021). Recent Updates on the Conversion of Pineapple Waste (*Ananas comosus*) to Value-Added Products, Future Perspectives and Challenges. *Agronomy*, 11, 1-27.

Hemery, Y., Chaurand, M., Holopainen, U., Lampi, A. M., Lehtinen, P., Piironen, V., Sadoudi, A. and Rouau, X. (2011). Potential of dry fractionation of wheat bran for the development of food ingredients, part I: influence of ultra-fine grinding. *J Cereal Sci*, 53, 1-8. Doi: <https://doi.org/10.1016/j.jcs.2010.09.005>

Hossain, M.F. (2016). World pineapple production: An overview. *African J. Food, Agric. Nutr. Dev.* 16, 11443–11456. [CrossRef]

Huang, H.C., Wei, C.M., Siao, J.H., Chang, T.M., Tsai, T.C., Ko, W.P., Chang, K.J. and Hii, C.H. (2016). Supercritical fluid extract of spent coffee grounds attenuates melanogenesis through down regulation of the PKA, PI3K/Akt, and MAPK signalling pathways. *Evidencebased Complementary and Alternative Medicine*, 2016: 5860296.

Hughes, S. R., López-Núñez, J.C. and Jones, M.A. (2014). Sustainable conversion of coffee and other crop wastes to biofuels and bioproducts using coupled biochemical and thermochemical processes in a multi-stage biorefinery concept. *ApplMicrobiolBiotechnol*;98(20): 8413-31.

ICO. (2021) Crop year production by country

Ilyasoglu, H. and El, S. N. (2014). Nanoencapsulation of EPA/DHA with sodium caseinate-gum Arabic complex and its usage in the enrichment of fruit juice. *Lebensmittel-Wissenschaft und -Technologie- Food Science and Technology*, 56(2), 461–468. <https://doi.org/10.1016/j.lwt.2013.12.002>.

- Irfanoglu, Zeynep Burcu, Baldos, Uris Lantz, Hertel, Thomas, van der Mensbrugge and Dominique. (2014). Impacts of Reducing Global Food Loss and Waste on Food Security, Trade, GHG Emissions and Land Use.
- Kader, A. A. (2005). Increasing food availability by reducing postharvest losses of fresh produce. *Acta Hort.* 682, 2169–2175. [CrossRef]
- Kalogeropoulos, N., Chiou, A., Pyriochou, V., Peristeraki, A. and Karathanos, V.T. (2012). Bioactive phytochemicals in industrial tomatoes and their processing byproducts. *LWT-Food Sci. Technol.* 49, 213–216. [CrossRef]
- Kappel, V. D., Cazarolli, L. H., Pereira, D. F., Postal, B. G., Madoglio, F. A., Buss, Z. S., Flávio Reginatto, F. H. and Silva, F. R. M. (2013). Beneficial effects of banana leaves (*Musa x paradisiaca*) on glucose homeostasis: multiple sites of action. *Brazilian Journal of Pharmacognosy*, 23(4), 706-715.
- Kasuya, M. C. M., Luz, J. M. R., Nunes, M. D., Silva, M. de C. S. da, Carvalho, D. R. de, Assunção, L. S. de, Paula, T. de A., Moura, C. and Bento, C. B. P. (2015). Production of Selenium-Enriched mushrooms in coffee husks and use of this colonized residue. coffee in health and disease prevention, [S.L.], p. 301–309. <https://doi.org/10.1016/B978-0-12-409517-5.00033-4>.
- Kaur D., Wani, A. A., Oberoi, D. P. S. and Sogi, D. S. (2008). Effect of extraction conditions on lycopene extractions from tomato processing waste skin using response surface methodology. *Food Chem.*, 108, 711–718. <https://doi.org/10.1016/j.foodchem.2007.11.002>
- Kaur G., Sandal, A. and Dhillon, N. (2017). Lycopene and human health-A review. *Agric. Rev.* 38, 282-289. <https://doi.org/10.18805/ag.R-1741>
- Kaur, G., Uisan, K., Ong, K.L. and Ki Lin, C.S. (2018). Recent Trends in Green and Sustainable Chemistry & Waste Valorisation: Rethinking Plastics in a Circular Economy. *Curr. Opin. Green Sustain.* 9, 30–39. [CrossRef]
- Ketnawa, S., Chaiwut, P., Rawdkuen, S. (2012). Pineapple wastes: a potential source for bromelain extraction. *Food and bioproducts processing*. Doi: 10.1016/j.fbp.2011.12.006
- Kim, J. H. and Kim, M. Y. (2016). The potential use of citrus juice waste as sources of natural phenolic antioxidants. *J. Appl. Pharm. Sci.* 2016,6, 202–205. [CrossRef]
- Kittiphoom, S. and Sutasinee, S. (2013). Mango seed kernel oil and its physicochemical properties. *Int Food Res J.*, 20, 1145-1149
- Kouame, C., Batchep, R. and Kamga, T. (2013). Evaluation des pertes post-recolte dans la chaîne de production et de commercialisation des légumes feuilles traditionnels à yaounde (Cameroun). *Agronomie Africaine.* 25, 61–70.
- Koubala, B. B., Kansci, G., Garnier, C., Thibault, J. F. and Ralet, M. C. (2013). Physicochemical properties of dietary fibres prepared from ambarella (*Spondias cytherea*) and mango (*Mangifera indica*) peels. *Food Bioprocess Tech.* 6: 591-597.
- Kourmentza, C., Tsafrakidou, P. and Kornaros, M. (2018). Spent coffee grounds make much more than waste: exploring recent advances and future exploitation strategies for the valorization of an emerging food waste stream. *Journal of Cleaner Production* 172, 980–992
- Kughur, P.G., Iornenge, G.M. and Ityonongu, B.E. (2015). Effects of postharvest losses on selected fruits and vegetables among small-scale farmers in Gboko local government area of Benue State, Nigeria. *Int. J. Innov. Sci. Res.* 19, 201–208.

- Kumer, P., Kumer, S., Joshi, L. (2015). Socio-economic and environmental implication of agricultural residue burning: A case study of Punjab India. *Springer Briefs Environ Sci.*, 4, 13-25. Doi: <https://doi.org/10.1007/978-81-322-2014-5-2>
- Kung, C. C., Kong, F. and Choi, Y. (2015). Pyrolysis and biochar potential using crop residues and agricultural wastes in China. *Ecol. Ind.*, 51, 139-145.
- Lau, B. F., Kong, K. W., Leong, K. H., Sun, J., He, X., Wang, Z., Mustafa, M. R., Ling, T. C. and Ismail, A. (2020). Banana inflorescence: Its bio-prospects as an ingredient for functional foods. *Trends in Food Science and Technology*, 97, 14-28.
- Lavanya, M. N., Venkatachalapathy, N. and Manickavasagan, A. (2017). Physicochemical characteristics of rice bran. In Brown rice, Manickavasagan A., Santhakumar C. and Venkatachalapathy N. (eds), 79-90. Doi: 10.1007/978-3-319-59011-0
- Lescot, T. (2017). La diversité génétique des bananiers. *Fruitrop Focus Banane* 208–212
- Li, T., Shen, P., Liu, W., Liu, C., Liang, R., Yan, N. and Chen, J. (2014). Major Polyphenolics in Pineapple Peels and their Antioxidant Interactions. *International Journal of Food Properties*, 17(8), 1805-1817.
- Lin, C. S. K., Pfaltzgraff, L. A., Herrero-Davila, L., Mubofu, E. B., Abderrahim, S., Clark, J. H., Koutinas, A. A., Kopsahelis, N., Stamatelatou, K., Dickson, F., Thankappan, S., Mohamed, Z., Brocklesby, R. and Luque, R. (2013). Food Waste as a Valuable Resource for the Production of Chemicals, Materials and Fuels: Current Situation and Global Perspective. *Energy Environ. Sci.*, 6, 426-464, doi: 10.1039/c2ee23440h
- Lipinski, B., Hanson, C., Lomax, J., Kitinoja, L., Waite, R. and Searchinger, T. (2013). Reducing food loss and waste. Installment 2 of “Creating a Sustainable Food Future”. Working Paper. Washington, DC, World Resources Institute ([http://www.unep.org/pdf/WRI\\_UNEP\\_Reducing\\_Food\\_Loss\\_and\\_Waste.pdf](http://www.unep.org/pdf/WRI_UNEP_Reducing_Food_Loss_and_Waste.pdf)).
- Lopez-Barrera, D. M., Vazquez-Sanchez, K., Loarca-Pina, M.G.F. and Campos-Vega, R. (2016). Spent coffee grounds, an innovative source of colonic fermentable compounds, inhibit inflammatory mediators in vitro. *Food Chemistry* 212, 282–290
- Løvdal, T., Van Droogenbroeck, B., Eroglu, E. C., Kaniszewski, S., Agati, G., Verheul, M. and Skipnes, D.(2019). Valorization of Tomato Surplus and Waste Fractions: A Case Study Using Norway, Belgium, Poland, and Turkey as Examples. 8, 229. [CrossRef] [PubMed]
- Luque, R. and Clark, J. H. (2013). Valorization of food residues: Waste to wealth using green chemical technologies. *Sustainable Chemical Processes*, 1(10), 1-3.
- Mahato, N., Sharma, K., Sinha, M. and Cho, M. H. (2018). Citrus waste derived nutraceuticals for health benefits: Current trends and future perspectives. *J. Funct. Foods*. 40, 307–316. [CrossRef]
- Maina, S., Kachrimanidou, V. and Koutinas, A. (2017). A Roadmap towards a Circular and Sustainable Bioeconomy through Waste Valorization. *Curr. Opin. Green Sustainable Chem.*, 8, 18-23, doi: 10.1016/j.cogsc.2017.07.007
- Majid, R. A., Mohammad, A. W. and May, C. Y. (2012). Properties of residual palm pressed fiber oil. *Journal of Oil Palm Research*, 24, 1310-1317.
- Malviya, R., Srivastava, P., Bansal, M. and Sharma, P. K. (2010). Mango peel pectin as a super disintegrating agent. *Journal of Scientific & Industrial Research* 69: 688-690.
- Manrich, A., Moreira, F. K. V., Otoni, C. G., Lorevice, M. V., Martins, M. A. and Mattoso, L. H. C. (2017). Hydrophobic edible films made up of tomato cutin and pectin. *Carbohydr. Polym.*, 164, 83–91. <https://doi.org/10.1016/j.carbpol.2017.01.075>
- Maria, D. and Freire, G. (2010). Sustainability and environmental issues in food processing

- fiomnuala. *Food Processing: Principles and Applications*, 2, 685-705.
- Markus, M. B., Bolwig, S., Hansen, T. and Tanner, A. N. (2019). Theoretical perspectives on innovation for waste valorisation in the bioeconomy. Doi: 10.4324/9780429460289-3
- Marn, F. R., Soler-Rivas, C., Benavente-Garcia, O., Castillo, J. and Perez-Alvarez, J. (2007). *Food Chem.* 100, 736–741.
- Martínez-Cámara, S., Rubio, S., Del Río, H., Rodríguez-Sáiz, M. and Barredo, J.-L. (2018). Lycopene Production by Mated Fermentation of *Blakeslea trispora*. In: Barreiro C., Barredo JL. (eds) *Microbial Carotenoids. Methods in Molecular Biology*, vol 1852. Humana Press, New York, NY. [https://doi.org/10.1007/978-1-4939-8742-9\\_15](https://doi.org/10.1007/978-1-4939-8742-9_15)
- Massaya, J., Pereira, A. P., Mills-Lampsey, B., Benjamin, J. and Chuck, C.J. (2019). Conceptualization of a spent coffee grounds biorefinery: A review of existing valorization approaches. *Food and Bioproducts Processing* 118, 149–166
- Mata, T. M., Martins, A. A. and Caetano, N. S. (2018). Bio-refinery approach for spent coffee grounds valorization. *Bioresource Technology Reports* 247, 1077–1084
- Matos, A.T. (2015). Tratamento e resíduos após colheita do café. F.M. Borem (Ed.), *Pós-Colheita Do Café*, Editora UFLA, Lavras (2015), pp. 160-202
- Mazzutti, S., Pedrosa, R. C., Salvador, F. S. R. (2020). Green processes in Foodomics. Supercritical fluid extraction of bioactives. Reference module in food science, *Elsevier*, 1-19, doi: [10.1016/b978-0-08-100596-5.22816-3](https://doi.org/10.1016/b978-0-08-100596-5.22816-3)
- Min, B., McClung, A. M. and Chen, M. H. (2011). Phytochemicals and antioxidant capacities in rice brans of different colour. *J Food Sci*, 76, 117-126. Doi: <https://doi.org/10.1111/j.1750-3841.2010.01929.x>
- Mirabella, N., Castellani, V. and Sala, S. (2014). Current options for the valorization of food manufacturing waste: a review. *Journal of Cleaner Production*, 65, 28-41. Doi: <http://dx.doi.org/10.1016/j.jclepro.2013.10.051>
- Mohapatra, D., Mishra, S. and Sutar, N. (2010). Banana and its by-product utilisation: an overview. *Journal of Scientific & Industrial Research*, 69, 323-329.
- Mohd Ali, M., Hashim, N., Abd Aziz, S. and Lasekan, O. (2020). Pineapple (*Ananas comosus*): A comprehensive review of nutritional values, volatile compounds, health benefits, and potential food products. *Food Res. Int.* 137, 109675. [CrossRef] [PubMed]
- Mohdal, A. A. A., Sarhan, M. A., Mahmoud, A., Ramadan, M. F. and Smetanska, I. (2010). Antioxidant efficacy of potato peels and sugar beet pulp extracts in vegetable oils protection. *Food Chemistry*, 123(4), 1019-1026. Doi: <https://doi.org/10.1016/j.foodchem.2010.05.054>
- Mua, R., Honga, X., Nia, Y., Lia, Y., Panga, J., Wang, Q., Xiaoc, J. and Zheng, Y. (2019). Recent trends and applications of cellulose nanocrystals in food industry. *Trends in Food Science and Technology*, 93, 136-144.
- Muchiri, D., Mahungu, S. and Gituanja, S. (2012). Studies on Mango (*Mangifera indica*, L.) Kernel Fat of Some Kenyan Varieties in Meru. *Journal of the American Oil Chemists' Society*, 89, 1567-1575.
- Murthy, P. S. and Naidu, M. M. (2012). Sustainable management of coffee industry by-products and value addition – A review. *Resources, Conservation and Recycling* 66, 45–58
- Mussatto, S. I., Carneiro, L. M., Silva, J. P. A., Roberto, I. C. and Teixeira, J. A. (2011a.) A study on chemical constituents and sugars extraction from spent coffee grounds. *Carbohydrate Polymers* 83(2), 368–374
- Mussatto, S. I., Machado, E. M. S., Martins, S. and Teixeira, J. A. (2011b). Production,

- composition, and application of coffee and its industrial residues. *Food and Bioprocess Technology* 4(5), 661–672
- Mussatto, S. L., Ballasteros, L. F., Martins, S. and Teixeira, J. A. (2011c). Extraction of antioxidant phenolic compounds from spent coffee grounds. *Separation and Purification Technology* 83, 173–179
- Musumeci, L., Maugeri, A., Cirimi, S., Lombardo, G.E., Russo, C., Gangemi, S., Calapai, G. and Navarra, M. (2020). Citrus fruits and their flavonoids in inflammatory bowel disease: An overview. *Nat. Prod. Res.* 34, 122–136. [CrossRef]
- Nayak, A. and Bhushan, B. (2019). An overview of the recent trends on the waste valorization techniques for food wastes. *J. Environ. Manag.* 2019, 233, 352–370. [CrossRef]
- Naziri, E., Nenadis, N., Mantzouridou, F. T. and Tsimidou, M.Z. (2014). Valorization of the major agrifood industrial by-products and waste from Central Macedonia (Greece) for the recovery of compounds for food applications. *FoodRes. Int.* 65, 350–358. [CrossRef]
- Neoh, B. K., Thang, Y. M., Zain, M. Z. M. and Junaidi, A. (2011). Palm pressed fibre oil: A new opportunity for premium hardstock? *International Food Research Journal*, 18, 769-773.
- Ngnikam, E. (2000). Evolution Environnementale et Economique du système de gestion des déchets solides municipaux: Analyse du cas de Yaounde au Cameroun. LAEPSI, Lyon. Institute National des Sciences Applique de Lyon.
- Nguyen, Q. A., Cho, E. J., Lee, D. S. and Bae, H. J., (2019). Development of an advanced integrative process to create valuable biosugars including manno-oligosaccharides and mannose from spent coffee grounds. *Bioresource Technology* 272, 209–216
- Ngwasiri, P. N., Adanmengwi, V. A., Ambindei, W. A., Ngangmou, N. T., Fonmboh, D. J., Ngwabie, N. M., Ngassoum, M. B. and Aba, E. R. (2021). Effect of Traditional Process Methods on the Physicochemical and Functional Properties of a Traditional Food Salt (Nikkih) Obtained from Waste Biomass Peels of *Musa paradisiaca* and *Musa acuminata*. *Hindawi International Journal of Chemical Engineering*, 1-11. doi: <https://doi.org/10.1155/2021/1995683>
- Njukwe, E., Onadipe, O., Kirscht, B. and Mbairanodji, A. (2014). Cassava processing among small-holder farmers in Cameroon: Opportunities and challenges. *Int. J. Agric. Pol. Res.*, 2, 113-124.
- Novoa-Muñoz, J., Simal-Gándara, J., Fernandez-Calviño, D., López-Periago, E. and Arias-Estévez, M. (2008). Changes in soil properties and in the growth of *Lolium multi florum* in acid soil amended with a solid waste from wineries. *Bioresour. Technol.* 99, 6771–6779. doi: 10.1016/j.biortech.2008.01.035
- Ogidi, I. A., Wariboko, C. and Alamene, A. (2018). Appraisal of some chemical elements of plantain (*musa paradisiaca*) cultivars in bayelsa state, nigeria. 6(4), 1-31.
- Okunade, A. F., Agboola, T. L., Ogunsina, B. S., Salau, A. O. and Salami, A. (2019). Physicochemical properties of four selected food spices of Nigerian origin: Alligator pepper, Black pepper, Ginger and Cloves. *Ife Journal of Technology*, 26(1): 7-12.
- Okwu, D. E. (2004). Phytochemicals vitamins content of indigenous spices of South Eastern Nigeria. *Journal of Sustainable Agricultural Environment*, 6: 30-34.
- Oliveira, L. S. and Franca, A. S. (2015). An overview of the potential uses for coffee husks. coffee in health and disease prevention, p. 283–291. 10.1016/B978-0-12-409517-5.00031-0 Google Scholar
- Oluwaseun, K., Adeoye, G. O. and Coker, A. O. (2006). Palm kernel waste management through composting and crop production. *Journal of Environmental Health Research*, 5.

- Ominski, K., McAllister, T., Stanford, K., Mengistu, G., Kebebe, E. G., Omonijo, F., ...Wittenberg, K. (2021). Utilization of by-products and food waste in livestock production systems: a Canadian perspective. *Animal Frontiers*, 11(2), 55–63. Doi:10.1093/af/vfab004
- Onwuka, U. N. and Okala, O. (2003). Effects of selected salts on the cooking time, protein content and sensory properties of African yam beans and cowpeas. *Food Serv Technol.* 3:3–7.
- Oyegbado, C. O., Iyagba, E. T. and Ofor, O. J. (2004). Solid soap production using plantain peels ash as source of alkali. *J. Applied Sci. and Environmental Management.* 6: 73-77.
- Oosterveld, A., Harmsen, J.S., Voragen, A.G.J. and Schols, H.A. (2003). Extraction and characterization of polysaccharides from green and roasted *Coffea arabica* beans, *Carbohydrate Polymers*, 52, (3), 285-296
- Oreopoulou, V. and Tzia, C.(2007). Utilization of plant by-products for the recovery of proteins, dietary fibers, antioxidants, and colorants. In: Oreopoulou, V., Russ, W. (Eds.), *Utilization of By-Products and Treatment of Waste in the Food Industry*. Springer Science+Business Media, New York, 209-232.
- Osano, A. M., Okong'o, E. R., Oyaro, N. and Kiptoo, J. (2013). Compositional and Structural Characterization of Three Basic Indigenous Salts Used in Kenya: A Case Study of "Ebara", "Magadi" and "Lebek" Crystalline Salts. *Int J Adv Res Technol.* 2:118–23.
- Özdestan, O., Erol, T. and Acar, B. (2014). Phytosterols in rice bran and usage of rice bran in food industry. *FOODBALT*, 19-23.
- Palafox-Carlos, H., Ayala-Zavala, F. and González-Aguilar, G. A. (2010). The role of dietary fiber in the bioaccessibility and bioavailability of fruit and vegetable antioxidants. *J Food Sci.*, 76(1), 6-15.
- Pandey, S.K. (2008). Potato research priorities in Asia and the Pacific region. In *Proceedings of the RAPPublication (FAO); FAO/RAP: Bangkok, Thailand.*
- Pandya, D. (2017). Standardization of Solvent Extraction Process for Lycopene Extraction from Tomato Pomace. *J. Appl. Biotechnol. Bioen.* 2, 12-16. <https://doi.org/10.15406/jabb.2017.02.00019>
- Pataro, G., Sinik, M., Capitoli, M. M., Donsì, F. and Ferrari, G. (2015). The influence of Post-harvest UV-C and Pulsed Light treatments on quality and antioxidant properties of tomato fruits during storage. *Innov. Food Sci. Emerg. Technol.* 2015, 30, 103–111. [CrossRef]
- Pateiro-Moure, M., Nóvoa-Muñoz, J. C., Arias-Estévez, M., López-Periago, E., Martínez-Carballo, E. and Simal-Gándara, J. (2009). Quaternary herbicides retention by the amendment of acid soils with a bentonite -based waste from wineries. *J. Hazard. Mater.* 164, 769–775. doi: 10.1016/j.jhazmat. 2008.08.071
- Patel, S. (2012). Cereal bran: the next super food with significant antioxidant and anticancer potential. *Med J Nutr Metab*, 5, 91-104. Doi: <https://doi.org/10.1007/s12349-012-0091-1>
- Peshev, D., Mitev, D., Peeva, L. and Peev, G. (2018). Valorization of spent coffee grounds – A new approach. *Separation and Purification Technology* 192, 271–277
- Pezeshk, S., Ojagh, S. M. and Alishahi, A. (2015). Effect of plant antioxidant and antimicrobial compounds on the shelf- life of seafood. A review. *Czech Journal of Food Sciences*, 33(3), 195–203.

- Pinela, J., Oliveira, M. B. P. and Ferreira, I.C.F. (2016). Chapter 3 Bioactive Compounds of Tomatoes as Health Promoters. In *Natural Bioactive Compounds from Fruits and Vegetables as Health Promoters Part II*; Bentham Science Publishers: Sharjah, United Arab Emirates. Volume 2, pp. 48–91.
- Pleissner, D., Qi, Q., Gao, C., Rivero, C. P., Webb, C., Lin, C. S. K. and Venus, J. (2016). Valorization of Organic Residues for the Production of Added Value Chemicals: A Contribution to the Bio-Based Economy. *Biochem. Eng. J.*, 116, 3– 16, doi: 10.1016/j.bej.2015.12.016
- Pujol, D., Liu, C., Gominho, J., Olivella, M. À., Fiol, N., Villaescusa, I. and Pereira, H. (2013). The chemical composition of exhausted coffee waste. *Industrial Crops and Products* 50, 423– 429
- Rabiu, Z., Maigari, U. F., Lawan, U. and Mukhtar, Z. G. (2018). Pineapple Waste Utilization as a Sustainable Means of Waste Management, *Sustainable Technologies for the Management of Agricultural Wastes*, 143-154.
- Rafiq, S., Kaul, R., Sofi, S., Bashir, N., Nazir, F. and Ahmad Nayik, G. (2018). Citrus peel as a source of functional ingredient: A review. *J. Saudi Soc. Agric. Sci.* 17, 351–358. [CrossRef]
- Rame (2018). Oil Palm Empty Fruit Bunches (OPEFB): Existing Utilization and Current Trends Bio Refinery in Indonesia. *E3S Web of Conferences*, 31, 03014. Doi: <https://doi.org/10.1051/e3sconf/20183103014>
- Raquel Madureira, A., Atatoprak, G., Sousa, F., Pullar, R. C. and Pintado, M. (2018). Extraction and characterisation of cellulose nanocrystals from pineapple peel. *Int. J. Food Stud.* 7, 24–33. [CrossRef]
- Rathinavelu, R. and Graziosi, G. (2005). Potential alternative use of coffee wastes and by-products” ICS-UNIDO Project, Available at: <http://www.ico.org/documents/ed1967e.pdf>, 2005. [viewed on 11/09/2016]
- Roda, A. and Lambri, M. (2019). Food uses of pineapple waste and by-products: A review. *Int. J. Food Sci. Technol.* 54, 1009–1017. [CrossRef]
- Roda, A., De Faveri, D.M., Dordoni, R., Cases, E.V., Jáuregui, N. N., Carbonell Barrachina, Á. A., Frutos Fernandez, M. J. and Lambri, M. (2017). Pineapple Wines Obtained from Saccharification of Its Waste with Three Strains of *Saccharomyces cerevisiae*. *J. Food Process.* 41, e13111. [CrossRef]
- Rodsamran, P. and Sothornvit, R. (2019). Preparation and characterization of pectin fraction from pineapple peel as a natural plasticizer and material for biopolymer film. *Food Bioprod. Process.* 118, 198–206. [CrossRef]
- Roldán-Gutiérrez, J. M. and Luque de Castro, M.D. (2007). Lycopene: The need for better methods for characterization and determination. *Trends Anal. Chem.* 26, 163–170. [CrossRef]
- Rossini, G., Toscano, G., Duca, D., Corinaldesi, F., Foppa Pedretti, E. and Riva, G. (2013). Analysis of the characteristics of the tomato manufacturing residues finalized to the energy recovery. *Biomass Bioenergy.* 51, 177–182. [CrossRef]
- Sabu, A., Pandey, A., Daud, M. J. and Szakacs, G. (2005). Tamarind seed powder and palm kernel cake: two novel agro residues for the production of tannase under solid state fermentation by *Aspergillus niger* ATCC 16620. *Bioresource Technology*, 96, 1223-1228. doi: 10.1016/j.biortech.2004.11.002

- Sahin, S. and Elhuseein, E. A. A. (2018). Valorization of a biomass: phytochemicals in oilseed by-products. *Phytochem Rev*, 17, 657-668. Doi: <https://doi.org/10.1007/s11101-018-9552-6>
- Saidou, H., Hamzaoui, A. H. and Mnif, A. (2015). Insoluble Content, Ionic Composition, Density, and X-Ray Diffraction Spectra of 6 Evaporites from the Niger Republic. *J Appl Chem*. 10.
- Saini, R. K. and Keum, Y. S. (2018). Carotenoid extraction methods: A review of recent developments. *FoodChem*. 2018,240,90–103. [CrossRef]
- Salemdeeb and Al-Tabbaa. (2015). A hybrid life cycle assessment of food waste management options: a UK case study Presented at the 2015 ISWA World Congress, International Solid Waste Association, Antwerp, Belgium.
- Schieber, A. and Aranda, M. D. (2009). Potato peels: A source of nutritionally and pharmacologically interesting compounds – A review. *Global Science Book*, 3, 23–29.
- Sepelev, I. and Galoburda, R. (2015). Industrial Potato Peel Waste Application in Food Production: A Review. *Research for rural development*, 1, 130-136.
- Sepúlveda, L., Romani, A., Aguilar, C. N. and Teixeira, J. (2018). Valorization of pineapple waste for the extraction of bioactive compounds and glycosides using autohydrolysis. *Innov. Food Sci. Emerg. Technol.* 47, 38–45.
- Serna-Loaiza, S., Miltner, A., Miltner, M. and Friedl, A. (2019). A Review on the Feedstocks for the Sustainable Production of Bioactive Compounds in Biorefineries. *Sustainability*, 11, 6765, doi: 10.3390/su11236765
- Shang, Y. F., Xu, J. L., Lee, W. J. and Um, B.H. (2017). Antioxidative polyphenolics obtained from spent coffee grounds by pressurized liquid extraction. *South African Journal of Botany* 109, 75–80
- Sharma, K., Mahato, N., Cho, M. H. and Lee, Y. R. (2017). Converting citrus wastes into value-added products: Economic and environmentally friendly approaches. *Nutrition*. 34, 29–46. [CrossRef] [PubMed]
- Sharma, M., Usmani, Z., Gupta, V. K. and Bhat, R. (2021). Valorization of fruits and vegetable wastes and by-products to produce natural pigments. *Crit. Rev. Biotechnol.* 41, 535–563. [CrossRef] [PubMed]
- Shomodder, A., Hossain, M. M., Das, P. C., Ranganathan, T. V. and Mazumder, M. A. R. (2021). Effect of Banana Leaf Extract, Pumpkin Seed Extract and Bee Honey Treatment on the Inhibition of Browning of Fresh Cut Guava. *The Journal of Agricultural Sciences, Sri-Lanka*, 16(2), 203-214.
- Siaka, D. (2014). Potential of mango (*Mangifera indica*) seed kernel as feed ingredient for poultry: A review. *World Poultry Science Journal*, 70, 279-288.
- Singh, P., Baisthakur, P. and Yemul, O. S. (2020). Synthesis, characterization and application of crosslinked alginate as green packaging material. *Heliyon* 6, e03026. [CrossRef] [PubMed]
- Sonia, N. S., Mini, C. and Geethalekshmi, P. R. (2016). Vegetable peels as natural antioxidants for processed foods – A review. *Agricultural Reviews*, 37(1), 35–41.
- Steiner, C., Das, K. C., Nathan, M. and Lakly, D. (2010). Reducing nitrogen loss during poultry litter composting using biochar. *J Environ Qual* 39:1236–1242. <https://doi.org/10.2134/jeq2009.0337>

- Stoica, R. M., Tomulescu, C., Cășărică, A. and Soare, M. G. (2018). Tomato by-products as a source of natural antioxidants for pharmaceutical and food industries—A mini-review. *Sci. Bull. Ser. F Biotechnol.* 2018, 22, 200–204.
- Strati, I.F. and Oreopoulou, V. (2014). Recovery of carotenoids from tomato processing by-products e a review. *Food Res. Int.* 2014, 65, 311–321. [CrossRef]
- Strati, I.F. and Oreopoulou, V. (2011). Process optimisation for recovery of carotenoids from tomato waste. *Food Chem.* 129, 747–752. [CrossRef]
- Stuart, T. (2009). *Waste: Uncovering the global food scandal*. London, W.W. Norton Co.
- Sudiyani, Y., Styarini, D., Triwahyuni, E., Sudiarmanto, Sembiring, K. C., Aristiawan, Y., Abimanyu, H. and Han, M. H. (2013). Utilization of biomass waste empty fruit bunch fiber of palmoil for bioethanol production using pilot-scale unit. *Energy Procedia*, 32, 31-38. doi: 10.1016/j.egypro.2013.05.005
- Sundalian, M., Larissa, D. and Suprijana, O. (2021). Contents and Utilization of Palm Oil Fruit Waste. *Biointerface Research in Applied Chemistry*, 11(3), 10148-10160. Doi: <https://doi.org/10.33263/BRIAC113.1014810160>
- Szabo, K., Cătoi, A. F. and Vodnar, D.C. (2018). Bioactive Compounds Extracted from Tomato Processing by-Products as a Source of Valuable Nutrients. *Plant. Foods Hum. Nutr.* 73, 268–277. [CrossRef]
- Szabo, K., Dulfb, F.V., Diaconeasa, Z. and Vodnara, D.C. (2019). Antimicrobial and antioxidant properties of tomato processing byproducts T and their correlation with the biochemical composition. *Food Sci. Technol.* 116, 108558. [CrossRef]
- Szabo, K., Diaconeasa, Z., Cătoi, A. F. and Vodnar, D.C. (2019). Screening of Ten Tomato Varieties Processing Waste for Bioactive Components and Their Related Antioxidant and Antimicrobial Activities. *Antioxidants*. 8, 292. [CrossRef]
- Tamrat, T., and Bruce S. (2016). *Valorisation of mango seed via extraction of starch*, Organic Waste book valorisation, South Africa.
- Tan, Y. (2006). By-products of palm oil extraction and refining. *Oilseeds and fats, Crops and Lipids*, 13(1), 9-11.
- Tata, P.I., Afari-Sefa, V., Ngome, A.F. and Billa, S.F. (2016). Policy and institutional frameworks impacting on vegetable seed production and distribution systems in Cameroon. *J Crop Improve* 30:196–216. <https://doi.org/10.1080/15427528.2016.1141134>
- Teh, S. S., Lau, H. L. N. and Mah, S. H. (2019). Palm-Pressed Mesocarp Fibre Oil as an Alternative Carrier Oil in Emulsion. *J. Oleo Sci.*, 68(8), 803-808. doi: 10.5650/jos.ess19098.
- Tesfaye, T., Begam, R., Sithole, B. B. and Shabaridharan, K. (2015). Dyeing Cotton with Dyes Extracted from Eucalyptus and Mango Trees. *The International Journal of Science and Technoledge*, 3(8), pp. 310.
- Thorat, R. L. and Bobade, H. P. (2018). Utilization of banana pseudo-stem in food applications. *International Journal of Agricultural Engineering*, 11, 86-89.
- Tian, T., Freeman, S., Corey, M., German, J. B. and Barile, D. (2017). Chemical characterization of potentially prebiotic oligosaccharides in brewed coffee and spent coffee grounds. *Journal of Agricultural and Food Chemistry* 65(13), 2784– 2792
- Tibolla, H., Pelissari, F., Martins, J., Vicente, A. and Menegalli, F. (2018). Cellulose nanofibers produced from banana peel by chemical and mechanical treatments: Characterization and cytotoxicity assessment. *Food Hydrocoll.* 75, 192–201. [Google Scholar] [CrossRef]

- Tirouchelvame, D., Pragalyaashree, M. M. and Dayanand, P. (2019). Process Development for the Value Addition of Banana Central Core (Pseudo Stem). *International Journal of Recent Technology and Engineering (IJRTE)*, 8(4), 1055-1060.
- Tiwari, B. K., Valdramidis, V. P., O'Donnell, C. P., Muthukumarappan, K., Bourke, P. and Cullen, P. J. (2009). Application of natural antimicrobials for food preservation. *Journal of Agricultural and Food Chemistry*, 57(14), 5987–6000. <https://doi.org/10.1021/jf900668n>
- Tobias, I. N. E., Enebaku, U. E. and Ogueke, C. (2014). Waste to Wealth-Value Recovery from Agrofood Processing Wastes Using Biotechnology: A Review. *British Biotechnology Journal*, 4(4), 418-481.
- Torres-Leon, C., Ramirez-Guzman, N., Londono-Hernandez, L., Martinez-Medina, G. A., Diaz-Herrera, R., Navarro-Macias, V., Alvarez-Perez, O. B., Picazo, B., Villarreal-Vazquez, M., Ascacio-Valdes, J. and Aguilar (2018). Food waste and by-products: An opportunity to minimize malnutrition and hunger in developing countries. *Front. Sustain. Food Syst.*, 2(52), 1-17.
- Tortoe, C., Johnson, P. N. T., Slaghek, T., Miedema, M. and Timmermans, T. (2013). Physicochemical, Proximate and Sensory Properties of Pineapple (*Ananas sp.*) Syrup Developed from Its Organic Side-Stream. *Food Nutr. Sci.* 4, 163–168. [CrossRef]
- Tremblay, A., Corcuff, R., Goulet, C., Godefroy, S. B., Doyen, A. and Beaulieu, L. (2020). Valorization of snow crab (*Chionoecetes opilio*) cooking effluents for food applications. *Journal of the Science of Food and Agriculture*, 100(1), 384–393. <https://doi.org/10.1002/jsfa.10066>.
- Tsouko, E., Alexandri, M., Fernandes, K. V., Freire, D. M. G., Mallouchos, A. and Koutinas, A. A. (2018). Extraction of Phenolic Compounds from Palm Oil Processing Residues and Their Application as Antioxidants. *Food Technology and Biotechnology*, 57(1), 29-38. doi: <https://doi.org/10.17113/ftb.57.01.19.5784>
- Tuck, C. O., Pérez, E., Horváth, I. T., Sheldon, R. A. and Poliakoff, M. (2012). Valorization of Biomass: Deriving More Value from Waste. *Science*, 338, 695, doi: 10.1126/science.338.6107.604-b
- Umaru, F. F., Esedafe, W. K., Obidah, J. S., Akinwotu, O. and Danba, E. (2015). Production of Vinegar from Pineapple Peel Wine Using *Acetobacter* Species. In Proceedings of the 3rd International Conference on Biological, Chemical & Environmental Sciences (BCES-2015), Kuala Lumpur, Malaysia, 21–22 September 2015.
- Upadhyay, A., Lama, J. P. and Tawata, S. (2013). Utilization of Pineapple Waste: A Review. *J. Food Sci. Technol. Nepal*. 6, 10–18. [CrossRef]
- Upadhyay, A., Lama, J. P. and Tawata, S. (2010). Utilization of Pineapple Waste: A Review. *J. Food Sci. Technol. Nepal*, 6, 10-18.
- Van Dyk, J. S., Gama, R., Morrison, D., Swart, S. and Pletschke, B. I. (2013). Food processing waste: Problems, current management and prospects for utilisation of the lignocellulose component through enzyme synergistic degradation. *Renew. Sustain. Energy*. 26, 521–531. [CrossRef]
- Varzakas, T., Zakynthinos, G. and Verpoort, F. (2016). Plant Food Residues as a Source of 1276 Nutraceuticals and Functional Foods. *Foods*, 5, 1-32. Doi: 10.3390/foods5040088
- Vázquez-Sánchez, K., Martínez-Saez, N., Rebollo-Hernanz, M., Del Castillo, M. D., Gaytán-Martínez, M. and Campos-Vega, R. (2018). In vitro health promoting properties of antioxidant dietary fiber extracted from spent coffee (*Coffea arabica* L.) grounds. *Food*

Chemistry 261, 253–259

- Vermande, P. and Ngnikam, E. (1994). Etude de la gestion et des traitement des ordures menageres de Yaounde (Cameroon), vol. I Actual Situation. Yaounde. Fond Special d'Equipement et d'Intervention Intercommunale (FEICOM). Maire d'arrondissement de Yaounde II, ENSP et l'Universite de Yaounde I.
- Wahyuningsih, K., Iriani, E. S. and Fahma, F. (2016). Utilization of Cellulose from Pineapple Leaf Fibers as Nanofiller in Polyvinyl Alcohol-Based Film. *Indones. J. Chem.*, 16 (2), 181-189.
- Wali, N. (2019). Pineapple (*Ananas comosus*). In Non vitamin and Non mineral Nutritional Supplements; Elsevier: Amsterdam, The Netherlands. pp. 367–373. ISBN 9780128124918.
- Wang, H. Y., Qian, H. and Yao, W. R., (2011). Melanoidins produced by the Maillard reaction: structure and biological activity. *Food Chemistry* 128, 573–584
- Whiting, A. and Azapagic, A. (2014). Life cycle environmental impacts of generating electricity and heat from biogas produced by anaerobic digestion. *Energy*, 70, 181-193, doi: [10.1016/j.energy.2014.03.103](https://doi.org/10.1016/j.energy.2014.03.103)
- Widjaja-Adhi, M. A. K., Ramkumar, S. and von Lintig, J. (2018). Protective role of carotenoids in the visual cycle. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 32, 6305–6315. [CrossRef] [PubMed]
- Widmer, W., Zhou, W. and Grohmann, K. (2010). *Bioresour. Technol.* 2010, 101, 5242– 5249.
- Wills, R., McGlasson, B., Graham, D. and Joyce, D. (2004). *Postharvest: Introduction to the Physiology and Handling Fruits, Vegetables and Ornamentals*, 4th ed.; University of New South Wales Press Ltd.: Sydney, Australia. 262p
- World Food Program. (2017). *Cameroon: Comprehensive Food Security and Vulnerability Analysis (CFSVA); Vulnerability Analysis Unit (VAM) (OSZAF)*: Rome, Italy, 183p.
- World Processing Tomato Council (WPTC). (2020). *The 2020 Processed Tomato Yearbook*.. Available online: <http://www.tomatonews.com/pdf/yearbook/2020/index.html#48> (accessed on 30 June 2021).
- Wunder, S., Keighley, M. F., Martin, H. G., *et al.* (2018). Food waste prevention and valorisation: Relevant EU policy areas. Report of the REFRESH project, D3.3 review of EU policy areas with relevant impact on food waste prevention and valorization.
- Wyman, Ch.E. (2001). Twenty years of trials, tribulations, and research progress in bioethanol technology. *Appl. Biochem. Biotech.* 91: 5-21.
- Xiong, X., Yu, I. K. M., Tsang, D. C. W., Bolan, N. S., SikOk, Y., Igalavithana, A. D., Kirkham, M. B., Kim, K. H. and Vikrant, K. (2019). Value-Added Chemicals from Food Supply Chain Wastes: State-of-the-Art Review and Future Prospects. *Chem. Eng. J.*, 375, 121983. Doi: [10.1016/j.cej.2019.121983](https://doi.org/10.1016/j.cej.2019.121983)
- Xu, H., Wang, W., Liu, X., Yuan, F. and Gao, Y. (2015). Antioxidative phenolics obtained from spent coffee grounds (*Coffea Arabica* L.) by subcritical water extraction. *Industrial Crops and Products* 76, 946–954
- Yadav, M. P., Hicks, K. B., Johnston, D. B., Hotchkiss Jr., A. T., Chau, H. K. and Hanah, K. (2015). Production of Bio-based Fiber Gums from the Waste Streams Resulting from the Commercial Processing of Corn Bran and Oat Hulls. *Food Hydrocolloids*. Doi: [10.1016/j.foodhyd.2015.02.017](https://doi.org/10.1016/j.foodhyd.2015.02.017)

- Yashin, A., Yashin, Y., Xiaoyan, X. and Nemzer, B. (2017). Antioxidant Activity of Spices and Their Impact on Human Health: A Review. Gordon and Breach Science Publishers, New York, pp. 1-31.
- Zainal Alam, M. N. H., Adrus, N., Abdul Wahab, M. F., Kamaruddin, M.J. and Sani, M.H. (2020). Utilization of Agro-Waste as Carbon Source for Biohydrogen Production: Prospect and Challenges in Malaysia. In Valorisation of Agro-industrial Residues—Volume I: Biological Approaches; Springer: Cham, Switzerland. pp. 131–147.
- Zarei, M., Ebrahimpour, A., Abdul-Hamid, A., Anwar, F. and Saari, N. (2012). Production of Defatted Palm Kernel Cake Protein Hydrolysate as a Valuable Source of Natural Antioxidants. *Int. J. Mol. Sci.*, 13, 8097-8111. doi:10.3390/ijms13078097
- Zenebe, W., Ali, M., Derbew, B., Zekarias, S. and Adam, B. (2015). Assessment of banana postharvest handling practices and losses in Ethiopia. *J. Biol. Agric. Healthcare*. 5, 82–96.
- Zhao, M., Lan, Y., Cui, L., Monono, E., Rao, J. and Chen, B. (2019). Formation, characterization, and potential food application of rice bran wax oleogels: Expeller-pressed corn germ oil versus refined corn oil. *Food Chemistry*, 125704. Doi: <https://doi.org/10.1016/j.foodchem.2019.125704>
- Zhao, S., Yao, S., Ou, S., Lin, J., Wang, Y., Peng, X., Li, A. and Yu, B. (2015). Preparation of ferulic acid from corn bran: Its improved extraction and purification by membrane separation. *Food and Bioproducts Processing*, 92, 309-313. Doi: <http://dx.doi.org/10.1016/j.fbp.2013.09.004>
- Zuorro, A. (2020). Enhanced lycopene extraction from tomato peels by optimized mixed-polarity solvent mixtures. *Molecules* 25. <https://doi.org/10.3390/molecules25092038>
- Zuorro, A. and Lavecchia, R. (2012). Spent coffee grounds as a valuable source of phenolic compounds and bioenergy. *Journal of Cleaner Production* 34, 49–56
- Zuorro, A., Fidaleo, M. and Lavecchia, R. (2011). Enzyme-assisted extraction of lycopene from tomato processing waste. *Enzym. Microb. Technol.* 49, 567–573.