

## Original Research Article

### Is *MTHFR*C677T Gene Polymorphism Associated with Hypertension in Nigerians?

**Running Title:** Association of *MTHFR*C677T Gene Polymorphism with hypertension in Nigerians.

#### ABSTRACT

**Background:** Essential hypertension is very common in Nigeria. The cause is unknown. Genetic factors have been postulated by some authors as a possible risk factor. Such genetic factors include the mutation of methylenetetrahydrofolate reductase (*MTHFR*) gene.

**Aim:** This study aimed to document the allelic and genotype frequencies and distribution among hypertensive and healthy Nigerian population.

#### Materials and Methods

This was a cross-sectional study involving 75 consenting subjects (50 cases and 25 controls) at the Cardiology Clinic of the Lagos University Teaching Hospital, Lagos, Nigeria. Structured interviewer administered questionnaire was used to obtain socio-demographic and clinical history of subjects. About 5mls of venous blood was collected from each subject by a trained phlebotomist into EDTA bottle and stored at 4°C until ready for analysis. Genomic DNA extraction was done after which polymerase chain reaction was carried out. This was followed by restriction enzyme digestion and agarose gel electrophoresis. The digestion products were then visualized with SYBR Safe (Monitagen) using Sygene bio-imaging system.

#### Results

When compared with hypertensive subjects, normotensive subjects had more CC (84% vs 74%) and CT (16% vs 12%) genotypes. Hypertension was significantly associated with mutant *MTHFR* genotypes (14% vs 0.0%) (OR = 3.995, 95% C.I: 1.101- 10.034; p=0.033). Except for age (OR= 1.771, 95% CI: 1.036 – 3.029; p=0.037), smoking (OR= 0.000; p=0.999), alcohol consumption (OR= 0.000; p=0.999), and sex (OR= 15.052, 95% CI: 0.196- 115.028; p=0.139) did not attain statistical significance.

**Conclusion:** The 677TT homozygous mutant had the highest risk of association with hypertension in Nigerians.

**Key Words:** Hyperhomocysteinemia; mutant; polymorphism; hypertension

## INTRODUCTION

Many multi-organ complications arise from hypertension. The prevalence of hypertension is said to be lowest in India (males: 3.4%; females: 6.8%) and highest in Poland (males: 68.9%; females: 72.5%)[1] A Nigerian review on hypertension published in 2015 revealed a prevalence of 28.9% (males 29.5%; females 25.0%) with a geographical prevalence of 30.6% in the urban population and 26.4% among the rural dwellers, respectively.[2] In 95% of subjects, the cause is unknown and is referred to as essential hypertension.[3] However, many authors now believe that genetic factors such as the mutation of methylenetetrahydrofolate reductase (*MTHFR*) gene play prominent roles in the pathogenesis of essential hypertension.[4-9]

The *MTHFR* gene has been shown to have many polymorphisms. A study in the Israeli women population reported an increase in the diastolic blood pressure in women with *MTHFR* C677T population irrespective of their blood lipid levels.[10] A recent study also reported the association between *MTHFR* C677T gene polymorphism and essential hypertension which is closely related to the increased level of homocystein (Hcy) [11]

The C677T polymorphism (in which valine substitutes alanine at the 222<sup>nd</sup> codon) is one of the clinically significant mutations. [9] *MTHFR* C677T is thermolabile because its activity decreases when the temperature is above 37°C.[12] The enzyme activity of the *MTHFR* 677TT genotype varies depending on whether it is homozygous (30%) or heterozygous (60%) [7] Whereas some authors have reported that the *MTHFR* C677T mutation did not increase cardiovascular risk, others have shown that the homozygous mutation was linked with diseases such as hypertension and stroke,[5,13] with different prevalences in different parts of the world. [14-16]

This study aimed to document the allelic and genotype frequencies and distribution among hypertensive and healthy Nigerian populations. The study also aimed to evaluate MTHFR 677CT SNP as a possible independent predictor of hypertension in relation to other traditional markers of hypertension such as age, gender, alcohol use and smoking habits among Nigerians.

## **MATERIALS AND METHODS**

### **Study site and population**

This cross-sectional study involving 75 subjects (50 subjects and 25 controls) was carried out at the Cardiology Clinic of the Lagos University Teaching Hospital after obtaining ethical approval from the institutions Ethics and Research Committee with assigned number ADM/DCST/HREC/1889. Twenty three males and 27 females were recruited as study group with 11 males and 14 females serving as controls.

### **Inclusion criteria**

Male and female subjects diagnosed with primary hypertensive disorders and normotensives of 18 years and above, were included in the study.

### **Exclusion criteria**

The following subjects were excluded from the study: Subjects below 18years; those with clinical or laboratory evidence of secondary hypertension; extradural and subdural haemorrhages. Subjects that survived from TIA or stroke events in cases of blood disease (e.g. leukaemia, polycythaemia vera); brain tumours or brain metastases.

### **Subject Selection, Sample Collection and Handling**

Structured interviewer administered Questionnaires were used to obtain socio-demographic and clinical history of subjects. Verbal and written informed consents were obtained from the subjects after clear explanation of study objectives, benefits and limitations. Subjects were aged 18 years and above with the diagnosis of essential hypertension confirmed by

cardiologists according to the International Society of Hypertension's criteria [systolic blood pressure (SBP)  $\geq$  140 mmHg or diastolic blood pressure (DBP)  $\geq$  90 mmHg]. [17] The hypertensive subjects were attending the Cardiology Clinic of the hospital regularly. Before recruitment and at each visit, blood pressure readings were taken twice in the sitting position with a mercury sphygmomanometer (Accoson MDF800 desk Mercury Thermometer, UK), 30 minutes apart and subjects with mean blood pressure readings above 140/90.

The control group were normotensive subjects (SBP < 140 mmHg or DBP < 90 mmHg) without any family history of diabetes, cardiovascular and thyroid diseases.

Approximately 5mLs of venous blood was collected by a trained phlebotomist into EDTA bottle and was stored at 4°C until ready for analysis.

#### **DNA Extraction and Restriction Enzyme Digestion**

After genomic DNA extraction with the QiagenDNeasy Kit according to the manufacturers' protocols, polymerase chain reaction (PCR) was done, followed by restriction enzyme digestion. A 25 $\mu$ l PCR reaction mixture comprised of 2 $\mu$ l genomic DNA, 16  $\mu$ l PCR water, 0.8 $\mu$ l dNTP (2.5mM), 0.2 $\mu$ l MgCl<sub>2</sub> (1.5mM), 0.5 $\mu$ l 5<sup>1</sup>-AAGATCAGAGCCCCAAAGC-3<sup>1</sup> forward primer, 0.5 $\mu$ l 5<sup>1</sup>- ACTCAGCGAACTCAGCACTC-3<sup>1</sup> reverse primer and 5.0 $\mu$ l Tag DNA polymerase, constituted inside the PCR tube. The genotyping protocol for the detection of the MTHFR C677T polymorphism was adapted from Frosst *al.*[18] The Hinf I restriction digestion and agarose gel electrophoresis were carried out according to standard protocols. The digestion products were visualized with SYBR Safe (Moltagen) using Sygene bio-imaging system.

#### **DATA ANALYSIS**

Data were analysed using the Statistical Package for Social Sciences (SPSS) version 25.0 Chicago, IL. For descriptive statistics,  $\chi^2$  test was used as univariate statistics to compare proportions in the distribution of demographic variables and MTHF C677T genotypes among the subjects in the study, and the student t-test was used to compare mean values of systolic and diastolic blood pressures among the two groups of subjects. Logistic regression model was adopted to describe associations between MTHFR genotypes and some socio-demographic factors with dichotomous response on presence or absence of hypertension among subjects, and determine

independent predictor variables for hypertension in the study. The critical level of significance was set at  $p < 0.05$ .

## RESULTS

### Socio-demographic characteristics of subjects

The socio-demographic characteristics of the study and control subjects with respect to their age range, sex and blood pressure are shown in Table 1. In both groups (study and controls), the age group 60 years and above (study 40%; Control 48%) constituted the bulk of the subjects in this study. This was followed by the age group 50-59 years (20% in both study group and control). Subjects  $\leq 30$  years (10.0%) in the study group and 30-39 years (8.0%) in the control group had the lowest frequency. The blood pressure readings of the study group (systolic  $147 \pm 18$ ; diastolic  $92 \pm 15$ ) were significantly higher than in the control (systolic  $115 \pm 5.8$ ; diastolic  $76 \pm 5.3$ ) ( $p = 0.000$ ).

### Distribution of *MTHFR* C677T genotypes

Normotensive subjects had more CC genotypes than the hypertensive subjects (84% vs 74%). The normotensive group also had more CT genotypes than the hypertensive group (16% vs 12%). There was a significant association of homozygous mutant genotype TT with hypertension; while 7 of the hypertensive subjects had T genotype, none of the normotensive subjects had the TT genotype (14.0% vs 0.0%;  $p = 0.049$ ). Table 2.

### The logistic regression for variables associated with hypertension

In Table 3, hypertension was significantly associated with mutant *MTHFR* genotypes (OR = 3.995, 95% C.I: 1.101- 10.034;  $p = 0.033$ ) among subjects. However, other traditional socio-demographic risk factors for hypertension such as smoking (OR= 0.000;  $p = 0.999$ ), alcohol consumption (Or= 0.000;  $p = 0.999$ ), and sex (OR= 15.052, 95% CI: 0.196- 115.028;  $p = 0.139$ ) did not attain statistical significance except age (OR= 1.771, 95% CI: 1.036 – 3.029;  $p = 0.037$ ) when evaluated with *MTHFR* mutation as a potential risk factor for hypertension among study group.

## DISCUSSION

This case-control study demonstrated that the *MTHFR*C677T gene polymorphism was significantly associated with increased risk of hypertensive disorders in Lagos, South-Western Nigeria. Previously, studies have been conducted to determine the association between *MTHFR* C677T polymorphism and hypertensive disorders, however, the conclusions were largely controversial. In a Moroccan study, the homozygous mutant SNPs for the *MTHFR* gene (677TT/CT) were associated with significant risk of hypertension similar to the findings in this study[5] A meta-analysis report by Kumar *et al.*, 2015 also found significant association between *MTHFR* C677T polymorphism and the risk of developing cerebrovascular accident or ischemic stroke[19]. Other published reports include studies from Spain. [20] Chinese Han [21,22], Taichung, China[23] and Turkey [24]. Yang *et al.*, also reported a strong association of *MTHFR* C677T with the risk of hypertension among Asians, Caucasians and Chinese subjects in a meta-analysis [25] However, few reports have observed no significant association between mutant *MTHFR* genotypes and hypertension. These included studies carried out on Chinese Caucasians [26] and Japanese[27] where the presence of the T allele did not predispose subjects to increased risk of hypertension.

A significant relationship between gender and mutant *MTHFR* C677T was not established in this study. In a similar study carried out in South West Cameroon, which considered *MTHFR* C677T genotype distribution by gender in both hypertensive and normotensive subjects, reported no gender preference in the genotype distribution for the *MTHFR* 677 SNP [28] The *MTHFR* genotype frequencies for CC, CT and TT among control group subjects were 84.0%, 16.0% and 0.0% respectively. The hypertensive group had 74.0%, 12.0%, 14.0 % for CC, CT and TT

respectively thus showing a significant difference in *MTHFR* 677CT genotype distribution between control and hypertensive subjects. (P=0.049). This finding is in agreement with reports from a study by Nasserredine *et al.*, in Morocco as further evidence that the homozygous mutant 677TT was significantly associated with the risk of hypertension in an African population.[5] Markan *et al.* also reported similar observations in the Indian population[29] A number of studies from Sub-Saharan Africa have suggested that genotype frequencies for *MTHFR* mutant variants are extremely low among African populations; suggesting that the C677T allele is less common in African populations than other ethnic groups[28] Therefore, studies indicating marginal elevations in the genotype frequencies of *MTHFR* C677T mutant SNP in a pathologic entity among African populations, should arouse strong suspicion for disease association studies.

**Conclusion:** The 677TT homozygous mutant had a high risk of association with hypertension in Nigerians.

**Recommendation:** *MTHFR* screening should be encouraged early in life or as part of routine health check. This may help identify individuals with increased risk of hypertensive heart diseases, stroke and metabolic syndrome for early life style modifications in those at risk.

**Study Limitation:** Limitation to this was the inability to obtain serum folate levels of the subjects. It was difficult to control all other factors that might have contributed to variations in homocysteine levels among participants.

#### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## REFERENCES

1. Kearney PM, Whelton M, Reynolds K, Whelton PK, and He J. Worldwide prevalence of hypertension: a systematic review. *J Hypertens* 2004; 1: 11-19.
2. Adeloye D, Basquill C, Aderemi AV, Thompson JY, and Obi FA. An estimate of the prevalence of hypertension in Nigeria: a systematic review and meta-analysis. *J Hypertens*. 2015; 33: 230-242.
3. Carretero OA, and Oparil S. Essential Hypertension. Part I: Definition and Etiology. *Circulation*. 2000; 101: 329–335.
4. Alghasham A, Settin AA, Ali A, Dowaidar M, and Ismail H. Association of MTHFR C677T and A1298C gene polymorphisms with hypertension. *Int J Health Sci*. 2012; 6: 3-11.
5. Nassereddine S, Kassogue Y, Korchi F, Habbal R, and Nadifi S. Association of methylenetetrahydrofolate reductase gene (C677T) with the risk of hypertension in Morocco. *BMC Res Notes*. 2015; 8: 775.
6. Akalin A, Alatas O, and Colak O. Relation of plasma homocysteine levels to atherosclerotic vascular disease and inflammation markers in type 2 diabetic subjects. *Eur J Endocrinol*. 2008; 158: 47-52.
7. Robien K, and Ulrich CM. 5,10-Methylenetetrahydrofolate Reductase Polymorphisms and Leukemia Risk. *Am J Epidemiol* 2003; 157: 571–582.
8. Kiseljaković E, Jadrić R, Hasić S, Skenderi F, Resić H, and Winterhalter-Jadrić M. Polymorphism in methylenetetrahydrofolate reductase gene: important role in diseases. *Bosn J Basic Med Sci*. 2008; 8: 165–169.
9. Alam MA, Husain SA, Narang R, Chauhan SS, Kabra M, and Vasisht S. Association of polymorphism in the thermolabile 5, 10-methylene tetrahydrofolate reductase gene and hyperhomocysteinemia with coronary artery disease. *Mol Cell Biochem*. 2008; 310: 111-117.
10. Heifetz EM, Birk RZ. MTHFR C677T polymorphism affects normotensive diastolic blood pressure independently of blood lipids. *Amer J Hypertens*. 2015; 28(3) : 387–392
11. Cai W, Yin L, Yang F, Zhang L, Cheng J. Association between Hcy levels and the CBS844 ins68 and MTHFR C677T polymorphisms with essential hypertension. *Biomed Rep* 2014; 2(6):861-868
12. Li C, Yichao J, Jiabin L, Yueting Z, Qin L, and Tonghua Y. Methylenetetrahydrofolate reductase gene polymorphism and risk of chronic myelogenous leukemia: a meta-analysis. *J BUON*. 2015; 20: 1534-1545.

13. Brattström L, Wilcken DE, Ohrvik J, and Brudin L. Common methylenetetrahydrofolate reductase gene mutation leads to hyperhomocysteinemia but not to vascular disease: the result of a meta-analysis. *Circulation*. 1998;98:2520–2526
14. Castro R, Rivera I, Ravasco P, Camilo ME, Jakobs C, Blom HJ, and de Almeida IT. 5,10-methylenetetrahydrofolate reductase (MTHFR) 677C-->T and 1298A-->C mutations are associated with DNA hypomethylation. *J Med Genet*. 2004; 41: 454-458.
15. Wilcken B, Bamforth F, Li Z, Zhu H, Ritvanen A, Renlund M, Stoll C, Alembik Y, Dott B, Czeizel AE, Gelman-Kohan Z, Scarano G, Bianca S, Ettore G, Tenconi R, Bellato S, Scala I, Mutchinick OM, López MA, de Walle H, Hofstra R, Joutchenko L, Kavteladze L, Bermejo E, Martínez-Frías ML, Gallagher M, Erickson JD, Vollset SE, Mastroiacovo P, Andria G, and Botto LD. Geographical and ethnic variation of the 677C>T allele of 5,10 methylenetetrahydrofolate reductase (MTHFR): findings from over 7000 newborns from 16 areas worldwide. *J Med Genet*. 2003; 40: 619-625.
16. Yang B, Liu Y, Li Y, Fan S, Zhi X, Lu X, Wang D, Zheng Q, Wang Y, Wang Y, and Sun G. Geographical distribution of MTHFR C677T, A1298C and MTRR A66G gene polymorphisms in China: findings from 15357 adults of Han nationality. *PLoS One*. 2013; 8: e57917.
17. Whitworth JA. 2003 World Health Organization (WHO)/International Society of Hypertension (ISH) statement on management of hypertension. *J Hypertens*. 2003; 21: 1983–1992.
18. Frosst P, Blom HJ, Milos R, Goyette P, Sheppard CA, Matthews RG, Boers GJ, den Heijer M, Kluijtmans LA, and van den Heuvel LP . A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nature Genetics*. 1995; 10: 111–113.
19. Kumar A, Kumar P, Prasad M, Sagar R, Yadav AK, Pandit AK, et al. (2015). Association of C677T polymorphism in the methylenetetrahydrofolate reductase gene (MTHFR gene) with ischemic stroke: A meta-analysis. *Neurol. Res*. 2015;37(7):568-77
20. Rodríguez-Esparragón F, Hernández-Perera O, Rodríguez-Pérez JC, Anábitarte A. The effect of methylenetetrahydrofolate reductase C677T common variant on hypertensive risk is not solely explained by increased plasma homocysteine values. *Clin. Exp. Hypertens*. 2003; **25**: 209-220.
21. Liu C, Li H, Qi Q, Lu L, et al. (2011). Common variants in or near FGF5, CYP17A1 and MTHFR genes are associated with blood pressure and hypertension in Chinese Hans. *J. Hypertens*. 2011; **29**: 70-75.

22. Li YY. Methylenetetrahydrofolate reductase C677T gene polymorphism and coronary artery disease in a Chinese Han population: a meta-analysis. *Metabolism*. 2012; **61**: 846-852
23. Lin PT, Cheng CH, Wei JC and Huang YC. Low plasma pyridoxal 5 $\alpha$ -phosphate concentration and MTHFR 677C->T genotypes are associated with increased risk of hypertension. *Int. J. Vitam. Nutr. Res.* 2008; **78**: 33-40
24. Bayramoglu A, Urhan Kucuk M, Guler HI, Abaci O, Kucukkaya Y, Colak E. Is there any genetic predisposition of MMP-9 gene C1562T and MTHFR gene C677T polymorphisms with essential hypertension? *Cytotechnology*. 2015; 67: 115–122.
25. Yang KM, Jia J, Mao LN, Men C, Tang KT, Li YY. Methylenetetrahydrofolate reductase C677T gene polymorphism and essential hypertension: a metaanalysis of 10,415 subjects. *Biomed Rep.* 2014; 2(5):699-708.
26. Javed Y, Fowdar, Marta V. Lason, Attila L. Szvetko, Rodney A. Lea, Lyn R. Griffiths, "Investigation of Homocysteine-Pathway-Related Variants in Essential Hypertension", *Int J Hypertension*. 2012; Article ID 190923, 9 pages.<https://doi.org/10.1155/2012/190923>
27. Nakata Y, Katsuya T, Takami S, Sato N. Methylenetetrahydrofolate reductase gene polymorphism: relation to blood pressure and cerebrovascular disease. *Am. J. Hypertens.* 1998; **11**: 1019-1023 gene polymorphism and essential hypertension in South West Cameroon.; *Genet.*
28. Ghogomu SM, Ngolle NE, Moulion RN, Asa BF. Association between the MTHFR C677T gene polymorphism and essential hypertension in South West Cameroon. *Gen and Mol Res.* 2015; **15**(1): gm.15017462.
29. Markan S, Sachdeva M, Sehrawat BS, Kumari S, Jain S, Khullar M. MTHFR 677CT/MTHFR 1298CC genotypes are associated with increased risk of hypertension in Indians. *Mol Cell Biochem.* 2007; 302:125–1231.

**TABLES**

**Table 1 : Demographic Characteristics of Subjects**

<b>Characteristics</b>	<b>Hypertensive(%)</b>	<b>Normotensive (%)</b>	<b>Frequency</b>	<b><math>\chi^2</math></b>	<b><i>p</i> value</b>
<b>Age Group (Yrs)</b>					
≤30	5(10)	3(12.0)	8	0.972	0.808
30-39	8(16)	2(8.0)	10		
40-49	7(14)	3(12.0)	10		
50-59	10(20)	5(20.0)	15		
≥60	20(40)	12(48.0)	32		
<b>Sex</b>					
Male	23(67.6)	11(32.4)	34(100)	0.246	0.619
Female	27(65.9)	14(34.1)	41(100)		
<b>Mean BP</b>					
Systolic (Mean±SD)	147±18	115± 5.8			
Diastolic (Mean±SD)	92±15	76±5.3			

**Table 2: Subjects and Gender Distributions of C677T Genotypes**

	<b>CC(%)</b>	<b>CT(%)</b>	<b>TT(%)</b>	<b>Total (%)</b>	<b><math>\chi^2</math></b>	<b><i>p</i> value</b>
Hypertensive	37(74.0)	6(12.0)	7(14.0)	50(100)	5.759	0.049
Normotensive	21(84.0)	4(16)	0(0.0)	25(100)		
Total	58	10	7	75		
<b>Sex</b>						
Male	30(83.4)	3(8.3)	3(8.3)	36(100)	2.727	0.256
Female	28(71.8)	7(17.9)	4(10.3)	39(100)		
Total	58	10	7	75		

**Table 3 : The Logistic Regression for Independent Predictors of Hypertension among Subjects**

<b>Variables</b>	<b>B</b>	<b>OR</b>	<b>95% CI</b>	<b>p value</b>
Presence of Mutant	1.939	3.955	1.101- 10.034	0.033
MTHFR genotypes				
Age	0.572	1.771	1.036 – 3.029	0.037
sex	5.011	15.052	0.196 - 115.028	0.139
Alcohol use	-21.747	0.000	0.000 - 0.000	0.999
Smoking	-21.809	0.000	0.000 - 0.000	0.999