

Review Article

THE USE OF EXOGENOUS ENZYMES FROM PLANTS IN FEED FOR VARIOUS CULTIVATION ORGANISMS; LITERATURE REVIEW

ABSTRACT

The presence of enzymes in the feed will increase the digestibility of fish to feed ingredients. Fish need enzymes to help speed up the process of digestion and hydrolysis. Enzymes function as catalysts that can increase the speed of chemical reactions. Therefore, exogenous enzymes are needed that can accelerate the protein hydrolysis process to accelerate fish growth. Several studies have been conducted regarding enzymes derived from plants that aim to accelerate fish growth. It is hoped that this journal can provide information related to the use of several enzymes that have been used for various cultured organisms.

Keywords: Cultivation, Enzyme, Growth, Hydrolysis, Protein

1. INTRODUCTION

Low daily growth in fish can be caused by several factors, including genetic factors, poor water quality, and the amount of feed that is not following the needs of the fish. Therefore, fish feed needs to be added with exogenous enzymes to provide good growth results.

The presence of enzymes in the feed will increase the digestibility of fish to feed ingredients. Fish need enzymes to help speed up the process of digestion and hydrolysis [1]. Enzymes can be produced from several sources, such as from animals and plants, in the form of liquid or solid extraction results. In this case, the enzyme extract in solid form has several advantages, such as being easier to store and relatively longer storage period than enzymes in liquid form.

The purpose of the review is to provide information about the use of several enzymes that have been used for various cultured organisms.

2. ENZYME

2.1 Definition, Characteristics, and Types

Enzymes are a group of proteins that regulate and carry out chemical changes in biological systems. Enzymes produced by organs in animals and plants are catalyzed by various reactions, such as hydrolysis, oxidation, reduction, isomerization, addition, radical transfer, carbon-chain termination [2]. In general, enzymes provide speed, specification, and control overreactions in the body. Enzymes function as catalysts, namely compounds that increase the speed of chemical reactions [3].

According to Sutrisno [4], enzymes have five properties, namely:

1. Enzymes are biodegradable, which means they can be biodegraded. Enzymes have safe and more environmentally friendly properties because they are part of natural living systems. When degraded, enzymes produce various amino acids that can be reabsorbed by nature.
2. Enzymes can be used repeatedly. It means that enzymes can be used and making them more efficient. This is because enzymes do not undergo permanent changes in the catalytic process.
3. Enzymes are unique or specific, which means that enzymes only work according to their substrates so that enzymes are only suitable for one or a group of substrates whose function and structure are more or less the same.
4. Enzymes are biocatalysts. Enzymes can change the rate of a reaction without changing its structure. Changing this speed means slowing down or speeding up the reaction.
5. The enzyme does not participate in the reaction because the enzyme has a fixed structure or will not change. The enzyme structure will remain the same, either before or after the reaction. Because enzymes only change the rate of reaction.

Enzymes consist of several groups. According to Zonneveld et al., [5], the enzymes that play a role in the digestion of fish are proteases, amylase, and lipase, which catalyze the breakdown of complex nutrients (proteins, carbohydrates, and fats) into simple nutrients.

Table 1. Types and Characteristics of Enzymes

No	Enzyme Type	Enzyme Characteristics
1.	Amylase	Splitting starch amylase molecules at -1,4-glycoside and -1,6-glycoside bonds [6].
2.	Lipase	It breaks the ester bonds in fats to become fatty acids and glycerol [7].
3.	Protease	Hydrolyze proteins into simpler fragments and hydrolyze polypeptides into amino acids.

2.2 Enzyme Working Mechanism in Digestion of Fish

According to Campbell [8], the mechanism of action of enzymes is that the substrate first binds to the enzyme and then is induced until it fits (induced fit), then the catalysis process, and finally, the product is released. The reaction product is released from the active site, and the enzyme remains in its original form. The enzyme can then leave the active site and be reused with a new substrate. As a biocatalyst, the work of enzymes can be influenced by several factors, including temperature, pH, substrate concentration, inhibitors, and activators. Some types of enzymes can only work at their optimum temperature and pH. The optimum temperature and pH for each type of enzyme are different. Enzymes cannot work too low or too high temperatures [8].




Figure 1. Enzyme mechanism of action
Source: Campbell 2002

3. APPLICATIONS OF VARIOUS ENZYMES IN CULTIVATION ORGANISMS

One of the things that need to be considered in cultivation activities is feeding. Feed is one of the most essential aquaculture inputs because the production cost from feed can reach 60% [9]. One of the critical elements in feed is protein because the protein content in feed determines fish growth. Protein is the most expensive feed nutrient compared to other feed nutrients, but the amount of protein available in the feed must be sufficient and not excessive [10]. Therefore, exogenous enzymes are needed to accelerate the protein hydrolysis process to accelerate fish growth.

Several studies have been conducted regarding enzymes derived from plants that aim to accelerate fish growth. Faizal [11] uses papain enzymes and bromelain enzymes to feed

Tilapia. Papain is a protease enzyme, one of which is found in papaya. The enzyme is used to break or decompose peptide bonds in proteins to break down into simpler peptide bonds because papain can catalyze hydrolysis reactions of a substrate.

Bromelain is an enzyme that can help dissolve mucus formation and speed up fat elimination through the kidneys. Bromelain also has citric and malic acids, which are essential and needed to improve removing fat and manganese and being an important component of certain enzymes needed in protein and carbohydrate metabolism [12].

The content of enzymes in some easily available fruits such as papaya and pineapple have proteolytic properties or can simplify protein into amino acids that fish can digest. This encourages the utilization of the enzymes contained in the fruit is vital. Putri [13] researched the effectiveness of turmeric flour to optimize the work of digestive enzymes, which results in increasing the activity of digestive enzymes and the growth performance of carp.

The research conducted by Insana [14] regarding the substitution of Curcuma flour (*Curcuma xanthorrhiza* sp) in feed on the growth and survival of tilapia fish (*Oreochromis niloticus*) gave significantly different results. Curcuma has various benefits for the fish body, especially for health and growth. In addition to containing antibiotics. Curcuma also contains essential oils and curcumin. Curcumin functions to increase appetite and plays a role in increasing the work of the digestive organs stimulates bile walls to secrete fluids and stimulates the release of pancreatic juice, which contains amylase, lipase, and protease enzymes to improve digestion of carbohydrate, fat, and protein feed ingredients.

The following is a table regarding types of enzymes, sources of enzymes, and test organisms that have been carried out in several studies that have been carried out previously.

Table 2. Types, Sources of Enzymes, Test Organisms

No.	Enzyme Source	Enzyme Type	Test Organism
1.	Papaya	Papain	Tilapia [11] Nilem Fish [15] Catfish [16] Tawes Fish [17] Cantang Grouper [18,19] Fresh Lobster [20]]
2.	Pineapple	Bromelain	Tilapia [11] Nilem Fish [15] Catfish [16]
3.	Turmeric Flour		Gourami [21] Goldfish [13] White Snapper [22,23]
4.	Curcuma Flour		Tilapia [14] Milkfish [24] Goldfish Koi [25]

4. CONCLUSION

Several studies have been conducted regarding the use of enzymes derived from plants that aim to accelerate fish growth. Several enzymes that have been known to give significant results on growth are papain enzymes and bromelain enzymes which can provide good growth results. In addition, turmeric flour and curcuma flour can also accelerate the protein

hydrolysis process because they make the performance of digestive enzymes more optimal in some freshwater and seawater fish commodities.

REFERENCES

1. Sari, W. A. P., Subandiyono, dan Sri Hastuti. 2013. Pemberian Enzim Papain untuk Meningkatkan Pemanfaatan Protein Pakan dan Pertumbuhan Benih Ikan Nila Larasati (*Oreochromis niloticus* Var.). *Journal of Aquaculture Management and Technology*. 2 (1): 1-12.
2. Supriyatna A, Amalia D, Jauhari A. A., Holydaziah D. 2015. Aktivitas Enzim Amilase, Lipase, Dan Protease Dari Larva *Hermetia illucens* yang diberi pakan jerami padi. *Jurnal Istek IX* (2): 18-32
3. Marks, D.B.A.D., Marks, C.M., Smith. 2000. *Biokimia Kedokteran Dasar: Sebuah Pendekatan Klinis*. Penerbit Buku Kedokteran EGC. Jakarta
4. Sutrisno, A. L. (2017). Studi aktivitas antidiabetes minuman herbal kayu manis (*cinnamomum burmanni*), jahe (*zingiber officinale*), dan jeruk nipis (*citrus aurantifolia*) secara in vitro (Doctoral dissertation, Universitas Pelita Harapan).
5. Zonneveld N, Huisman EA, Boon JH. 1991. *Prinsip-prinsip budidaya ikan*. Jakarta: Gramedia Pustaka.
6. Richana, N. (2000). Prospek dan Produksi Enzim Alfa-amilase dari Mikroorganisme. *Tinjauan Ilmiah Riset Biologi dan Bioteknologi Pertanian*, 3(2), 78.
7. Poedjiadi, A., & Supriyanti, F. T. (2006). *Dasar-dasar Biokimia Edisi Revisi*. UI-Press. Jakarta.
8. Campbell, N.A., Reece, J.B., & Mitchell, L.G. (2002). *Biologi*. Jilid 1. Edisi Kelima. Alih Bahasa: Wasmen. Jakarta: Penerbit Erlangga.
9. Afrianto, E. dan E. Liviawaty. 2005. *Pakan Ikan*. Kanisius. Yogyakarta.
10. Halver, J.E, Hardy,R.W,. 2002. *Fish Nutrition*. Third Edition. Academy Press Inc, New York
11. Faizal D, Rostika R, Yustiati A, Andriani Y, Zidni I. 2017. Pengaruh Penambahan Kombinasi Ekstrak Enzim Kasar Papain Dan Bromelin Pada Pakan Buatan Terhadap Pertumbuhan Benih Ikan Nila (*Oreochromis niloticus*). *Jurnal Perikanan dan Kelautan*. VIII (1): (56-63)
12. Winastia, B. 2011. Analisa Asam Amino pada Enzim Bromelin dalam Buah Nanas (*Ananas comusus*) Menggunakan Spektrofotometer. [Tugas Akhir]. Prodi Diploma III Teknik Kimia Universitas Diponegoro, Semarang.
13. Putri, I. W., Setiawati, M., Jusadi, D. Enzim pencernaan dan kinerja pertumbuhan ikan mas, *Cyprinus carpio* (*Linnaeus, 1758*) yang diberi pakan dengan penambahan tepung kunyit *Curcuma longa* Linn.
14. Insana, N., Wahyu, F. 2015. Subtitusi Tepung Temulawak (*Curcuma xanthorhiza* Sp) Pada Pakan Dengan Dosis Berbeda Terhadap Pertumbuhan Dan Sintasan

Benih Ikan Nila (*Oreochromis niloticus*). OCTOPUS : Jurnal Ilmu Perikanan. 4(2): 381-391

15. Rostika, R., Sunarto, H.N. Sugiyanto, L.P. Dewanti,. 2018. The Effectiveness of Crude Papain- Enzyme Supplementation for Tilapia's (*Oreochromis niloticus*) Growth at the Cultured at Floating Net Cage of Cirata Reservoir. Earth and Environment Science Proceeding 139.
16. Rostika, R., T. Herawati , Dewanti, L.P., Suhendar, A.H.K. 2018. The Effect of Additional Papain Crude Enzyme In Feed On The Growth Rate Of Barb Juvenile (*Osteochilus hasselti*) Fry In Floating Net Cage Aquaculture Cirata Reservoir. Global Scientific Journal 6(5):61-66
17. Rostika R, A. Nurhayati, I.D. Buwono, A. Rizal, L.P. Dewanti, T. Maulana, 2018. Papain and bromelain crude enzyme extract in commercial feed, effectiveness toward pisciculture production of striped catfish (*Pangasianodon hypophthalmus*) in aquaculture facility. AACL Bioflux 11(5)
18. Rostika, R. A. Yustiati, A. Rizal, M. Rudyansyah, L.P. Dewanti. 2019. Intestinal Histology of the grouper *Epinephelus lanceolatus-fuscoguttatus* Administered with *Simplicia* Papaya. Biotropia 26(3): 208 – 213
19. Rostika, R. , L.P. Dewanti, M. Rudyansyah., 2018. Additional Of Papain Crude Enzyme Extract, In Feed On The Growth Rate Of Cantang Grouper (*Epinephelus Fuscogutattus Lanceolatus*) Fry In Floating Net Cage At Pangandaran Coastal. International Journal of Agriculture and Environmental Research 4(2)
20. Rachmawati, D., A. A. Prihanto, R. H. Setyobudi, O. Anne. 2018. Effect of Papain Enzyme in Feed on Digestibility of Feed, Growth Performance, and Survival Rate in Post Larvae of Freshwater Lobster [*Cherax quadricarinatus* (Von Martens, 1868)]. Proceedings of the Pakistans Academy of Sciences: B. Life and Environmental Sciences 55 (3) 31 – 39
21. Arifin, P.P. 2015. Evaluasi Pemberian Ekstrak Kunyit Curcuma Longa Linn. pada Pakan Terhadap Enzim Pencernaan dan Kinerja Pertumbuhan Ikan Gurame *Osphronemus Gouramy*. [Skripsi]. Bogor: Institut Pertanian Bogor.
22. Santika, L., Diniarti, N., Astriana. 2021. Pengaruh Penambahan Ekstrak Kunyit Pada Pakan Buatan Terhadap Pertumbuhan Dan Efisiensi Pemanfaatan Pakan Ikan Kakap Putih (*Lates calcarifer*). Jurnal Kelautan Indonesian Journal of Marine Science and Technology 14(1):48-57
23. Anriyono, Henky, I., & Wiwin, K.A.P. (2018). Pertumbuhan Benih Ikan Kakap Putih (*Lates calcarifer*) dengan Pemberian Dosis Pakan yang Berbeda. Jurnal Akuakultur Indonesia, 2(3), 2-19
24. Prabowo, A. S., Madusari, B. D., Mardiana, T. Y. 2017. Pengaruh Penambahan Temulawak (*Curcuma xanthorrhiza*) Pada Pakan Buatan Terhadap Pertumbuhan Ikan Bandeng (*Chanos chanos*). PENA Akuatika 15(1): 40 – 48
25. Monoarfa, VD, Mansyur, K., Tis'in, M., Ndobe, S. 2020. Penambahan Tepung Temulawak (*Curcuma xanthorrhiza roxb*) Padapakan Terhadap Pertumbuhan Ikan

