

1 **Reviewer Details:**

2
3 **Name: Gemechu Yigezu Ofgeha**

4 **Department, University & Country Wollega University, Ethiopia**

5
6 **Soil erosion monitoring indicators: An approach towards natural resource**
7 **management in Kuresoi South, Kenya**

8
9 Chepkirui Gloria Baliach¹, Andrew Kiplagat², Anne Sitienei*¹

10 *1 Department of Agroforestry, Environmental Studies and Integrated Natural Resources,*
11 *University of Kabianga*

12 *2 Department of Environmental Monitoring, Planning and Management, University of Eldoret*

13 **Corresponding Author*

14 **Abstract**

15 Soil erosion is still one of the most important land problems and it is linked to land use
16 and land cover changes. These have negative effects on land resource which ultimately
17 affects agricultural productivity and water quality. Local monitoring systems constitute
18 an almost compulsory component of any program or project dealing with sustainable
19 management of natural resources. The purpose of this study was to identify soil erosion
20 monitoring indicators in Kuresoi South, Kenya. The study was comprised of a total
21 representative sample population of 68 respondents from Kuresoi south catchment
22 which was achieved using Nassiuma coefficient of variation formulae. Our findings
23 reveal a positive significant relationship between soil erosion monitoring indicators and
24 natural resource management. Taken together, soil erosion monitoring indicators can
25 be used in detecting change over time in soil resource.

26 **Keywords: Soil erosion, Environment, Food Security,**

27 **1. Introduction**

28 Soil erosion is a major problem confronting land and water resources. The rate of
29 erosion is primarily determined by the erosive events (e.g., short duration and high-
30 intensity rainfall events), soil type, and characteristics of the terrain (Wei et al., 2007).
31 The impacts of accelerated soil erosion processes can be severe, not only through land
32 degradation and fertility loss but through a conspicuous number of off-site effects such
33 as sedimentation, siltation, and eutrophication of waterways or enhanced flooding
34 (Borrelli et al., 2017).


35 Soil erosion rates are exacerbated for the arid and semi-arid regions due to barren
36 mountains with scattered vegetation that provide direct exposure to heavy rainfall (Vijith
37 et al., 2018). In Kenya, Soil erosion is one of the most important land problems and it is

38 linked to land use and land cover changes. For instance, this problem has persisted in
39 Kuresoi South where its negative effects on land resource, soil productivity, available
40 agricultural land, and water resources due to sedimentation has been dominant.

41 Soil erosion monitoring indicators are strategies that are used in detecting change over
42 time in soil resource. They help during natural resource decision making during
43 environmental planning and management. Soil erosion is manifested in crop yield
44 reduction. (Munodawafa, 2012), reduced quality of the water (Zhai, 2014), building up of
45 rills and gullies (Blanco-Canqui and Lal 2008), exposure of roots and rocks (Stoffel et
46 al., 2013) as well as Sediment deposition (Shen&Julien, 1993; Okoba, B. O., &Sterk, G.,
47 2006).

48 Low vegetation covers and poorly developed soils intensify wind erosion (Jones et al.
49 2013; Blanco-Canqui and Lal 2008).

50 To date, various studies have been conducted to determine the strategies employed in
51 monitoring soil erosion (Shen et al., 2009; Wang et al., 2009; Wu & Chen, 2012; Xu, Xu,
52 Wu, & Tang, 2012). Nonetheless, less focus has been directed to Kuresoi South ward
53 yet it is an agriculturally productive area which is experiencing high population growth.
54 In fact, farmers in this area continue to experience soil erosion despite effort to
55 conserve soil. This threatens agronomic productivity, the environment, food security,
56 quality, and the well-being of many small-scale farmers. Therefore, the present study
57 seeks to answer the question on whether there any soil erosion Monitoring Strategies in
58 Kuresoi. Providing an answer to this question will immensely help the land use
59 planners, environmental planners, and policy makers to identify and execute site
60 specific best management practices to bring soil erosion rates within the permissible
61 limits at the local environment.



62

63

64 **Figure 1: An illustration on the association between soil erosion monitoring**
65 **indicators and natural resource management**

66

67 2. Materials and methods

68 **2.1 Study area**

69 This study focuses on Kuresoi ward in Nakuru County, within a latitude of -0.3015° S
70 and a longitude of 35.5307° E. Its elevation is 2551 meters feet. It is located next to the
71 South West Mau Forest and is experiencing high population growth and people engage
72 in wide range of agricultural activities such as farming, poultry and herd keeping for their
73 livelihood. The area under study is under significant human pressure through
74 encroachment to the remaining parts of the forest, charcoal burning, grazing, illegal tree
75 logging, smallholder agriculture and subsistence farming. The area is therefore prone to
76 soil erosion due to the many human activities taking place. It was therefore important to
77 identify some soil erosion monitoring strategies to manage the problem of soil erosion.

78 **2.2 Research Design**

79 This study used descriptive research design to describe soil erosion monitoring
80 strategies as indicators of soil erosion and natural resource degradation. This was
81 deemed since it helped to provide answers to the raised questions (Bryman &
82 Cramer.,1997) and how associated with soil erosion monitoring strategies.

83 **2.3 Target Population**

84 Kothari (2004) defines target population as the total number of items that the study
85 intends to investigate them. The target population in this study was the small holder
86 farmers of Kuresoi South constituency, Kuresoi ward with specific focus to three
87 villages; Mwaragania, Kibugat and South B within the affected regions. The total
88 population of this area is 6,649 (2019 Census) and is distributed in the table below.

89 **2.4 Sample and Sampling Procedures**

90 Sampling is the selection of subset of units, people, or items, from the target population.
91 This is for the purpose of collecting information which is used to draw deductions about
92 the entire population (Kothari, 2014). A sample is the subset of units that are selected
93 and they are used to represent the entire population (Mugenda & Mugenda, 2003).
94 According to Abraham and Rusell (2008), a sample size should be greater than 30 for
95 inferential statistics to be conducted. In this study, the sample was 68 households and
96 was obtained using Nassiuma Coefficient of variation formula (Nassiuma, 2000).

$$97 \quad S = \frac{N(Cv^2)}{Cv^2 + (N-1) e^2} \quad \text{Where:}$$

98 S = Sample size

100 N = Total size of the population (6,649)

101 Cv = the Coefficient of Variation (25%)

102 e = marginal error (3%)

103

104 **2.5 Data Collection Instruments**

105 Primary data collection was used in the collection of data where open and closed ended
106 questionnaires adopted as well as the use of camera for capturing picture. Open ended
107 questions gave deeper information about the soil erosion monitoring strategies while
108 closed ended provided quantitative analysis for the study.

109 **2.6 Validity of the Research Instrument**

110 This is the adequate reporting of the objectives under study and the measure of
111 accuracy (Cohen et al, 2000). Instrument's validity is significant for logical premises and
112 accuracy (Oso and Onen, 2008). The instruments were interrogated by the supervisors
113 in the university together with the peers and the way forward was decided. The pre-
114 testing of the instruments enabled for the evaluation of the content's validity.

115 2.7 Reliability of the Research Instrument

116 A pilot study was conducted in order to ascertain the reliability of the research
117 instruments, detect any ambiguities, identify the questions that are constructed poorly
118 and cannot be understood together with those questions that are irrelevant. Mugenda
119 and Mugenda (2003) recommended 10% of the sample size is appropriate for pilot
120 study. A pilot of 7 respondents from the target population was selected randomly to test
121 the questionnaires. Cronbach alpha with a set lower limit of Cronbach alpha 0.6
122 acceptability was used to analyze the results of the pilot test. The study found an overall
123 Cronbach alpha results of 0.762 which is more than the recommended threshold of 0.6.

124 2.8 Data analysis


125 Descriptive statistics were used to analyze the data in this study. This describes and
126 explains what the data shows about soil erosion monitoring strategies. After data
127 collection, the researcher edited, coded, and presented the results in the form of
128 frequency tables, graphs and pie-charts for easier understanding and interpretation.
129 Descriptive statistics such as mean was used to summarize the data. Regression model
130 was also used to establish the relationship between soil erosion indicators and soil
131 management.

132

133 3. Results and discussion

134 3.1 demographic characteristics of respondents

135 The age and education level of the respondents were established and the results
136 provided in figure 2 below



137

138 **Figure 2: Age and education level of the respondents**

139

140 The largest number came from age bracket of 20 – 40 years with 24(41%), followed by
141 40 – 60 years with 22(37%) then 60 and above with 9(15%) and finally 18 and below
142 with 4(7%). It is evident that most of the respondents are youths in the society who are
143 active and engage in economic activities like farming and animal rearing which are the

144 major factors causing soil erosion. The highest percentage of respondents completed
 145 primary and secondary education with 57% and 22% respectively. This indicates that
 146 most respondents achieved basic education while a few advanced their studies.
 147


148 3.2 Landscape characteristics

149 How the area was when the respondents first came was also assessed. It was provided
 150 in table1 below;
 151

152 **Table 1: The state of the area initially**

	Frequency	Percent	Valid Percent	Cumulative Percent
Forested	39	66.1	66.1	66.1
Grassland	5	8.5	8.5	74.6
Valid Agricultural land	9	15.3	15.3	89.8
Bare land	6	10.2	10.2	100.0
Total	59	100.0	100.0	

153
 154 The findings indicated that 39(66%) of the respondents indicated that the area was
 155 forested when they first came. 9(15%) indicated that it was an agricultural land while
 156 6(10%) indicated that it was a bare land and 5(9%) indicated that it was a grassland.



157
 158 **Figure 3: The state of the area initially**
 159

160
 161 Most respondents (66%) indicated that the area was forested when they first came. This
 162 shows that soil erosion was not a serious environmental problem at that time. However,
 163 15% responded that it was an agricultural land, to show that as time went by, there was
 164 massive migration of people to this land and the residents began to clear the forests in
 165 their land to create space for farming.
 166



167
 168 **Figure 4: Soil erosion monitoring indicators (a) Rills, (b) Gullies, (c) Tree root**
 169 **exposure, (d) Rock exposure, (e) Pedestal, and (f) Decolorized River water.**

170
 171 Decline in crop yield averagely indicated the presence of soil erosion (mean of 2.8136).
 172 Small channels greatly indicated the presence of rill erosion (mean of 3.4068 whereas
 173 deep channels indicated the occurrence of gully erosion (mean of 3.6102). The river
 174 water had been significantly decolorized and roots of plants exposed due to soil erosion
 175 (mean of 3.2203 and 4.2034 respectively). At the same significant rate, rocks located in
 176 the rivers had been exposed and sediments found along the drainages and river banks
 177 to indicate the presence of soil erosion (mean of 3.0169 and 4.2034 respectively).
 178 Change in soil color, tree mounds and pedestals indicated the presence of soil erosion
 179 at a low rate since they are observed after a long period of time (mean of 2.6271,
 180 2.6102 and 2.0169 respectively).

181
 182 Soil erosion monitoring indicators were therefore significant in assessing the
 183 effectiveness of natural resource management by a mean of 3.9492. Studies by
 184 Ypsilantis (2011) support these studies. He proposes that soil erosion monitoring
 185 indicators are important in providing qualitative assessment of erosion so that the sites
 186 that indicate potential erosion problems be red flagged and mitigating measures can be
 187 implemented to correct the problem. Rills and gullies were presented at a highly
 188 significant rate to show that rill erosion and gully erosion had taken place in most places
 189 and the necessary mitigation measures had to be put in place.

190
 191 **Table 2: Response on Soil erosion monitoring indicators**

Soil erosion monitoring indicator	1(SD)	2(D)	3(N)	4(A)	5(SA)	Mean	STD
-----------------------------------	-------	------	------	------	-------	------	-----

There is decline in crop yield in my farm due to soil erosion	0(0.00%)	12(20.3%)	11(18.6%)	12(20.3%)	24(40.7%)	2.8136	1.18139
There are small channels in my farm that indicate rill erosion	0(0.00%)	0(0.00%)	35(59.3%)	24(40.7%)	0(0.00%)	3.4068	.40545
There are deep channels in the fields and pathways which indicate the presence of gully erosion	0(0.00%)	0(0.00%)	35(59.3%)	12(20.3%)	12(20.3%)	3.6102	.80979
The river water have been decolorized due to soil erosion and thus not fit for human consumption	0(0.00%)	23(39.0%)	0(0.00%)	36(61.0%)	0(0.00%)	3.2203	.98380
The roots of plants and trees have been exposed due to soil erosion	0(0.00%)	0(0.00%)	0(0.00%)	47(79.7%)	12(20.3%)	4.2034	.40598
Rocks located in the fields and rivers have been exposed due to prolonged soil erosion	0(0.00%)	23(39.0%)	24(40.7%)	0(0.00%)	12(20.3%)	3.0169	1.10628
There are sediments on the drainages/river banks which shows that soil erosion has taken place	0(0.00%)	0(0.00%)	0(0.00%)	47(79.7%)	12(20.3%)	4.2034	.40598
The color of the soil in the farm has changed due to soil erosion	11(18.6%)	12(20.3%)	24(40.7%)	12(20.3%)	0(0.00%)	2.6271	1.01537
The soil under the trees in the field is at a higher level than the soil in the surrounding area which indicates prolonged soil erosion	0(0.00%)	0(0.00%)	36(61.0%)	23(39.0%)	0(0.00%)	2.6102	.49190
There is a column of soil that stands out from the general eroded surface on the fields as a result of prolonged soil erosion	23(39.0%)	12(20.3%)	24(40.7%)	0(0.00%)	0(0.00%)	2.0169	.90003
If soil erosion monitoring strategies are identified, it will help to manage soil erosion	0(0.00%)	0(0.00%)	19(32.2%)	24(40.7%)	16(27.1%)	3.9492	.77512

192 SD=Strongly disagree, D=Disagree, N=Neutral, A=Agree, SA= Strongly agree

193

194

195 **Table 3: ANOVA soil erosion monitoring indicators**

	Sum of Squares	Df	Mean Square	F	Sig.
Between Groups	5.561	2	2.781	27.684	.000
Within Groups	5.625	56	.100		
Total	11.186	58			

196 The results of the findings above revealed that at the level of significance 0.05 soil
197 erosion monitoring indicators identified by the community are significant in natural
198 resource management (F = 27.684, P<0.05).

199 **Discussion**

200 The findings revealed that most respondents experienced decline in crop yield as result
201 of soil erosion. The soil erosion has caused rills, gullies, tree root exposure,
202 decolorisation of the river water, rock exposure and sedimentation as physical indicators
203 of soil erosion in their farms and fields. Change in soil color, tree mounds and pedestals
204 were experienced by few respondents. Therefore, in order to obtain high success rates
205 in natural resource management, there is need to implement management techniques
206 that will curb rill, gully, and sheet erosion since these types of erosion are responsible

207 for decline in crop yield, decolorisation of the river water, sedimentation, tree/plant root
208 exposure and rock exposure. Soil erosion monitoring indicators significantly affect
209 natural resource management. It was also reflected in the regression model where soil
210 erosion monitoring indicators was the second leading variable on predicting natural
211 resource management

212 **Conclusion**

213 The study established that soil erosion monitoring indicators (rills, gullies, Tree root
214 exposure, Rock exposure, Pedestal, and Decolorized River water) are important in
215 natural resource management. These indicators are used in detecting change over time
216 in soil resource. They help during natural resource decision making during
217 environmental planning and management. **Therefore, training needs and capacity
218 building on the adoption of soil erosion monitoring indicators at a local scale**

219 **References**

220
221 Wei, W., Chen, L., Fu, B., Huang, Z., Wu, D., & Gui, L. (2007). The effect of land uses
222 and rainfall regimes on runoff and soil erosion in the semi-arid loess hilly area,
223 China. *Journal of hydrology*, 335(3-4), 247-258.

224
225 Borrelli, P., Robinson, D. A., Fleischer, L. R., Lugato, E., Ballabio, C., Alewell, C., ... &
226 Panagos, P. (2017). An assessment of the global impact of 21st century land use
227 change on soil erosion. *Nature communications*, 8(1), 1-13.

228
229 Vijith, H., Hurmain, A., & Dodge-Wan, D. (2018). Impacts of land use changes and land
230 cover alteration on soil erosion rates and vulnerability of tropical mountain ranges in
231 Borneo. *Remote Sensing Applications: Society and Environment*, 12, 57-69.

232
233 Munodawafa, A. (2012). The effect of rainfall characteristics and tillage on sheet erosion
234 and maize grain yield in semiarid conditions and granitic sandy soils of Zimbabwe.
235 *Applied and environmental soil science*, 2012.

236 Zhai, X., Xia, J., & Zhang, Y. (2014). Water quality variation in the highly disturbed Huai
237 River Basin, China from 1994 to 2005 by multi-statistical analyses. *Science of the Total
238 Environment*, 496, 594-606.

239 Blanco-Canqui, H., & Lal, R. (2008). Principles of soil conservation and management.
240 Springer Science & Business Media.

241 Shen, H. W., & Julien, P. Y. (1993). Erosion and sediment transport. Handbook of
242 Hydrology. McGraw-Hill, Inc, New York, NY, 12-1.

243 Okoba, B. O., & Sterk, G. (2006). Quantification of visual soil erosion indicators in
244 Gikuuri catchment in the central highlands of Kenya. *Geoderma*, 134(1-2), 34-47.

245 Stoffel, M., Corona, C., Ballesteros-Cánovas, J. A., & Bodoque, J. M. (2013). Dating
246 and quantification of erosion processes based on exposed roots. *Earth-Science
247 Reviews*, 123, 18-34.

248 Shen, Z. Y., Gong, Y. W., Li, Y. H., Hong, Q., Xu, L., & Liu, R. M. (2009). A comparison
249 of WEPP and SWAT for modeling soil erosion of the Zhangjiachong Watershed in the
250 Three Gorges Reservoir Area. *Agricultural Water Management*, 96(10), 1435-1442.

251 Wang, L. Qu, J. (2009). Satellite remote sensing applications for surface soil moisture
252 monitoring: A review. *Front Earth Sci Chin.* 3(2)237-247.

253 Xu, H. S., Xu, Z. X., Wu, W., & Tang, F. F. (2012). Assessment and spatiotemporal
254 variation analysis of water quality in the Zhangweinan River Basin, China. *Procedia*
255 *Environmental Sciences*, 13, 1641-1652.

256 Bryman, A., & Cramer, D. (1997). *Quantitative data analysis with SPSS for Windows: A*
257 *guide for social scientists*. Routledge.

258 Kothari, C. R. (2004). *Research methodology: Methods and techniques*. New Age
259 International.

260 Abraham, W. T., & Russell, D. W. (2008). Statistical power analysis in psychological
261 research. *Social and Personality Psychology Compass*, 2(1), 283-301.

262 Mugenda, O. M., & Mugenda, A. G. (2003). *Research methods. Quantitative and*
263 *qualitative approaches*, 46-48.

264 Nassiuma, D. K. (2000). *Survey sampling. Theory and methods*, 10.

265 Oso, W. Y., & Onen, D. (2008). *Writing research proposal and report: A handbook for*
266 *Public Personnel Management*.

267 Cohen, M. E., & Marino, R. J. (2000). The tools of disability outcomes research
268 functional status measures. *Archives of physical medicine and rehabilitation*, 81, S21-
269 S29.

270 Kombo, D. K. (2004). *Girl parents in secondary schools in Kenya: An examination of pre*
271 *and post pregnancy performance*. Unpublished thesis, Egerton University). Nakuru.

272 Ypsilantis, W. G., Biggam, P., Bryant, L., Davis, R., Herrick, J. E., Horn, E., ... & Smith,
273 C. (2011). *Upland soil erosion monitoring and assessment: An overview*. US
274 Department of the Interior, Bureau of Land Management, National Operations Center.

275