

Original Research Article

UPTAKE AND SIDE EFFECTS PROFILE OF COVID-19 VACCINE AMONGST MEDICAL DOCTORS AT A TERTIARY HOSPITAL IN SOUTH-SOUTH NIGERIA

ABSTRACT

Background: The war against COVID-19 pandemic is currently being fought using the COVID-19 vaccine. To break the chain of transmission of the infection and establish herd immunity, the COVID-19 vaccine is that veritable tool to be used but its uptake is hindered by the fear of its side effects. As with all medications, COVID-19 vaccine has side effects that requires discovery as the disease and its vaccine are novel.

Aim: To determine the uptake of, the incidence, pattern of side effects to the COVID-19 vaccine amongst medical doctors at the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa state, Nigeria as well as the association between COVID-19 vaccine side effects and chronic medical illnesses.

Study design: This was a descriptive cross-sectional study.

Place and duration of Study: This study was carried out among medical doctors at the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa state, Nigeria, from 15th June to 30th July, 2021.

Methodology: Semi-structured questionnaires were used to obtain data from 102 participants recruited by consecutive sampling. Data obtained were the socio-demographic characteristics of the participants, the presence of co-morbidities and the presence of side effects to the COVID-19 vaccine for those vaccinated.

Result: Out of 102 medical doctors who participated in the survey, only 30 (29.4%) had received the COVID-19 vaccine and 26 of them (86.7%) reported different forms of side effects such as pain at the injection site, fever, muscle pain and joint pain. Pain at the injection site was the commonest vaccine-associated side effect (76.7%). The first day after COVID-19 vaccination was the commonest time of onset of the reported side effects. There was no statistically significant relationship between COVID-19 vaccine side effects and presence of chronic medical conditions in the participants.

Conclusion: Uptake of COVID-19 vaccine among medical doctors was low and is worrisome. Incidence of COVID-19 vaccine side effects was high; however, the symptoms were mild to moderate. In the face of this challenging pandemic, COVID-19 vaccination still remains a safe means of ensuring protection against the disease.

KEYWORDS: COVID-19 vaccine, Side effects, Medical doctors, Gender, Co-morbidities

INTRODUCTION

COVID-19 is an illness caused by a new virus called severe acute respiratory syndrome Corona virus 2 (SARS-CoV-2). It was identified first in the middle of an outbreak of respiratory illnesses in Wuhan City, Hubei province, China.¹ It was reported to WHO on the 31st of December 2019, declared a global health emergency by WHO on the 30th of January 2020 and declared a global pandemic by WHO on the 11th of March 2020.²⁻⁴

COVID-19 has serious deleterious effect globally. During the conduct of this study; as of 30th June 2021, there were 181,344,224 confirmed cases of COVID-19 including 3,934,252 deaths reported to WHO globally.⁵ Of these, Nigeria had 167,543 confirmed cases and 2,120 deaths as reported by Nigeria centre for disease control.⁶ This put so much strain on health and global economy with increased unemployment rate; as more man power, infrastructure, equipment, drugs, consumables are needed to combat this pandemic. This effect is more in developing countries including Nigeria that has an already fragile economy and health system.

Since respiratory droplets carrying infectious virus is the principal mode of transmission, the following measures are employed to prevent the spread of COVID-19: washing of hands with soap and water or use of alcohol-based hand sanitizer; maintaining a safe distance from anyone who is coughing or sneezing; wearing of facemask; avoid touching eyes, nose and mouth; covering of nose and mouth with bent elbow or a tissue when coughing or sneezing; staying at home when feeling unwell; and seeking medical attention when having fever, cough, or difficulty in breathing.⁷

While observing these preventive measures, vaccination is key to ending this pandemic. There are about 52 candidate vaccines in human trials as at December 2020.⁸ Pfizer/ bioNTech vaccine was the first to receive emergency validation by the WHO on the 31st of December, 2020.⁹ Others approved by WHO for emergency use include: AstraZeneca/Oxford vaccine, Johnson and Johnson vaccine, Moderna, Sinopharm, and Sinovac.¹⁰ Nigeria received her first batch of COVID-19 vaccine (AstraZeneca/Oxford vaccine) on the 2nd of March 2021.¹¹ As of 29th June, 2021, Two million, Two Hundred and Ninety-nine Thousand , Nine Hundred and Eighty-seven (2,299,987) people have received their first dose of COVID -19 vaccine while 1,209,591 people have received their second dose in Nigeria.¹² Bayelsa state flagged off COVID-19 vaccination on 12th of March 2021.¹³

Like other vaccines, COVID-19 vaccines are associated with some side effects which are mostly mild to moderate in severity and short lasting.^{14,15} This include pain, redness and swelling at injection site; fever; fatigue; headache; muscle pain; chills; nausea; vomiting; diarrhoea; mild flu-like symptoms; joint pain; weakness; loss of appetite; malaise; cough; and facial swelling.^{15,16} Others include transverse myelination, disorientation, hypersensitivity to light and noise, personality behaviour change. Severe allergic reactions have occurred though rarely to some COVID-19 vaccines.¹⁴⁻¹⁶ Severe allergic reaction such as anaphylaxis is a potential but rare side effect with any vaccine.¹⁶

Medical personnel, the military and other essential workers in Nigeria were among the first to be given the vaccine because of the limited supply of the vaccine to the country.¹⁶ It would be necessary to study the side effect profile of COVID-19 vaccine amongst vaccinated medical doctors because they would be in a good position to tell their experiences to other health care workers as well as their public. This study would also act as a reference tool for further research on COVID-19 vaccine uptake and side effects in Africans.

The study aimed at determining the uptake of, and the incidence and pattern of side effects to the COVID-19 vaccine amongst medical doctors at the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa state, Nigeria as well as the association between COVID-19 vaccine side effects and chronic medical illnesses.

METHODOLOGY

This was a descriptive cross-sectional study carried out among medical doctors who were members of staff at the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa state,

Nigeria, from 15th June to 30th July, 2021. The hospital is one of the two tertiary hospitals in the state. It resides within the suburbs of the state and 0.69 km from the state capital, Yenagoa. There are approximately 154 medical doctors working at the facility. **Medical doctors who were not staff and health workers who were not doctors were excluded from the study.**

Using the formula for calculating sample size in cross-sectional studies¹⁷ and based on a prevalence of COVID-19 vaccination among health workers of 27.4% from another study in Yenagoa,¹⁸ a sample size of 102 was derived for this study after adjusting for an anticipated non-response rate of 30%. A semi structured questionnaire was used to obtain data from 102 participants by consecutive sampling. **The questionnaire had four sections: section A covered the bio-data of the participants, section B explored the presence of chronic medical conditions, section C asked about the uptake of COVID-19 vaccination and Section D, the occurrence of COVID-19 vaccination side effects and day of commencement of those side effects. The data collected from the participants were age, marital status, religion, presence of chronic medical condition and the type, COVID-19 vaccination status, associated side effects of the vaccine, and the day of commencement of the side effects. All data extracted were subjected to statistical analysis using the Statistical Package for Social Sciences (SPSS) version 25.0. Categorical variables were summarized in frequencies and percentages and presented in figures and tables. Relationship between chronic medical illnesses and COVID-19 side effects, were explored using chi-squared test. The statistical significance was set at p-value of 0.05.**

RESULTS

Socio-demographic characteristics of participants

A total of one hundred and two medical doctors from the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa state, Nigeria participated in the survey. Of the 102 participants, about three-fifths were males (62.7%) while females were 37.3% (Table1). Thirty-nine (38.2%), 45 (44.1%), and 12 (11.8%) medical doctors were in the third, fourth and fifth decades of life respectively (Table 1). Majority of the participants were single (62.7%) and Christians by religion (93.1%).

Chronic medical conditions among participants

Chronic non-communicable diseases (NCD) were reported among seventeen participants (16.7%). The NCDs included hypertension (11.8%), diabetes mellitus (2.9%) and asthma (2.9%). Other conditions (2.9%) reported were chronic inflammatory synovitis and rheumatoid arthritis (Table 2).

Table 1: Socio-demographic characteristics of participants

Characteristics	Frequency (Percent) n (%)
Sex	

Male	64 (62.7)
Female	38 (37.3)
Age group	
20 - 29 years	39 (38.2)
30 - 39 years	45 (44.1)
40 - 49 years	12 (11.8)
> 50 years	6 (5.9)
Marital Status	
Single	64 (62.7)
Married	38 (37.3)
Religion	
Christian	95 (93.1)
Others	7 (6.9)

Table 2: Chronic medical conditions among respondents

Characteristics	Frequency (Percent)
	n (%)
Chronic medical condition	
No chronic medical condition	85 (83.3)
Chronic medical condition present	17 (16.7)

Type of chronic medical condition

Hypertension	12 (11.8)
Diabetes	3 (2.9)
Asthma	3 (2.9)
Sickle Cell Disease	1 (1.0)
Others*	3 (2.9)

*Others – Allergic drug reaction, rheumatoid disease, chronic inflammatory arthritis

COVID 19 vaccine uptake and reported side effects among respondents

Thirty (30) out of 102 medical doctors (29.4%) had taken the vaccine about 3 months after it was provided at the hospital for vaccination of medical personnel (Figure 1).

Of the 30 medical doctors who had taken the vaccine, 26 of them (86.7%) reported different forms of side effects (Table 3). The most common reported side effect was pain at the injection site, which was reported in 23 participants (76.7%). Other side effects included fever (60.0%), muscle and joint pain (46.7%), fatigue (40.0%) and headache (40.0%). Most of those (46.7%) who reported side effects reported it on day 1 after the injection (Table 3).




Figure 1: Vaccine uptake among Participants

Table 3: Reported side effects among respondents.

Side effects	Frequency (Percent)
	n (%)

Side effects reported	26 (86.7)
No side effect reported	4 (13.3)
Reported Site effects	
Pain in injection site	23 (76.7)
Fever	18 (60.0)
Muscle and joint pain	14 (46.7)
Fatigue	12 (40.0)
Headache	12 (40.0)
Chills	9 (30.0)
Nausea	3 (10.0)
Vomiting	1 (3.3)
Allergy	1 (3.3)
Day COVID-19 Side effects started	
Day 1	14 (58.8)
Day 2	11 (42.3)
Day 3 to 7	1 (3.8)

Association between chronic medical conditions and vaccine side effects

There was no significant difference ($p > 0.05$) in the occurrence of side effects among participants who reported chronic medical conditions and those who did not (Table 4)

Table 4: Association between chronic medical conditions and vaccine side effects

Characteristic	Total N = 30	Chronic medical condition	Chi square	P-value
----------------	-----------------	---------------------------------	------------	---------

	(%)	Present N = 3 (%)	Absent N = 27 (%)		
Side effect reported	26 (86.7)	3 (100.0)	23 (85.2)	0.37	0.54
No side effect reported	4 (13.3)	0 (0.0)	4 (14.8)		
Pain at injection site	23 (76.7)	3 (100.0)	20 (74.1)	1.01	0.31
No Pain at injection site	7 (23.3)	0 (0.0)	7 (25.9)		
Fever	18 (60.0)	3 (100.0)	15 (55.6)	2.22	0.26
No fever	12 (40.0)	0 (0.0)	12 (44.4)		
Muscle/Joint pain	14 (46.7)	1 (33.3)	13 (48.1)	0.24	0.63
No muscle/Joint pain	16 (53.3)	2 (66.7)	14 (51.9)		
Fatigue	12 (40.0)	1 (33.3)	11 (40.7)	0.06	0.80
No Fatigue	18 (60.0)	2 (66.7)	16 (59.3)		
Headache	12 (40.0)	2 (66.7)	10 (37.0)	0.99	0.55
No headache	18 (60.0)	1 (33.3)	17 (63.0)		
Chills	9 (30.0)	2 (66.7)	7 (25.9)	2.13	0.53
No Chills	21 (70.0)	1 (33.3)	20 (74.1)		
Vomiting	1 (3.3)	0 (0.0)	1 (3.7)	0.12	0.74
No Vomiting	29 (96.7)	3 (100.0)	26 (96.3)		

Nausea	3 (10.0)	0 (0.0)	3 (11.1)	0.37	0.54
No nausea	27 (90.0)	3 (100.0)	24 (88.9)		

DISCUSSION

COVID-19 vaccine is that veritable tool in the war against COVID-19 pandemic, needed to break the chain of transmission of the infection and establish herd immunity, but its uptake may be hindered by the fear of its side effects. COVID-19 vaccine side effects are yet to be well understood as the disease and its vaccine are novel.

More males than females participated in this study. This was due to the fact that more males than females presently work at the facility. Most of the participants were single and Christians as the study was carried out in a Christian-dominated locality.

From this study, most of the medical doctors had no chronic medical condition while the nature of chronic medical conditions reported by 16.7% of them was predominantly hypertension and diabetes. This is in keeping with similar studies done in Bayelsa, Lagos, China and Saudi Arabia respectively.¹⁹⁻²² Presence of co-morbidities predisposes to adverse clinical outcomes in patients with COVID-19 and may result in greater disease severity when compared with those with no co-morbidity.²¹ Thus, approved and authorized COVID-19 vaccines are especially important in people with underlying medical conditions.

COVID-19 vaccine uptake among medical doctors was low in this study; only 29.4% of the respondents took the vaccine. This low uptake rate is congruent with 27.7% recorded in Congo,²³ 37.4% in Jordan²⁴ and 39.3% in Ghana.²⁵ It is however not in consonance with 64.9% and 86.2% recorded in Saudi Arabia and China respectively.^{26,27} The low uptake rate among medical practitioners in this study is worrisome as it will likely affect the uptake of the vaccines by their patients. Maintaining confidence in vaccination depends on the interaction between patients and providers.²³ Attitude and utilization of vaccination by healthcare professionals (HCP) is a major factor that is consistently associated with patient acceptance and vaccination, adherence to vaccination schedules, and reduced hesitation/aversion.²⁸ In addition, vaccinated HCPs also have a noticeable effect on patients' decision to take a vaccine. The low uptake rate among the top echelon of the health care team may be related to the side effects of the vaccine and its overall safety. Thus, more vaccine clinical trials have to be done to provide more convincing data about the safety of this vaccine and settle all controversies about the drug once and for all.

The finding that 86.7% of those who took the vaccine in this study reported different forms of side effects is similar to studies done by Khaleduzzaman et al²⁹ and Oriji et al¹⁹, however slightly higher than the rate in studies done by Hatmal et al³⁰ and Menni et al.³¹ The most common reported side effect of pain at the injection site from this study was in consonance with a similar study among health workers in a sister facility by Oriji et al.¹⁹ It however disagrees with other studies.^{29,30} Other side effects include fever (60.0%), muscle and joint pain (46.7%), fatigue (40.0%) and headache (40.0%). Vaccines are designed to give immunity without the dangers of getting the disease however developing mild to moderate side effects is common. This is

because the immune system is instructing the body to react in certain ways increasing blood flow so more immune cells can circulate, and it raises the body temperature in order to kill the virus. Mild-to-moderate side effects, like a low-grade fever or muscle aches, are normal and not a cause for alarm: they are signs that the body's immune system is responding to the vaccine, specifically the antigen and is gearing up to fight the virus. These side effects usually last for a few days.^{31,32}

In this present study, most of the side effects (46.7%) were reported within the first day of receiving the vaccine. This is consistent with several other studies²⁹⁻³³ but inconsistent with a study done in Bangladesh²⁹ where most symptoms were reported after the third day of vaccination. Spike proteins are viral protein found on the surface of viruses and also apply to vaccines which follow the natural process of viral infection to mount an immune response. An immediate response is elucidated by the innate immune system initiating an inflammatory response resulting in fever and pain at injection site while the adaptive immunity which produces T cells and antibodies provides protection on subsequent exposure hence a delayed response.

From this study, there was no association between COVID-19 vaccine side effects and presence of chronic medical conditions. **It was likely due to the small number of vaccinated respondents.** This is consistent with the knowledge that COVID-19 vaccines may be administered to most people with underlying medical conditions.^{1,34,35}

Experiencing side effects after getting vaccinated means the vaccine is working and the immune system is responding as it should.¹⁶ After vaccination, it usually takes a few weeks for the body to build immunity against SARS-CoV-2. So, it's possible a person could be infected with SARS-CoV-2 just before or after vaccination and still get sick with COVID-19. This is because the vaccine has not yet had enough time to provide protection.

CONCLUSION

Uptake of COVID-19 vaccine among medical doctors was low and is worrisome, as it is likely to affect the uptake of the vaccines by their patients. Incidence of COVID-19 vaccine side effects was high; however, the symptoms were mild to moderate. The most common side effect was pain at injection site, observed especially within 24 hours of vaccination. In the face of this challenging pandemic, COVID-19 vaccination still remains a safe means of ensuring protection against the disease.

ETHICAL APPROVAL

This was obtained from the Research and Ethics Committee of the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa. The ethical approval number for this study is NDUTH/REC/2021/0029/0451. Informed consent for the study was obtained from respondents.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. CDC. COVID-19 (2019 Novel Coronavirus) Research Guide. CDC. Available from: <https://www.cdc.gov/library/researchguides/2019NovelCoronavirus.html>. Last reviewed September 2, 2020. Accessed August 2, 2021.
2. Gallegos A. WHO Declares Public Health emergency for novel coronavirus. Medscape Medical News. Available from: <https://www.medscape.com/viewarticle/924596>. Published January 30, 2020. Accessed August 2, 2021.
3. Ramzy A, McNeil NG. W.H.O. Declares Global Emergency as Wuhan Coronavirus Spreads. The New York Times. Available from: <https://nyti.ms/2RER70M>. Published January 30, 2020. Accessed August 2, 2021.
4. McNeil Jr DG. Coronavirus Has Become a Pandemic, W.H.O. Says. The New York Times. Available from: <https://www.nytimes.com/2020/03/11/health/coronavirus-pandemic-who.html>. Published March 11, 2020. Accessed August 4, 2021.
5. WHO Coronavirus (COVID 19) Dashboard. WHO. Available from: <https://covid19.who.int>. Accessed September 13, 2021.
6. Covid-19 Nigeria. Nigeria Centre for disease control. Available from: <https://covid19.ncdc.gov.ng>. Accessed September 13, 2021.
7. Advice for the public: Coronavirus disease (COVID-19). WHO. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public>. Accessed September 13, 2021.
8. The different types of COVID-19 vaccines. WHO. Available from: <https://www.who.int/news-room/feature-stories/detail/the-race-for-a-covid-19-vaccine-explained>. Published January 12, 2021. Accessed September 30, 2021.
9. WHO issues its first emergency use validation for a COVID-19 vaccine and emphasizes need for equitable global access. WHO. Available from: <https://www.who.int/news/item/31-12-2020-who-issues-its-first-emergency-use-validation-for-a-covid-19-vaccine-and-emphasizes-need-for-equitable-global-access>. Published December 31, 2020. Accessed September 30, 2021.
10. WHO validates sinovac COVID-19 vaccine for emergency use and issues interim policy recommendations. WHO. Available from: <https://www.who.int/news/item/01-06-2021-who-validates-sinovac-covid-19-vaccine-for-emergency-use-and-issues-interim-policy-recommendations>. Published June 1, 2021. Accessed August 21, 2021.
11. COVID-19 vaccines shipped by COVAX arrive in Nigeria. Unicef. Available from: <https://www.unicef.org/nigeria/press-releases/covid-19-vaccines-shipped-covax-arrive-nigeria#>. Published March 2, 2021. Accessed August 30 2021.
12. COVID-19 Vaccination Update. National Primary Health Care Development Agency. Available from: <https://m.facebook.com/NPHCDA/posts/4451541584870460>. Published June 29, 2021. Accessed 30 August 2021.
13. Owe O. Covid-19: Bayelsa State flags off vaccine campaign, Deputy Governor takes first jabs. Ait.live. Available from: <https://ait.live/covid-19-bayelsa-state-flags-off-vaccine-campaign-deputy-governor-takes-first-jabs/>. Accessed August 29, 2021.
14. Possible Side effects After Getting a COVID-19 Vaccine. CDC. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/expect/after.html>. Updated June 21, 2021. Accessed August 20, 2021.
15. Poole J. Covid vaccine side effects. Researchgate. Available from: https://www.researchgate.net/publication/348327013_Covid_vaccine_side_effects_Jan_8_2020. Accessed August 20, 2021.
16. Side Effects of COVID-19 Vaccines. WHO. Available from: <https://www.who.int/news-room/feature-stories/detail/side-effects-of-covid-19-vaccines>. Published March 31, 2021. Accessed 20 August 2021.

17. Oyidarín OT, Usman SA, Osoba ME, Olukorode SO, Lucero-Prisno III DE. Towards effective and efficient covid-19 vaccination in Nigeria. *J Glob Health Reports*. 2021;5:e2021023. doi:10.29392/001c.21404.
18. Araoye MO. Subject Selection. In: Research Methodology with statistics for Health and Social sciences. Ilorin, Nigeria: Nathedex Publishers; 2003:115-129.
19. Oriji PC, Allogoa DO, Wagio TJ, Obagah L, Tekenah ES, Ozori ES. Hesitancy of COVID-19 Vaccination among Health Workers (other than Doctors) in a Tertiary Health Hospital in South-South, Nigeria. *Asian J Res Infect Dis*. 2021;7(1):21-31.
20. Oriji PC, Allagoa DO, Obagah L, Oguche OI, Ohaeri OS, Tekenah ES et al. Side Effect Profile of COVID-19 Vaccine among Health Workers in a Tertiary Health Institution in South-South Nigeria. *Eur J Med Health Sci*. 2021;3(3):59-64.
21. Osibogun A, Balogun M, Abayomi A, Idris J, Kuyinu Y, Odukoya O, et al. Outcomes of COVID-19 patients with comorbidities in southwest Nigeria. *PLoS One*. 2021;16(3):e0248281. doi:10.1371/journal.pone.0248281.
22. Guan WJ, Liang WH, Zhao Y, et al. Comorbidity and its impact on 1590 patients with Covid-19 in China: a nationwide analysis. *Eur Respir J*. 2020;55(5):2000547. doi:10.111183/1393003.00547-2020.
23. Fathi M, Vakili K, Sayehmiri F, Mohamadkhani A, Hajiesmaeili M, Rezaei-Tavirani M. The prognostic value of comorbidity for the severity of COVID-19: A systematic review and meta-analysis study. *PLoS One*. 2021;16(2):e0246190. doi:10.1371/journal.pone.0246190.
24. Kabamba Nzaji M, Kabamba Ngombe L, Ngoie Mwamba G, et al. Acceptability of Vaccination Against COVID-19 Among Healthcare Workers in the Democratic Republic of the Congo. *Pragmat Obs Res*. 2020;11:103-109.
25. El-Elimat T, AbuAISamen MM, Almomani BA, Al-Sawalha NA, Alali FQ. Acceptance and attitudes toward COVID-19 vaccines: A cross-sectional study from Jordan. *PLoS One*. 2021;16(4):e0250555. doi:10.1371/journal.pone.0250555.
26. Agyekum MW, Afrifa-Anane GF, Kyei-Arthur F, Addo B. Acceptability of COVID-19 Vaccination among Health Care Workers in Ghana. *Adv. Public Health*. 2021:Article ID 9998176. <https://doi.org/10.1155/2021/9998176>.
27. Elharake JA, Galal B, Alqahtani SA, Kattan RF, Barry MA, Temsah M, et al. COVID-19 Vaccine Acceptance among Health Care Workers in the Kingdom of Saudi-Arabia. *Int J Infect Dis*. 2021;109:286-293.
28. Xu B, Gao X, Zhang X, Hu Y, Yang H, Zhou YH. Real-World Acceptance of COVID-19 Vaccines among Healthcare Workers in Perinatal Medicine in China. *Vaccines (Basel)*. 2021;9(7):704. doi.org/10.3390/vaccines9070704.
29. Asma S, Akan H, Uysal Y, et al. Factors affecting influenza vaccination uptake among health care workers: a multi- center cross-sectional study. *BMC Infect Dis*. 2016;16:192. doi:10.1186/s12879-016-1528-9.
30. Khaleduzzaman HM, Nafisa Jabin Mishu NJ. Frequency of Side Effects after First Dose of Vaccination against COVID-19 Among the People of Bangladesh. *Eur J Med Health Sci*. 2021;3(3):22-24.
31. Hatmal MM, Al-Hatamleh MA, Olaimat AN, Hatmal M, Alhaj-Qasem DM, Olaimat TM, et al. Side Effects and Perceptions Following COVID-19 Vaccination in Jordan: A Randomized, Cross-Sectional Study Implementing Machine Learning for Predicting Severity of Side Effects. *Vaccines*. 2021;9(6):556. doi.org/10.3390/vaccines9060556.
32. Menni C, Klaser K, May A, Polidori L, Capdevila J, Louca P, et al. Vaccine side-effects and SARS-CoV-2 infection after vaccination in users of the COVID

Symptom Study app in the UK: a prospective observational study. *Lancet Infect. Dis.* 2021;21(7):939-949.

33. Djanas D, Yusirwan, Martini RD, et al. Survey data of COVID-19 vaccine side effects among hospital staff in a national referral hospital in Indonesia. *Data Brief.* 2021;36:107098. doi:10.1016/j.dib.2021.107098.
34. Arvanitis M, Opsasnick L, O'Connor R, Curtis LM, Vuyyuru C, Benavente JY, et al. Factors associated with COVID-19 vaccine trust and hesitancy among adults with chronic conditions. *Prev Med Rep.* 2021(24)101484.
35. Xiao J, Cheung JK, Wu P, Ni MY, Cowling BJ, Liao Q. Temporal changes in factors associated with COVID-19 vaccine hesitancy and uptake among adults in Hong Kong: Serial cross-sectional surveys. *Lancet Reg. Health West. Pac.* 2022;23:100441.

UNDER PEER REVIEW