

# Effects of Legumes Tree Leaf Mulch Placement and N- Mineralization on Maize Productivity in a Tropical Rainforest Area

## Abstract

Nutrient depletion in the rainforest area is a land use constraint that need urgent attention. This study was carried out to investigate the effects of legume tree leaf mulch placement and N-mineralization on maize productivity in a rain forest area. Three common indigenous nitrogen fixing leguminous trees (*Albizia ferruginea*, *Albizia zygia* and *Spondias mombin*) were randomly selected around the experimental site. Fifty (50g) of freshly collected leaves from these species was bulked and weighed into 20cm x 25cm litter bags. Three litter bags were placed above-ground and three below- ground (5cm deep) at three replicates per treatment. Maize was planted at a spacing of 90cm x 30cm. Fresh samples of each mulch were applied in a ring form to the three selected plant per plot in above-ground and below-ground (5cm deep) pattern two weeks after planting. The results showed that *Albizia ferruginea* had the highest percentage nitrogen (5.49%) and *Spondias mombin* had the lowest (3.49%). The percentage calcium composition of *Albizia ferruginea* and *Albizia zygia* was the same order of magnitude (0.22%), while that of *Spondias mombin* was 0.16%. Decomposition rate of the samples was highest in *Spondias mombin* with the lowest remaining weight of 2.92g and 3.16g in above and below ground litter bags respectively at week 10. *Albizia ferruginea* leaf (61.59%) had the highest nitrogen mineralization at above ground placement and *Spondias mombin* had the lowest (52.18%). Nitrogen mineralization in the below ground placement was highest in *Spondias mombin* (67.32%) and lowest in *Albizia zygia* (40.39%). Below ground mulch placement decomposed faster than the above ground mulch placement in the selected leguminous leaf samples. *Albizia zygia* was found to have outstanding performance on height, girth and yield of maize. It is therefore recommended to poor resource farmers to allow *Albizia zygia* to thrive in and around their farms so that the leaf could be used as mulch for crop production

**Keywords:** N- Mineralization, Legume Tree Leaf Mulch Placement, Rainforest, Maize

## Introduction

Nutrient losses in tropical forests are compensated by small inputs from atmospheric deposition and through nitrogen fixation (Lu *et al.*, 2015). Fraterrigo *et al.*, (2005) reported that land-use practices can have effect on the distribution and supply of soil nutrients by directly altering soil properties and by influencing biological transformations in the rooting zone. A well-designed agroforestry system has potential to address problems of land degradation and declined in soil fertility. The aim of agroforestry in soil fertility initiative is to reverse the detrimental effects on agriculture that result in soil degradation and nutrient depletion by developing innovative soil and crop management practices that focus on soil fertility restoration and maintenance.

Provision of adequate supply of nutrients to crops has always been a leading objective of soil management. The practice of supplying deficient nutrients simply by adding fertilizers has given place to a new approach of integrated soil fertility management. This is directed at using low to moderate quantities of fertilizers in combination with biological methods of soil management. Myers *et al.*, (1994) observed that a farmer with limited cash flow cannot afford to change from a traditional system to one with high productivity through capital investment. Shifting cultivation, a sustainable traditional system of land cultivation, is increasingly becoming less efficient because of population growth and attendant land use pressures.

Current concerns about sustainability have also focused attention on the need to conserve nutrients and other resources as agriculture intensifies. The release of nutrients from decomposing organic input can be manipulated to coincide with period of nutrient demand by the crop. A relationship exists between plant growth and up-take of N. Leguminous plants can serve as an important N source for crops in many parts of the tropics where fertilizer use is not economically feasible. In the tropics, mulching has been identified as an effective strategy of changing crop growing conditions to enhance yield and improve product quality by suppressing weed development, lowering soil temperature, conserving soil moisture, reducing soil erosion, improving soil structure, and increasing soil organic matter content (Egbe *et al.*, 2012). Thus application of leaf mulch through agroforestry approach (e.g. alley cropping, tree fallow) with annual crops has the potential to improve soil fertility and productivity in the tropics.

Nutrient depletion in humid tropics is a land use constraint that need urgent attention. Knowledge of the rate at which mulch decomposes and releases nutrients could help tropical farmers improve or maintain soil fertility (Lu *et al.*, 2015). Agroforestry technology therefore has potential to address the problems. Thus, this study investigated the process of nutrient replenishment through mulch application.

## **MATERIALS AND METHODS**

### **Experimental site**

The experiment was carried out at the Agroforestry Site of the Teaching and Research Farm, Federal University Of Technology Akure, Nigeria (latitude 7° 17' N, longitude 5° 10' E) with an annual mean rainfall and temperature of 1500 mm and 26°C respectively (Adesuyi *et al.*, 2018). Relative humidity ranges between 85% and 100% during rainy season and less than 60% during harmattan period. The soils are classified as Ferruginous Tropical Soil

(Aifisol) on crystalline rock of basement complex. The soil profile shows some medium to light textured materials near the surface, followed by sandy clay subsoil and a layer of sub angular quartz gravel below. The soil of the experimental site is impoverished, this was observed through the maize grown on the surrounding ploy.

## **Experimental procedures**

### **Sample preparation and Analysis (macro nutrient composition)**

The macro-elements (Magnesium, Potassium, Calcium and Phosphorus) were determined by ashing using muffle furnace. One gram of each oven-dried sample was ashed at a temperature of 450°C for 4 hours. It was cooled in a desiccator for one hour. Each sample was now dissolved in 10ml HCl and filtered. The filtrate was diluted to 100ml. However, Calcium and Magnesium were determined by EDTA titration, Potassium was determined using flame photometer, while Phosphorus was determined by spectro-photometer. Nitrogen was also determined by Micro-Kjeldahl acid digestion procedure(Onyekwelu, 2017).

### **Decomposition and Mineralization**

Fifty (50g) of freshly collected leaves from *Albizia ferruginea*, *Albizia zygia* and *Spondias mombin* was bulked and weighed into litter bags of 20cm x 25cm made of polyvinyl material with mesh size of 2mm which was large enough to prevent compression of the enclosed litter and allow free access to most group of micro-organisms was prepared. Three litter bags were placed above-ground and three below- ground (5cm deep) at three replicates per treatment. The litter bags (three replicates each) were retrieved from the field every two weeks for a total of 12 weeks. The bags were carefully lifted up at each harvesting period to reduce losses of particulate materials. The retrieved bags were securely packed and transported to the laboratory. The experimental materials were then air dried for five days after which the weights were determined.

### **Effects of added mulch on maize growth**

Three common indigenous nitrogen fixing leguminous trees were selected around the experimental site. These were *Albizia ferruginea*, *Albizia zygia* and *Spondias mombin*. The land was manually cleared and maize (zea mays) planted early July. The land was demarcated into plot of 3m x 3m and each ploy replicated three times giving a total of twelve plots. Maize was planted at a spacing of 90cm x 30cm. Fresh samples of each mulch were applied in a ring form to three selected plant per plot in above-ground and below-ground (5cm deep) pattern two weeks after planting. The growth of the maize was measured at two weeks height interval: by metre rule and girth by micro-metre screw

gauge. At maturity, maize cobs were harvested and the grains removed from the husks and the weight per plot per treatment was determined in the laboratory by weighing.

### Statistical Analysis

The means of plant nutrient content were calculated. Analysis of Variance (ANOVA) was carried out to determine if there were significant differences among the treatment means (Onyekwelu et al., 2017). Least significant difference (LSD) was used to identify the means that were significantly different from each other. Analysis of Variance (ANOVA) was also carried out to determine significance difference in the treatment plots; time of decomposition and nitrogen mineralization and their effects on maize growth and yield. Also differences in decomposition and N-mineralization rate of mulches were determined by the same analytical tool.

## RESULTS

### Macro-nutrients composition of selected leaf sample.

The results of the laboratory analysis of leaf samples for Nitrogen, Calcium, Potassium and Phosphorus are shown in Table 1. Analysis of the micro-nutrients showed that *Albizia ferruginea* had the highest percentage nitrogen (5.49%) followed by *Albizia zygia* (5.41%) and *Spondias mombin* had the lowest percentage (3.49%). The percentage calcium composition of *Albizia ferruginea* and *Albizia zygia* was the same order of magnitude (0.20%), while that of *Spondias mombin* was 0.16%. Magnesium content was highest in *Albizia zygia* with 0.21% followed by *Albizia ferruginea* (0.17%) and lowest in *Spondias mombin* (0.14%). The highest percentage of phosphorus was obtained from *Albizia ferruginea* (0.22%), *Spondias mombin* had 0.12% and the lowest from *Albizia zygia* (0.12%). Results of the Analysis of Variance (ANOVA) followed by the Fishers' Least Significant difference (LSD) test for comparing means of macro-nutrients composition of leaf samples showed significant difference ( $p \leq 0.05$ ) in the means of nitrogen contents of the samples. There was no significant difference ( $p \geq 0.05$ ) in the means of calcium contents of both *Albizia zygia* and *Albizia ferruginea*. Ash contents of *Albizia zygia* significantly differs from that of the other two samples

**Table 1: Percentage chemical composition of leaf samples of three leguminous species**

Macro-nutrients	<i>Albizia zygia</i>	<i>Albizia ferruginea</i>	<i>Spondias mombin</i>
Nitrogen (%)	5.41 <sup>b</sup>	5.49 <sup>a</sup>	3.49 <sup>c</sup>
Calcium (%)	0.20 <sup>a</sup>	0.20 <sup>a</sup>	0.16 <sup>b</sup>
Magnesium (%)	0.21 <sup>a</sup>	0.17 <sup>b</sup>	0.14 <sup>c</sup>
Potassium (%)	3.07 <sup>a</sup>	2.05 <sup>b</sup>	2.56 <sup>c</sup>
Phosphorous (%)	0.12 <sup>c</sup>	0.22 <sup>a</sup>	0.12 <sup>b</sup>

Ash (%)

1.60<sup>b</sup>8.89<sup>a</sup>8.90<sup>a</sup>

Means followed by the same alphabets on the same row are not significantly different at 0.05 level

### Decomposition and nitrogen release pattern of selected leaf samples.

The decomposition rate of leaf samples of three leguminous species is presented in table 2. Decomposition rate of the samples was highest in *Spondias mombin* with the lowest remaining weight of 2.92g and 3.16g in above and below ground litter bags respectively at week 10. The decomposition rate in the *Albizia* species was similar with residual undecomposed litter for both above and below-ground as (16.89g) and (18.43g). However, in *Albizia ferruginea*, it was 10.08g and 16.70g respectively. Generally, decomposition rate was higher in below ground than above ground in all the leaf samples.

**Table 2: Decomposition rate of leaf samples of three leguminous species**

Time(week)	<i>Albizia zygia</i>		<i>Albizia ferruginea</i>		<i>Spondias mombin</i>	
	AG(g)	BG(g)	AG(g)	BG(g)	AG(g)	BG(g)
2	30.86	26.84	32.7	31.43	17.05	19.00
4	28.49	25.88	28.91	31.4	16.64	14.36
6	26.11	21.67	26.95	23.07	6.00	9.27
8	18.30	21.07	25.15	20.91	4.80	7.45
10	16.89	18.43	10.08	16.70	2.92	3.16

AG- Above ground, BG- Below ground,

\*The mean value is the value of weight remaining. The initial weight subjected to decomposition experiment was 50g at week 0

The results of percentage nitrogen released pattern of the three leguminous three species are presented in table 3. Nitrogen released from leaf of *Albizia ferruginea* and *Spondias mombin* (33.43% and 28.27%) mulches in above ground placement was higher at week 2 than that of *Albizia zygia* (24.84%). Also, the below-ground placed mulch of *Albizia ferruginea* (33.43%) and *Albizia zygia* (27.43%) had a higher rate of N-release compared to *Spondias mombin* (24.28%) at week 2. At week 10, *Albizia ferruginea* leaf (61.59%) had the highest nitrogen mineralization at the above ground placement, this was followed by *Albizia zygia* (55.94%) and lowest in *Spondias mombin* (52.18%). Nitrogen mineralization in the below ground placement was highest in *Spondias mombin* (67.32%), followed by *Albizia ferruginea* (61.59%) and lowest in *Albizia zygia* (40.39%).

**Table 3: Percentage Nitrogen released pattern of three leguminous three species**

Time(week)	<i>Albizia zygia</i>	<i>Albizia ferruginea</i>	<i>Spondias mombin</i>
------------	----------------------	---------------------------	------------------------

	AG(%)	BG(%)	AG(%)	BG(%)	AG(%)	BG(%)
2	24.84	27.43	33.43	33.43	28.27	24.28
4	62.62	66.31	46.23	59.03	32.25	16.31
6	63.72	68.9	33.43	35.99	20.3	24.28
8	58.53	68.9	56.47	53.91	44.21	56.16
10	55.94	40.39	61.59	61.59	52.18	67.32

AG- Above ground, BG- Below ground

#### Effects of added tree mulch on maize growth (girth and height) and yield

The results of the effects of added tree mulch on maize growth (girth height) and yield are presented in table 4. The growth rate of maize plant (total height and girth) were measured. The weight of maize grains obtained per plot was used to represent the yield. The application of *Albizia zygia* mulch significantly increased height growth, girth and weight of maize grains. Mean height of 132.5, 119.5, 105.7 and 93.7cm were recorded for *Albizia zygia*, *Albizia ferruginea*, *Spondias mombin* and the control respectively. *Albizia zygia* had the highest mean girth of 6.7cm and the lowest was recorded for the control (5.1cm). No significant effect was observed between plots with *Albizia ferruginea* and *Spondias mombin* on girth growth. Mean grains weight ranged from 32.5-72.7g. The lowest value of 32.5g was recorded for the control and highest value for *Albizia zygia* (72.7g).

**Table 4 : Effects of added tree mulch on maize growth (girth and height) and yield**

Mulch/Treatment	Mean Height(cm)	Mean Girth(cm)	Mean Grains Weight(g)
<i>Albizia zygia</i>	132.5 <sup>a</sup>	6.7 <sup>a</sup>	72.7 <sup>a</sup>
<i>Albizia ferruginea</i>	119.5 <sup>b</sup>	6.1 <sup>b</sup>	41.8 <sup>b</sup>
<i>Spondias mombin</i>	105.7 <sup>c</sup>	5.9 <sup>b</sup>	40.2 <sup>b</sup>
Control	93.7 <sup>c</sup>	5.1 <sup>c</sup>	32.5 <sup>b</sup>

Means followed by the same alphabets on the same column are not significantly different at 0.05 level

## DISCUSSION

### Foliar analysis for macro-nutrient determination

Mulches of *Albizia zygia*, *Albizia ferruginea* and *Spondias mombin* are statistically different in their macro-nutrients compositions. The differences in the mulch macro-nutrient composition can be adduced to the differential effects on the growth and yield rate of the test crop. It can also be attributed to the differences in decomposition rate of the mulches (Awal & Khan, 2000).

### **Patterns of mulch decomposition and N-mineralization.**

The recycling of nutrients through breakdown of tree biomass, mostly litterfall or pruning, but also root that is added to the soil, is one of the essential acceptances of agroforestry. The quantity and nutrient content of the biomass introduced, as well as the rate at which it decomposes will obviously affect the extent of the advantages derived. Soil mulching using organic material is a method used to protect soil water while also assisting in maintaining a steady soil temperature inside the root system of crops (Samaila *et al.*, 2011).

The results of this study showed that *Spondias mombin* had a single-phase decomposition and nitrogen release pattern. Unlike the other mulches, which had an initial slow decomposition rate, which was followed by second phase of comparatively higher decomposition rate. No significant difference was found between the rate of decomposition of the mulches on the surface and below the surface. Soil incorporated mulch is generally observed to decompose faster than surface-placed mulch (Egbe *et al.*, 2012). The absence of significant differences in decomposition rate of *Spondias mombin*, *Albizia zygia* and *Albizia ferruginea* mulches placed-below and on the soil surface therefore is interesting. There were more termites activities during the experiment, therefore the relatively large mesh size (5mm) of the litter bags used might have resulted in the loss of mulch through the activities of termites. However, the differences in decomposition and N-mineralization rates may also be due to narrow C:N ratio of the materials and this is attributed to factors such as polyphenols and lignin + polyphenol to N ratio (Gentile *et al.*, 2009) and also to the variation in the macro-nutrients composition of the materials. Sugihara *et al.*, (2012) hypothesized that during decay of plant residues, individual biochemical components are lost at a rate that is proportional to the amount of each component present. Using several decomposability indicators, differences in decomposition rate have been linked to substrate quality. The relative amounts of labile and recalcitrant materials, the nutritional content of tissue, the presence of inhibitory compounds, and the concentration of biochemical components such as lignin and cellulose are all factors employed in these indexes (Gentile *et al.*, 2009).

The rate of N-release from *Albizia zygia* was highest during the first six weeks after application, when 60-70% of the mulch N was mineralized. Despite the differences in initial chemical compositions of the species, there was no difference between them in their N release characteristics. The reverse was the case in *Spondias mombin* mulch, there was slower rate of N-mineralization in the first six weeks after which there was a rapid release of nitrogen. In many forest species and in some legumes species, it has been recognized that nutrient rich leaf materials may be slow to release N and that this may be associated with lignin concentration. Vallis and Jones (1973) observed that *D.intortum* was slow to release N. He reported that it might be attributed to the presence of polyphenols in the materials. This is plausible since polyphenols are reactive compounds that can form stable polymers with many forms of N (Gentile *et al.*, 2009). Decomposition rate of *Spondias mombin* at the first six weeks was highest.

However, mineralization was low unlike the other mulches where the rate of decomposition and N-mineralization was simultaneous.

### **Effects of added tree mulch on maize growth (girth and height) and yield**

Mulch application had influence on soil properties giving rise to significantly better root growth and yield of maize compared to no mulch treatment (control) due to increase soil water content resulting from reduced evaporation and increased infiltration. Generally, plant height was highest under mulch treatment compared to the unmulch, this might be due to moisture retention in the soil and decomposition of organic matter in the soil. According to Koller *et al.*, (2013) soil biota increase under mulched soil environment thereby improving nutrient cycling and organic matter build up over a period of several years.

The height growth of maize was influenced mostly by *Albizia zygia* mulch with mean height growth of (132.5cm), followed by *Albizia ferruginea* (119.5cm) and lowest was recorded in the control plot (93.7cm). The girth growth of maize treated with *Albizia zygia* was more pronounced (6.7cm) and it was lowest in the control plot (5.1cm). The decomposition and N-mineralization rates would expect *Spondias mombin* to influence the growth and yield of maize than other mulches. However, reverse is the case in this study. This is because the mulch decomposed and the nutrient was locked up (humification) during the time maize needed the nutrients for growth and development. However, plots treated with *Albizia zygia* mulch were significantly influenced by the mulch application. Thus, after six (6) weeks, the nutrients were released into the soil (immobilization). Myers *et al.*, (1994) suggested that manipulation of mineralization rates through mixtures of plant residue with varying qualities could form the basis of practical management systems for efficient use of nutrients and for minimizing losses.

### **Conclusion and Recommendations**

The study shows that the legume tree leaf mulch (*Albizia ferruginea*, *Albizia zygia* and *Spondias mombin*) varied in their macro-nutrient compositions. It was also observed that below ground mulch placement decomposed faster than the above ground mulch placement in the selected leguminous leaf samples. The indigenous nitrogen fixing tree legumes were found to increase the growth and productivity of maize in the study area and hence it is recommended for use as mulching material to supplement the use of high input resource such as inorganic fertilizer. Specifically, *Albizia zygia* was found to have outstanding performance on height, girth and total yield of maize. It is therefore

recommended to poor resource farmers to allow *Albizia zygia* to thrive in and around their farms so that the leaf could be used as mulch for crop production.

## REFERENCES

- Adesuyi FE, Arinzechi CI, Iyagin FO, Omomoh BE, Akinbowale AS & Dafiewhare HB (2018) Effects of leaf mulch application of *Acacia crassicarpa* Benth., *Albizia adianthifolia* (Schum.) W. Wight and *Albizia zygia* (DC.) J. F. Macbr. on soil properties. *Tropical Plant Research* 5(3): 370–374
- Awal MA & Khan MAH (2000): Mulch induced ecophysiological growth and yield of maize. *Pakistan Journal of Biological Sciences* 3: 61–64.
- Egbe, E.A., Fonge, B.A., Mokake, S.E., Besong, M., Fongod, A.N., (2012). The effects of green manure and NPK fertilizer on the growth and yield of maize (*Zea mays* L.) in the mount Cameroon region. *Agriculture and Biology Journal of North America* 3(3); 82-92
- Gentile R, Vanlauwe B, Van Kessel C, Six J (2009) Managing N availability and losses by combining fertilizer-N with different quality residues in Kenya. *Agr Ecosyst Environ* 131:308–314.
- Fraterrigo, J. M., Turner, M. G., Pearson, S. M. and Dixon, P. (2005). Effects of past land use on spatial heterogeneity of soil nutrients in southern Appalachian forests. *Ecological Monographs*, 75(2): 215 - 230.
- Koller, R., Rodriguez, A., Robin, C., Scheu, S. & Bonkowski, M. (2013) Protozoa enhance foraging efficiency of arbuscular mycorrhizal fungi for mineral nitrogen from organic matter in soil to the benefit of host plants. *New Phytologist*, 199, 203– 211.
- Lu, Y.; Wang, R.; Zhang, Y.; Su, H.; Wang, P.; Jenkins, A.; Ferrier, R.C.; Bailey, M.; Squire, G.(2015) Ecosystem health towards sustainability. *Ecosyst. Health Sustain.* 1, 1–15.
- Myers, R. J. K., Palm, C. A., Cuevas, E., Gunatilleke, T. U. N and Brossard, M. (1994): The synchronization of nutrient mineralization and plant demand. In Woolmer, P.L. and Swift, M. J.ed. *The Biological Management of Tropical Soil Fertility*. John Wiley and Sons, Chichester, 81-136.
- Onyekwuelu, J.C., Reinhard, M. and Bernd, S. (2007). Tree species diversity and soil status of two natural forest ecosystems in lowland humid tropical rainforest region of Nigeria. Conference on International Agriculture Research for Development. Tropentag 2007

Samaila, K.I.; Marcus, N.D and Momale, S.B (2011) Efficient Management of Resources Wastewater Application in Dry Season Farming In Kaduna Urban Area, Nigeria: *Agricultural Journal: Medwell Journals Pakistan*, pp. 188-193.

Sugihara S, Funakawa S, Kosaki T (2012) Effect of land management on soil microbial N supply to crop N uptake in a dry tropical cropland in Tanzania. *Agr Ecosyst Environ* 146:209–219.  
doi:10.1016/j.agee.2011.11.008

Vallis, I., & Jones, R. J. (1973). Net mineralization of nitrogen in leaves and leaf litter of *Desmodium intortum* and *Phaseolus atropurpureus* mixed with soil. *Soil Biology and Biochemistry*, 5(4), 391-398.

UNDER PEER REVIEW