

Hydatid cyst of the Interventricular septum

Case report and literature review

Abstract

Hydatidosis is a frequent pathology which is still endemic in the kingdom of Morocco. Cardiac localization of hydatid cysts is rare representing 0.5 to 2% of all clinical forms of this condition.

Hydatid cysts are usually observed in the left ventricle whilst interventricular septum localization is exceptional. The principal diagnostic and preoperative investigations are serology,

imaging by transthoracic echocardiography and computed tomography. The

authors report a case of hydatid cyst of the interventricular septum in which transthoracic echocardiography (the

key investigation in all cases of hydatid cyst) computed tomography and nuclear magnetic resonance

imaging (MRI) was contributive to the diagnosis. Computerized tomography scan (CT-scan) investigation also aided in the diagnosis of secondary localization of the hydatid cyst especially in the lungs and brain tissues. Diagnostic or therapeutic delay can lead to

severe and even fatal complications.

Introduction

Hydatidosis is a frequent pathology which is still endemic in the kingdom of Morocco [1].

Cardiac localization of hydatid cysts is rare (0.5% to 2%). The revealing symptomatology is unspecific. The most frequent cardiac localizations are the free wall of the left ventricle (LV) and the septum. The auricles can also be affected. Diagnostic or therapeutic delay can lead to severe and even fatal complications [2,3,4].

Only few series of cardiac hydatid disease and primary hydatid disease of the pericardium have been published in the literature. We report a case of hydatid cyst of the interventricular septum revealed by an intracardiac mass.

Case Report

We report the case of a 32-year-old young man of rural origin admitted for exploration of NYHA stage II dyspnea accompanied by palpitation and atypical chest pain, occurring especially on low intensity effort.

The patient has a history of epilepsy since the young age of 10 years under medical treatment with prescribed drugs like tegretol 40mg per day and gardenal 50mg per day. He is also diagnosed hypertension with a likely nephropathy origin under dual therapy based on amlodipine 10mg and indapamide 2, 5mg per day.

The somatic examination is without abnormalities. The electrocardiogram (ECG) showed electrical left ventricular hypertrophy (LVH) with apico-lateral biphasic T wave repolarization disorder (Figure 1). The chest X-ray reveals a round left para-cardiac opacity at the level of the lower lobe of the lung on the same side




Figure 1: Electrocardiogram: Sinus rhythm: electrical LVH and biphasic T waves in the apico-lateral region




Figure 2: Chest X-ray ; Frontal view: limited round opacity localized at the lower lobe of the left lung without filling of the left costo-diaphragmatic cul-de-sac or parenchymal abnormality (arrow).

Transthoracic echocardiography showed a cystic-like oval formation located in the interventricular septum wall (IVS) measuring 30mm x 15mm with border edges protruding into the right ventricular cavity with no other signs of cardiac involvement (figure 3).



Figure 3: Transthoracic echocardiography: A- long-axis parasternal view: Presence of a cystic formation in the IVS wall measuring 30mm x 15mm (arrow) with borderline edges. B- 4 chambers view: shows the same mass protruding into the ventricular chamber.

Biological assessment showed hypereosinophilia without any other anomaly.

Based on the rural origin of the patient, radiology and echocardiographic findings, hydatid cyst of the heart is suspected whereby a hydatid serology by ELISA carried out proved positive.

The radiology findings of multiple localizations of hydatid cyst were then completed by a cardiac MRI which concluded a cystic mass with a fibrous capsule of the interventricular septum measuring 50mm x 30mm, of hydatid origin without intra-LV obstruction. This mass does not enhance in the late phase and is not vascularized in the arterial phase. Absence of extracardiac anomaly apart from a 3 mm pericardial effusion

blade opposite the lower wall of the LV/RV (figure 4).




Figure 4: Cardiac MRI, transversal cross-section: Cystic mass with fibrous capsule of the IVS measuring 50mm x 30mm (red arrow) with an effusion blade next to the inferior wall of the LV/RV (blue arrow).

An assessment of extension of the cystic disease including extracardiac assessment by cerebral and thoraco-abdomino-pelvic CT scan (TAP CT scan) revealed 4 cerebral parenchymal formations largely calcified and hypodense with annular enhancement related to hydatid cysts. CT scan showed left lung lesions of hydatid origin (figure 5 and 6), left renal hypodense patch on chronic pyelonephritic kidney and small aspect of the left renal artery




Figure 5: TAP CT scan at the thoracic level: Cross section passing through C5:

A- Parenchymal window: Stage I simple right cyst of the lower lobe of the left lung (Blue arrow).

B- Bone window: Cystic lesion of the interventricular septum (red arrow)



Figure 6: TAP CT scan at the thoracic level: Longitudinal section: Simple hydatid cyst

Stage I of the lower lobe of the left lung (arrow)

The patient was refused surgery due to a poor prognosis of his general condition, who died after hemodynamic complications by anaphylactic shock one month after the diagnosis was confirmed despite the isolated oral medical treatment by albendazole at a dose of 15mg/kg/day.

Discussion

The hydatid cyst of the heart is secondary to the intracardiac development of *Echinococcus granulosus*. The intermediate host of this parasite is sheep. Humans, the incidental host, become infected by ingesting embryonated eggs which soil a dog's coat or food. The embryo, released in the stomach, crosses the wall of the intestine, the hepatic and pulmonary filter to reach the left heart. From there, it can be distributed in the coronary circulation and end its course in the myocardium where it encysts [4].

The left ventricle (LV) being twice as irrigated as the right, hydatid damage is preponderant on the left (left ventricle 60%, right ventricle 10%, interventricular septum 4%, right atrium 4%) [5,6,7]. The development of intracardiac hydatid cyst is usually subepicardial for localizations in the left heart and subendocardial for those in the right heart, given the low-pressure regime of the right cavities. This explains the possibility of intra-pericardial rupture [5,6] in left localizations hydatid of the heart into the right cavities of the heart.

The clinical presentation of hydatid cyst of the heart (HCH) is very unspecific. The notion of exposure to dogs and sheep, the endemic context or the personal history of other hydatid localizations should suggest the diagnosis [4] as in our case.

The ECG may be normal or show ST segment elevation or depression in the event of myocardial compression or cyst rupture [7]. Biologically, hypereosinophilia can be noted. Hydatid serology is of great diagnostic value but it suffers from false negatives and cross-reactions with other parasitosis [4,9].

Attila has shown that transthoracic echocardiography can reveal unilocular or multivesicular cysts with well-defined contours, with uniform anechoic content, which may calcify, and hypoechoic with multiple septa. Infected hydatid cysts can sometimes appear as echogenic lesions resembling secondary cardiac tumors [7, 10]. More detailed images can be obtained

by transesophageal echocardiography. MRI and CT scan have better sensitivity and specificity than echocardiography in evaluating pericardial masses and analyzing their relationship to surrounding structures. Pulmonary and mediastinal hydatid cystic lesions can also be found [9,11,12]

The evolution of hydatid cyst of the heart can lead to severe complications such as [4]:

- intracardiac rupture with systemic or pulmonary embolic accidents, anaphylactic shock,
- intrapericardial rupture,
- conductive disorders (especially septal location),
- myocardial ischemia

For this, the treatment of hydatid cyst of the heart must be rapid. It is based on surgical excision. In our case, the patient was rejected for surgery because of the multiple cerebral and pulmonary localizations of the hydatid cyst,

which was complicated by hemoptysis after rupture of hydatid cyst in the lungs and cerebral engagement, followed by death.

Certain studies showed that only high-risk patients or those with a small calcified cyst (indicating a dead cyst) and

without hemodynamic complications should be considered for conservative therapy [8, 13]. The

option of isolated medical treatment has also been proposed as an alternative to surgery in certain cases such as in the elderly or in the event of refusal of surgery [4] as in our case. A combination of corticosteroids, H1 and H2 receptor blockers, and benzodiazepines may be advocated for anaphylactic coverage [14, 15].

Therapy is often in oral form with albendazole, a dose of 15mg/kg/day.

Conclusion

Cardiac hydatid is a very rare manifestation of hydatid disease. It can have fatal complications of intracardiac rupture, myocardial infarction, ventricular tachycardia,

conduction disorder such as AV- bloc in septal localization, sudden cardiac arrest or systemic pulmonary embolism.

The diagnosis is suggested by the identification of cysts in a multi septate lesion, presence of hydatid sand and calcification of the walls. Transthoracic echocardiography is the tool of choice for the diagnosis of cardiac hydatid cysts due to its availability, high sensitivity, and ability to concurrently analyze hemodynamic consequences. CT scans and MRI can help pinpoint the location of the lesion and detect other extracardiac localizations.

This case describes an unusual localization of *E. granulosus* in the heart with brain and lung involvement associated with a poor prognosis.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

References

1. Omar Ouzzad, Hicham Kechna, Mohamed Karim Moudden, Khalid Chkoura, Sidi Mohamed Hanafi. Arrêt cardiaque au cours d'une chirurgie de kyste hydatique du foie Pamj ;2015; 6909; p. 22-32
2. **Edlira Rruci, Ketjon Menkshi, Saimir Kuçi, Afërdita Veseli, Gentian Vyshka**
Cardiac Hydatid Cyst: A Case Report
Global Journal of Medical Case Reports; 2022; 2; p. 1-4
3. Alaa S. Bahjat, Ahamd Mohammad Sharif Tahir, Ayad Ahmad Mohammed
Hydatid cyst of the heart with mitral valve stenosis Case report
Ann Med Surg (Lond). 2020; p. 49: 49–52.
4. Bhagwati Pant, Ananthkrishnan Ramesh, Raja Selvaraj
Hydatid cyst of the interventricular septum – A rare cause of heart block
Indian Pacing Electrophysiol J. 2019; 19: p. 79–80.
5. Kosecik M, Karaoglanoglu M, Yamak B. Pericardial hydatid cyst presenting with cardiac tamponade. Can J Cardiol. 2006; 22: p. 145-7.
6. Sascha Kahlfuß, Robert Rainer Flieger, Torsten Kai Roepke, Kadir Yilmaz
Diagnosis and treatment of cardiac echinococcosis
Heartjnl 2016; 102; p. 1348-1353
7. S K MUTHU, A THIAGARAJAN, S GOVINDARAJAN, C M A COUTINHO, W J GUNAWARDENA, T S SWAMINATHAN, Hydatid cyst of the cardiac interventricular septum: report of two cases

UNDER PEER REVIEW

8. Yuksel Besir, Arif Gucu, Suleyman Surer, Orhan Rodoplu, Mehmet Melek, Omer Tetik, Giant cardiac hydatid cyst in the interventricular septum protruding to right ventricular epicardium.
Indian heart journal 2013; 65; p. 81-83
9. J. Bouassida, C. Pricopi, G. Mangiameli, A. Arame, J.B. Auliac, B. Gorbatai, M. Riquet, F. Le Pimpec Barthes. **Compression cardiaque d'origine hydatique.**
Rev de Pneumol clin 2018; 676 :03-009
10. Ceviz M, Becit N, Kocak H. Infected cardiac hydatid cyst.
Heart 2001;86: p. 13.
11. Naqibullah Foladi, Farhad Farzam, Mer Mahmood Shah Hoshang, Mohammad Saboor Rastin, Mohammad Tahir Aien,
Incidental Left Ventricular Myocardial Hydatid Cyst – A case report
Radiol Case Rep. 2022; 17: p. 496–501.
12. Yan F, Huo Q, Abudurehman M, Qiao J, Ma S, Wen H. Surgical treatment and outcome of cardiac cystic echinococcosis.
Eur J Card Thor Surg 2015; 47: p. 1053–8.
13. R, Eryilmaz S, Yazicioglu L, Eren NT, Durdu S, Uysalel A, et al. Surgery for cardiac hydatid disease: an Anatolian experience.
Anadolu Kardiyol Derg 2003; 3: p. 8–44
14. Koppen S, Wejda B, Dormann A, Seesko H, Huchzermeyer H, Junghanss T. Anaphylactic shock caused by rupture of an echinococcal cyst in a 25-year-old asylum seeker from Georgia.
Dtsch Med Wochenschr 2003; 128: p. 663–6.
15. W. OUECHTATI, K. BAJJAR, S. SIDHOM, L. BEZDEH, A. BELHENI. Kyste hydatique du coeur - À propos d'une observation
Cardiologie pratique.com, axis santé: 2011 :p. 1-6

Abbreviation

AV-BLOC: Atrio-ventricular bloc
CT-SCAN: Computerized Tomography Scan
ECG: Electrocardiogram
ELISA: Enzyme Linked Immuno Sorbent Assay
LV: Left Ventricle
LVH: Left Ventricular Hypertrophy
LV/RV: Left ventricular/ Right ventricle
MRI: Magnetic Resonance Imaging
TAP CT scan: Thoraco-abdomino-pelvic CT scan