

ABSTRACT:

Aims: To report a very unusual delayed complication after tubularised incised plate repair with preputioplasty for a distal penile hypospadias

Presentation of Case: A 2^{1/2}-year-old boy presented with a distal penile hypospadias. He underwent an uneventful tubularized incised plate repair with preputioplasty (on parents' request) in September 2018. 1 month after surgery there was a small preputial ulcer on the ventrum. Preputial retraction was done under general anaesthesia and the meatus and neourethra were normal. However, at 2 years the child presented with a fistula. During repair it was found that the neourethra was normal and the meatus was communicating with the prepuce giving the appearance of a fistula. Prepuce was retracted and the inner and outer prepuce layers were closed. Recovery was uneventful.

Discussion: Urethral fistula is the commonest complication after hypospadias surgery, the usual site being the neourethra. Suture line dehiscence and secondary phimosis are the commonest complications of preputioplasty. A fistula between the neomeatus and the prepuce occurring after 2 years is extremely uncommon. Treatment is separating the prepuce from the meatus and meticulously closing the two preputial layers.

Conclusion: A very unusual complication is reported after preputioplasty. Causes and treatment are discussed.

INTRODUCTION: Hypospadias repair is a delicate challenging surgery wherein the commonest complication that can occur are urethral fistula, meatal stenosis and diverticula of the neourethra. We report a very unusual delayed fistula after hypospadias surgery.

CASE REPORT: A 2^{1/2}-year-old boy underwent a tubularized incised plate repair with preputioplasty (on parents' request) for a distal penile hypospadias in Sept 2018. The

postoperative period was uneventful and the urethral stent was removed on the 8th postoperative day. The child voided well in a good stream through the neo meatus. At 1-month follow up there was a small preputial ulcer and the parents were unable to retract the prepuce. Gentle preputial retraction and meatal calibration were done under general anaesthesia 2 months after surgery. At 3 months follow up the child was voiding well and the preputial ulcer had healed. In Sept 2020 the child presented with voiding from a fistula. On examination there was preputial stenosis and a fistula in the distal prepuce.(Fig 1)The child was taken for repair of the fistula. Under caudal block, the prepuce was retracted. To our surprise the neomeatus was adherent to the prepuce giving the appearance of a urethral fistula.(Fig 2)The prepuce was completely retracted and separated from the meatus and the inner and outer preputial skins were closed. A 6 French stent was kept. Post opt period was uneventful. The stent was removed on the 4th post-operative day.

- **DISCUSSION:** Hypospadias is one of the commonest congenital anomalies of the male genital system the incidence of hypospadias is about 1 to 2 in 200 to 300 live births to 3 in 1000 live births⁽¹⁾. The commonest site is the distal penile or a coronal hypospadias⁽²⁾. The surgical goals of hypospadias repair are full straightening of the penis, formation of a hairless urethra of uniform caliber and adequate size, positioning of the meatus at the tip of the glans and normal penile appearance with minimum complications⁽¹⁾. A preputioplasty is performed to avoid making the child conscious about a surgical procedure having been done⁵. The TIP repair described by Snodgrass in 1994 is easy, has good cosmetic results and has minimum complications. A preputioplasty can be combined with this technique to provide a normal penile appearance. The preputioplasty can be done for the following reasons: 1. Religious. 2. Psychological benefit for the child.3. Foreskin plays an important role in normal sexual intercourse.4. It serves as a tissue reserve if the hypospadias fails.⁽³⁾ Reconstruction of the prepuce should be done only if the prepuce can be approximated without tension in the midline at the site of the coronal groove⁽³⁾. A preputioplasty is also the best method to correct the most visible part of the malformation and obtain a normal looking

penis. By adding a preputioplasty, there is no added risk of urethral complications and the incidence of complications of a preputioplasty is the same as circumcision⁶. The usual complications of any hypospadias surgery are urethral fistula, meatal stenosis, urethral diverticula, flap necrosis, penile torsion, infections, persistence chordee and complete dehiscence.⁽²⁾ the commonest being urethral fistula and meatal stenosis.⁽²⁾ However, the common complications of the TIP repair are meatal stenosis and fistula⁽²⁾ Whereas the complications of a preputioplasty as a procedure are dehiscence and secondary phimosis. The prevalence of these may be from 0% to 30%. Preputial dehiscence may be partial or complete. If a partial dehiscence occurs between the proximal and distal segments which have healed properly and completely, it can lead to an immediate preputial fistula. Successful closure of a preputial fistula has been reported in patients with a preputial fistula or complete dehiscence⁷. However, in a 20-year review and metaanalysis of preputial reconstruction, there is a 8% risk of specific complications (dehiscence of the prepuce and secondary phimosis needing circumcision) but there is no increase in the complications of the urethroplasty⁹. Gunjan Shoor et al presented their results of preputioplasty in tip urethroplasty in 48 children. The reported complications were dehiscence, irregular preputial hood, ventral tethering and preputial tightness. No case of preputial fistula was seen.⁸

The urethrocutaneous fistula may occur anywhere along the neourethra. The usual site of the fistula is in the neourethra usually proximally⁽⁴⁾. However, the other common sites include the site of the original meatus, glans penis, at the coronal level in tubularization urethroplasty and at the site of anastomosis in flap urethroplasty.⁽⁴⁾

A fistula between the neomeatus and the prepuce occurring as a delayed complication is extremely uncommon and has not been reported. Usually postoperatively, the first retraction of the foreskin is delayed for 2 to 12 months until there is adequate maturation of the scar.⁶

Even after this time period, reluctance to retract the prepuce, poor hygiene and adhesion of the prepuce to the meatus followed by infection can give rise to this delayed complication.

Treatment would be separation of the prepuce from the meatus followed by closing the inner and outer preputial layers in 3 or 4 layers. In a situation where there is difficulty in retracting the prepuce or reluctance, circumcision should be offered.

CONCLUSION: A case of delayed formation of a meato cutaneous fistula after TIP repair and preputioplasty is presented for its rarity. The etiology and the treatment options are discussed.

REFERENCES:

1. Radi Alsharbaini, MD b and Hamdi Almaramhy, FRCS Snodgrass urethroplasty for hypospadias repair: A retrospective comparison of two variations of the technique
Journal of Talibah University Medical sciences (2014) 9(1), 69=73
2. Warren t. Snodgrass Tubularized incised plate hypospadias repair: indications, technique, and complications Urology 54: 6–11, 1999
3. Amilal Bhat Preputioplasty in hypospadias repair Chapter 23 Surgical techniques in paediatric and adolescent Urology
Publisher: J P Brothers September 2019
4. Management of Post Hypospadias Urethral Fistula
In book: Current Concepts of Urethroplasty Chapter 5 Pg. 47-60 July 2011
Authors: Sherif M K Shehata ,Mohamed Hashish
5. Pradyumna Pan A prospective study comparing modified foreskin reconstruction versus circumcision with tubularized incised plate urethroplasty for distal and mid penile hypospadias *Journal of Pediatric Urology* August 3, 2020

6. Yves Heloury opposing views journal of urology vol. 191, 17-19, January 2014)
7. Castagnetti M, Bagnara V, Rigamonti W, Cimador M, Esposito C, Preputial reconstruction in hypospadias repair, Journal of Pediatric Urology Vol 13, Issue 1, Pg102=109 February 01, 2017(2016)
8. Shoor G, Sugandhi N, Acharya SK, Chakraborty G, Teckchandani N, Dixit A, Kour H, Bagga D, Outcomes Of Preputioplasty In Patients Undergoing TIP Urethroplasty (Tubularization Of Incised Urethral Plate) For Distal and Mid Penile Hypospadias, Journal of Pediatric Urology VOLUME 16, ISSUE 3, P319.E1-319.E7, JUNE 01, 2020
9. Marco Castagnetti, Michele Gnech, Lorenzo Angelini , Waifro Rigamonti , Vincenzo Bagnara and Ciro Esposito. Does Preputial Reconstruction Increase Complication Rate of Hypospadias Repair? 20-Year Systematic Review and Meta-Analysis Frontiers in paediatrics vol 4 article 41, April 2016

ACKNOWLEDGEMENTS: None

COMPETING INTERESTS: None

AUTHORS' CONTRIBUTIONS:

Author 1: Managed the literature search

Author 2: Wrote the draft of the manuscript