

Original Research Article

Influence Of Irrigation Water Deficit On Forage Yield, and Water Utilization Efficiency For Sorghum and Cowpea Forage Crops

ABSTRACT

The present study was carried out at Sakha Agricultural Research Station (clayey soils) during 2018 and 2019 to find out the role of water deficit on yield and some water relations for sorghum and cowpea summer forage crops. Three irrigation levels were executed; 100% (full irrigation), 90% (10% deficit) and 80% (20% deficit). Results showed that increasing water deficit, markedly led to decreasing fresh, dry yield and plant height at the three cuts in the two seasons. Regarding water relations, irrigation level with 90% (10% deficit) resulted in several advantages; almost same yield as recorded under full irrigation, 10% water saving as well as the highest values of water productivity (WP) and productivity of irrigation water (PIW). The mean values of fresh sorghum forage yield can be arranged in descending order as; 2917.0 > 2857.5 > 2561.0 kg.plot⁻¹, while the corresponding values of the dry weight were; 381.6 > 367.8 > 319.6 kg.plot⁻¹. Plant height for cowpea slightly affected with different irrigation treatments. On the contrary, both fresh and dry yields were significantly affected with water level or deficit irrigation. Moreover, the effect was highly significant in connection with the combined analysis. In this regard, mean values of fresh yield descending arranged as; 1502.5 > 1447.0 > 1321.5 kg.plot⁻¹ for treatments A(control), B (90%) and C (80%), respectively. The corresponding values regarding dry yield are 253.5, 241.0 and 215.4 kg.plot⁻¹. Comparing the deficit treatments with the control irrigation (Trt. A), overall mean decrease in fresh yield was 3.7% for treatment B (90%) and 12% for treatment C (80%). While, the corresponding decrease in dry yield were 4.9% and 15.0%. We can conclude that forage yield of sorghum and cowpea not affect by water irrigation deficit about 10% at Middle North Nile Delta area.

Keywords: Forage crops, sorghum, cowpea, water deficit, water productivity, productivity of irrigation water.

INTRODUCTION

Egypt has unique features regarding its water status, it is the solely country worldwide that its agricultural production is mainly under irrigation i.e., irrigated agriculture due to the prevailing aridity conditions. Nile River is the main water resource for Egypt with its inlets outside the boundaries of the country. Capita share from water per annum becomes less than the water

poverty edge of 1000 m³ and it decreasing rapidly towards the scarcity line of less than 500 m³ in the few coming decades. At this situation, it is difficult to make progress in any national economic sector.

In addition, Egypt is facing a pronounced water shortage, particularly for irrigating summer crops. In other words, irrigation is the main critical factor in agricultural production. Agricultural irrigation is the main sector in water consumption with more than 80% from the annual national water supply. Therefore, water productivity becomes one of the main targets under the umbrella of "**effective on-farm irrigation management**". Meaningfully, **more crop per drop**.

Regarding to irrigation management under water deficit status becomes necessity

Crop productivity and its water functions under the prevailing water shortage are among the first priorities of many researches in the world such as; {1,2,3,4,5}

{6} demonstrated that the consumptive use (ET) of 110 to 130-day sorghum crops range between 450 and 750 mm, depending on evaporative demand. Seasonal water use is higher for late maturing genotypes because of longer growing periods. They also stated that surface irrigation methods of furrow, border, basin or corrugation are commonly practiced. Water deficit and drought are the most limiting factors affecting plant growth; reduce crop production and threatening food security in the world {7}

On the other hand, sorghum and cowpea are among other summer crops for animal feeding stuff. Summer forage crops are essential to provide the necessary such animal feeding. Animal feeding stuff is lack in summer due to the less cultivated area in comparison to that cultivated with the winter principal forage crop of Egyptian clover.

Sorghum *Sorghum bicolor L. Moench* is among the most important forage crops in summer season. A successful forage sorghum breeding program to develop hybrid sorghum, which had more tolerant for unsuitable environment and more productivity {8}.

Cowpea *Vigna unguiculata L.* is an important source of protein in human diet {9} and is equally important as nutritious fodder for livestock {10}. As a legume fodder, it provides high quality forage rich in protein with

14-24% {11}. It has ability to tolerate drought and can withstand heat and can utilize soil moisture efficiently as it has a well-developed root system {12} {13} and {14}. So, the aim of this work was to find out the response of yield and some water functions for sorghum and cowpea forage crops to water deficit irrigation owing to produce "more per less".

Materials and Methods

A field experiment was carried out during the two successive summer for sorghum and forage cowpea at the Research Farm of Sakha Agricultural Research Station 2018 and 2019 seasons forage. The site is located in middle North of Nile Delta area with 30°-57 N latitude, 31°-07E longitude with an elevation of about 6 meters above mean sea level. Table 1 represents the climatic elements of the area during the two field trial seasons. The soil of the site is clayey in texture as shown in Table 2.

Table (1). Climatic elements ; air temperature (T, C°), relative humidity (RH, %), wind speed (U₂, m.sec⁻¹) and evaporation pan (Ep, mm.d⁻¹).

A.1 st season, 2018							b-2 nd season, 2019						
Month	T, C°			RH, %	U ₂ , m.sec ⁻¹	Ep, mm.day ⁻¹	Month	T, C°			RH, %	U ₂ , m.sec ⁻¹	Ep, mm. day ⁻¹
	max	min	Mea n					max	min	Mea n			
May-18	31.2	23.8	27.5	60	1.1	6.3	May-19	31.9	25.4	28.3	57.2	0.79	6.8
June "	32.6	25.3	28.9	62	1.14	7.8	June "	33	27	30	65.7	1.19	8.5
July "	34.2	25.4	29.8	67	1.03	7.4	July "	33.5	28.4	30.9	70.5	0.97	9.4
Aug. "	33.9	25.3	29.6	67	0.88	6.4	Aug. "	34.2	29.1	31.7	70.8	0.8	6.8
Sept. "	32.8	23.5	28.2	66	0.8	5	Sept. "	32.4	27.9	30.2	68.2	0.89	5.9
Oct. "	29.5	20.6	25.1	66	0.67	3.2	Oct. "	30.2	26.7	28.5	70.8	0.66	3.8
Mean	32.4	24	28.2	65	0.94	6	Mean	32.5	27.4	29.9	67.2	0.88	6.9

Physical and chemical characteristics of the studied site.

Soil samples were collected from different depths: 0-15, 15-30, 30-45 and 45-60 cm to determine soil-water constants of field capacity (F.C) and permanent wilting point (PWP) according to {15} as well as soil bulk density (Db). Physical properties of particle size distribution were determined according to {16} The obtained analysis indicated that the soil is clayey in texture as shown in Table 2. Chemical properties of total soluble salts, soil reaction (pH), both soluble cations and anions were also determined according to {17} Sulphate (So₄⁻) was computed by the difference between soluble cations and anions, both in meq L⁻¹ as tabulated in Table 3.

Table (2):Particle Size distribution and soil-water constants of the studied experimental site.

Soil depth, cm.	Particle size distribution			Texture class	F.C, %	W.P, %	AW, %	Db, Mg/m ⁻³
	Sand, %	Silt, %	Clay, %					
0 – 15	19.5	27.6	52.9	Clay	46.40	25.21	21.19	1.14

15 – 30	20.4	27.4	52.2	Clay	41.38	22.49	18.89	1.18
30 – 45	24.9	25.7	49.4	Clay	38.50	20.92	17.58	1.22
45 – 60	25.6	26.8	47.6	Clay	36.90	20.05	16.85	1.30
Mean	22.6	26.9	50.5	Clay	40.80.	22.17	18.63	1.21

Where: F.C, % = soil field capacity, W.P, % = wilting point, AW, % = available soil water and Db, Mg.m⁻³ = soil bulk density.

Table (3): Chemical properties of the experimental site:

Soil depth, cm	Ec, dsm ⁻¹	PH (1: 2.5) soil water suspension	Soluble ions, meqL ⁻¹							
			Cations				Anions			
			Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	CO ₃ ⁻	HCO ₃ ⁻	Cl ⁻	SO ₄ ⁻⁻
0-15	1.78	8.30	6.20	3.20	8.20	0.27	0.00	5.20	8.10	4.57
15-30	2.40	8.15	9.63	4.70	9.54	0.22	0.00	4.76	7.94	11.38
30-45	2.56	8.20	9.20	5.95	10.34	0.20	0.00	4.53	7.38	13.78
45-60	2.94	8.01	10.70	6.70	11.86	0.19	0.00	3.78	7.56	18.11
Mean	2.42		8.93	5.14	9.99	0.22	0.00	3.57	7.75	11.96

Agronomic practices:

The cultivated summer forage crops were sorghum (*Sorghum bicolor L. Moench cv. piper black*) and cowpea (*Vigna unguiculata cv. Baladi*), from Forage Corps Res. Dept, grown in a randomized complete block design (R.C.B.D) with three replications for each crop. Each crop grown in nine plots, every plot area was 140m² [(25 m long x 5.6 m width (8 ridges, 0.7m wide))].

Sowing date (S) in the two growing seasons were 20th and 24th of May in the two seasons, respectively. Seeding rates were 20 kg/fad for sorghum and 30 kg /fed for cowpea. Twenty unit of nitrogen in the form of urea applied in the three times for each crop, at the second irrigation and after the first and the second cut. Harvesting date (H) in the two seasons at three cuts were taken for every crop, the first, second and third cuttings were taken after 55 , 100 and 140 days from sowing in the two seasons, respectively.

Data recorded for each crop in the three cuts in the two seasons at the middle plot (4 ridges with 5 m long which mean 2.8 x 5 m for fresh forage yield and dry forage yield/kg (14m²) and converted to kg/plot (140m²), as well as plant height cm.

Statistically analysis:

The recorded data were statistically analyzed according to {18} Treatment means were compared by the least significant difference test using {19}. Homogeneity of experiments variances were computed according to {20}.

Irrigation treatments:

Three irrigation treatments based on water deficit levels were assigned as follows:

- A- Irrigation with 100% of crop water requirements (control).
- B- Irrigation with 90% of control A (10% deficit).
- C- Irrigation with 80% of control A (20% deficit).

Data collection:

a. Water parameters:

• Irrigation water (IW)

Irrigation water was controlled and measured by irrigation pump with a discharge of 18 L per second. Control irrigation treatment A was irrigated till the end of the assigned cultivated plot.

• Consumptive use (CU)

Actual consumptive use (CU) or so-called crop evapotranspiration (ET) was computed based on soil moisture depletion (SMD) in the effective root zone of 60 cm as follows {21}.

$$Cu = \frac{FC - \theta}{100} * \frac{Db}{Dw} * d$$

Where:

CU = consumptive use or actual crop- water consumed, cm.

FC = percent soil moisture content on weight basis at field capacity

θ = percent soil moisture content on weight basis before irrigation as well as at harvesting.

Db = soil bulk density, Mg.m⁻³

Dw = density of water = 1

d = effective root zone of 60 cm.

- **Crop-water functions**

- i. Water productivity (WP):**

Water productivity as defined by {22} is the parameter of crop-water functions which reflects the capability of water consumed by growing crop in producing marketable yield as follows:

$$WP = Y/CU$$

Where:

WP = productivity of crop- water consumed

Y = marketable yield, and

CU = consumption use.

- ii. Productivity of irrigation water (PIW):**

This parameter of PIW refers to the capability of applied irrigation water in producing marketable yield as defined by {22}.

$$PIW = Y/ IW$$

Where:

PIW = productivity of applied irrigation water,

Y = marketable yield, and

IW = applied irrigation water.

3. RESULTS AND DICUSSION

3-a. Mean squares:

Analysis of variance of two years and their combined for fresh and dry forage yield and plant height for each summer forage crop were presented in Tables (4) and (5). Data revealed that differences among irrigation levels were highly significant and significant effects for most fresh and dry forage yield as well as plant height at the three cuts and total yield for the two forage crops in the two seasons and combined analysis were detected similar results obtained by {23}, {24}and {25}.

Table (4) Analysis of variance for two years and combined analysis for yield and plant height of forage Sorghum under water deficit.

2018 season													
S.of.V.	df	Fresh forage yield				Dry forage yield				Plant height cm			
		Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
Replication	2	75	8.33*	1.33*	46.3	0.053**	1.5	0.343*	4.33*	4.11	14.3	28	3.81
Irrigation	2	7275.0*	10675**	5772.00**	69937**	217.56**	372.4 **	400.33* *	2915.29**	352.44**	523.0*	241.33**	363.15**
Error	4	637.5	508	131.33	111	8.42	17	7.87	95.8	5.77	32.3	6.33	6.53
Total	8	-	-	-	-	-	-	-	-	-	-	-	-
2019 season													
S.of.V.	df	Fresh forage yield				Dry forage yield				Plant height cm			
		Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
Replication	2	36.33	22.33*	6.33*	144.0*	0.260*	0.040**	0.04	0.823*	2.778*	5.78	7.11	3.02
Irrigation	2	13129.0**	18157.0**	4501.0**	100069.0**	378.02**	627.19**	216.19**	3501.31**	287.44*	560.77**	214.77*	338.01**
Error	4	578.8	515	231.33	3822	10.2	12	5.52	80.6	20.8	20.9	18.1	17.2
Total	8	-	-	-	-	-	-	-	-	-	-	-	-
combined													
S.of.V.	df	Fresh forage yield				Dry forage yield				Plant height cm			
		Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
Replication	2	8.167	24.5	1.5	35.2	0.23	1	0.25	3.85	6.50*	9.06	20.2	2.53
(Y)Year	1	1002152**	125000.0**	19404.5**	285012.50**	522.29**	926.651**	477.405**	990.125**	206.722**	2403.55**	600.88**	168.05*
Error	2	103.2	6.17	6.167	155	0.08	0.6	0.14	1.31	0.39	11.5	14.9	4.3
Irrigation(I)	2	19974.5**	28333.5**	10233.5**	168655.5**	584.44**	983.138**	602.105**	6402.88**	630.167**	1078.22**	455.72**	698.087**
(Y)x(I)	2	429.5*	499*	39.5*	1350.50**	11.1	17	14.4	13.7	9.72	5.56	0.389*	3.08

Error	8	608.2	512	181.33	1967	9.31	14	6.7	88.2	13.3	26.6	12.2	11.9
Total	17	-	-	-	-	-	-	-	-	-	-	-	-

*, ** P ≤ 0.05 and 0.01 respectively

Table (5) Analysis of variance of two years and combined analysis for yield and plant height of forage Cowpea under water deficit.

2018 season													
S.of.V.	df	Fresh forage yield				Dry forage yield				Plant height cm			
		Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
Replication	2	4	16	6.333*	72.33*	0.173*	0.013**	0.1	0.360**	0.190*	0.163*	0.653*	0.298*
Irrigation	2	1351	2496.0*	2547.00*	18811.0*	53.3	112.00*	139.08*	880.36*	13.9	49.6	43.3	33.04
Error	4	256	224	174	1944	12.4	9.1	9.2	91	18.5	9.96	15.1	14.24
Total	8	-	-	-	-	-	-	-	-	-	-	-	-
2019 season													
S.of.V.	df	Fresh forage yield				Dry forage yield				Plant height cm			
		Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
Replication	2	52	9.00*	4.00*	147	0.154**	1.377*	0.493*	4.960*	0.97	0.33	0.39	0.493
Irrigation	2	3411.0*	6247.00**	2179.00**	39361.0*	136	248.58*	104.160*	1414.359**	30.8	39	128.170*	58.96
Error	4	460	155	98	1911	23.2	34	13	204	23.8	10.8	16.5	16.55
Total	8	-	-	-	-	-	-	-	-	-	-	-	-
combined													
SOV	df	Fresh forage yield				Dry forage yield				Plant height cm			
		Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
Replication	2	42	24.50*	10.167*	211.167*	0.3	0.6	0.5	4	1.01	0.43	1.022*	0.776

(Year(Y	1	214512.50**	186050.00**	8580.500**	642978.00**	3397.902**	4418.940**	364.50**	11160.178**	121.680**	345.845**	5070.245**	448.002**
Error	2	14	0.5	0.17	8.2	0.03	0.8	0.1	1.3	0.16	0.07	0.02	0.016
Irrigation(I)	2	4524.50 NS	8319.500**	4718.00**	51591.50**	179.507**	347.113**	241.980**	2263.219**	43	88.265*	160.235**	90.141*
(Y)x(I)	2	238	424	8	1112	9.54	13	1.3	32	1.68	0.37	11.255*	1.871
Error	8	358	189	136	1928	17.8	22	11	148	21.1	10.4	15.8	15.4
Total	17	-	-	-	-	-	-	-	-	-	-	-	-

*, ** P ≤ 0.05 and 0.01 respectively

NS; not significantly different

3.b. Mean Performance:

3.b.1. Forage Sorghum

In comparison to the full irrigation control treatment A, the obtained data (Table 6) demonstrated that water deficit irrigation has a substantial impact on feed sorghum crop output. Table 6 shows the average performance of plant height, fresh and dry forage yield for sorghum. According to {26} and {27}., plant height was influenced by irrigation levels.

The lowest applied water, 80 percent water irrigation (deficient 20%), had the smallest plant height at the three cuts in the two trial seasons and combined analyses. Plant height can be ordered in descending order based on the levels of applied irrigation water that resulted from the tested water deficit treatments: no deficit (control) > 10% deficit > 20% deficit, respectively.

For fresh and dry sorghum forage yields, the highest mean values were obtained from the control normal watering. On the other hand, the low-level irrigation which associated with deficit treatments caused the low mean values at the three cuts in the two seasons and their combined data. The mean values of fresh sorghum forage yield can be arranged in descending order as; 2917.0 > 2857.5 > 2561.0 kg.plot⁻¹, while the corresponding values of the dry weight are; 381.6 > 367.8 > 319.6 kg.plot⁻¹ (area of plot = 140m² = 0.033fed = 0.014 ha). The stated values for treatments A (100% no deficit), B (90% of A) and C (80% of A), respectively.

Comparing with treatment A, 98% and 88% were obtained under deficit treatments B and C, respectively. Regarding dry weight, the corresponding percentages are 96% and 84%.

Therefore, by implementing 10% water deficit irrigation (Trt. B), only 2 and 4% (Table 11) decreasing in fresh and dry yield of sorghum were recorded comparing to the traditional irrigation and the decrease were insignificant, as combined data. While C treatment were recorded 12 and 16% decreased in fresh and dry yield compared to control A treatment and highly significant decreased, (Table 11) it can be concluded that 10% water deficit irrigation on forage sorghum not affect on fresh and dry yield. General mean of total and dry yield is very high in the second season for two forage crops, it is due to may be climatic elements (Table 1) The obtained results are nearly as obtained by {28}, {29}, {30}, {31} and {32}.

Table (6) Effect of irrigation levels in two years and combined data for fresh and dry forage yield and plant height of forage Sorghum under water deficit.

2018 season												
	Fresh forage yield kg/plot				Dry forage yield kg/plot				Plant height cm			
Irrigation levels	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
100%	1045.0	750.0	554.0	2349.0	125.4	101.3	89.7	316.4	261.6	185.0	164.0	203.5
90%	1032.0	735.0	542.0	2309.0	121.8	97.0	86.7	305.5	251.3	165.0	157.0	191.1
80%	931.0	660.0	481.0	2072.0	107.2	83.8	75.4	266.4	235.6	157.3	146.0	179.6
Grand Mean	1002.6	715.0	525.6	2243.33	118.1	94.0	83.9	296.1	249.5	169.1	155.6	191.4
Significance	*	**	**	**	*	**	*	*	**	*	**	**
LSD0.05	61.1	43.9	33.5	138.0	8.9	8.3	8.53	25.75	10.5	12.0	10.8	8.2
LSD0.01	-	72.9	55.6	228.8	-	13.7	-	-	17.4	19.9	-	13.7
2019season												
	Fresh forage yield kg/plot				Dry forage yield kg/plot				Plant height cm			
Irrigation levels	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
100%	1624.0	1288.0	573.0	3485.0	188.4	167.4	91.0	446.8	248.3	161.3	155.6	188.4
90%	1596.0	1254.0	556.0	3406.0	182.6	160.4	87.2	430.2	242.6	158.3	151.3	184.1
80%	1437.0	1120.0	493.0	3050.0	158.0	126.7	76.0	372.8	228.0	144.0	132.6	168.2
Grand Mean	1552.3	1220.6	540.6	3313.6	176.3	138.8	84.7	416.6	239.6	154.5	146.5	180.2

Significance	**	**	*	**	*	**	*	*	*	**	*	**
LSD0.05	50.4	49.4	40.0	139.6	17.2	13.3	8.3	38.8	-	-	7.1	13.7
LSD0.01	83.5	81.9	-	231.4	-	22.1	-	-	-	-	11.8	-
Combined												
	Fresh forage yield kg/plot				Dry forage yield kg/plot				Plant height cm			
Irrigation levels	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
100%	1334.5	1019.0	563.5	2917.0	156.9	134.3	90.3	381.6	255.0	173.1	159.8	196.0
90%	1314.0	994.5	549.0	2857.5	152.2	128.7	86.9	367.8	247.0	161.6	154.1	187.6
80%	1184.0	890.0	487.0	2561.0	132.6	111.3	75.7	319.6	231.8	150.6	139.3	173.9
Grand Mean	1277.5	967.8	533.1	2778.5	147.2	124.7	84.3	356.3	244.6	161.8	151.1	185.8
Significance	**	**	**	**	**	**	**	**	**	**	**	**
LSD0.05	46.5	38.8	30.6	115.3	11.4	9.2	6.9	27.3	10.1	8.2	5.3	6.6
LSD0.01	67.7	56.5	44.6	167.7	16.6	13.4	10.1	39.8	14.7	12.0	7.8	9.6

*, ** P ≤ 0.05 and 0.01 respectively

NS; not significantly different

3.b.2. Cowpea

The mean performance of plant height, fresh and dry cowpea forage yield in the two seasons as well as the combined data are presented in Table 7. Plant height slightly affected with different irrigation treatments. On the contrary, both fresh and dry yields were significantly affected with water level or deficit irrigation. Moreover, the effect was highly significant in connection with the combined analysis. In this regard, mean values of fresh yield descending arranged as; 1502.5 > 1447.0 > 1321.5 kg plot⁻¹ for treatments A (control), B (90%) and C (80%) from A, respectively. The corresponding values regarding dry yield are 253.5, 241.0 and 215.4 kg plot⁻¹. Comparing the deficit treatments with the control irrigation (Trt. A), overall mean decrease in fresh yield was 3.7% for treatment B (90%) and 12% for treatment C (80%). While, the corresponding decreases in dry yield were 4.9% and 15.0%, [33]. The decreasing in treatment B (deficit 10%) were insignificant effect of fresh and dry forage yield as combined data, while treatment c (20% deficit) irrigation were highly significant effect fresh and dry forage yield. This finding could be attributed to the less applied irrigation water under 80% level comparing with 90%. The obtained results emphasized the results that reported by [34], [35] and [36]. Obtained data also showed that average plant height of cowpea forage crop is not affected with irrigation level in the two seasons of study. Combined analysis showed that the highest plant height was resulted with control irrigation i.e., no deficit and vice versa for deficit irrigation treatments. The average combined values arranged in descending order as 67.8 > 65.6 > 60.2 cm. for treatments A, B and C, respectively.

In general, increasing water deficit stress notably affected the physiological processes and nutrient uptake by fodder cowpea crop. Therefore, growth, development and yield are negatively affected [37]

Table (7) Effect of irrigation levels in two years and combined data for fresh and dry forage yield and plant height of forage Cowpea under water deficit.

2018												
	Fresh forage yield kg/plot				Dry forage yield kg/plot				Plant height cm			
Irrigation levels	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
100%	470.0	424.0	407.0	1301.0	73.8	74.6	78.2	226.6	67.0	63.0	57.0	62.3
90%	459.0	408.0	389.0	1256.0	71.4	70.6	73.6	215.6	65.7	61.1	54.8	60.5
80%	429.0	368.0	350.0	1147.0	65.6	62.6	64.8	193.0	62.8	55.2	49.6	55.9
Grand Mean	452.6	400	382.0	1234.6	70.8	69.2	72.2	211.7	65.1	59.7	53.8	59.5
Significance	NS	*	*	*	NS	*	*	*	NS	NS	NS	NS
LSD0.05	-	33.9	29.8	99.9	-	6.8	6.8	21.6	-	-	-	-
LSD0.01	-	-	-	-	-	-	-	-	-	-	-	-
2019												
	Fresh forage yield kg/plot				Dry forage yield kg/plot				Plant height cm			
Irrigation levels	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
100%	700.0	642.0	362.0	1704.0	103.6	108.4	68.4	280.4	73.0	54.0	93.0	73.3
90%	679.0	615.0	344.0	1638.0	99.2	102.8	64.4	266.4	71.3	52.0	88.9	70.7

80%	634.0	553.0	309.0	1496.0	90.4	90.6	56.8	237.8	66.8	47.0	80.2	64.6
Grand Mean	671.0	603.3	338.3	1612.6	97.7	100.6	63.2	261.5	70.3	51.0	87.3	69.5
Significance	*	*	**	*	NS	*	*	*	NS	NS	*	NS
LSD0.05	48.5	28.1	22.4	99.1	-	13.2	8.2	32.4	-	-	9.2	-
LSD0.01	-	-	37.2	-	-	-	-	-	-	-	-	-
Combined												
	Fresh forage yield kg/plot				Dry forage yield kg/plot				Plant height cm			
Irrigation levels	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Total	Cut 1	Cut 2	Cut 3	Average
100%	585.0	533.0	384.5	1502.5	88.8	91.5	73.3	253.5	70.0	58.5	75.0	67.8
90%	569.0	511.5	366.5	1447.0	85.3	86.0	69.0	241.0	68.5	56.5	71.8	65.6
80%	531.5	460.5	329.5	1321.5	78.0	76.6	60.8	215.4	64.5	51.1	64.9	60.2
Grand Mean	561.8	501.6	360.1	1423.6	84.0	84.9	67.7	236.6	67.7	55.3	70.5	64.5
Significant	NS	**	**	**	**	**	**	**	NS	*	**	*
LSD0.05	35.6	25.9	21.95	82.6	7.7	8.7	8.7	22.9	-	6.07	7.48	7.38
LSD0.01	51.8	37.6	31.94	120.3	11.5	12.7	12.7	33.3	-	-	10.8	-

*, ** P ≤ 0.05 and 0.01 respectively NS; not significantly different

3.c. Water relations

3.c.1. Applied irrigation water (IW).

Seasonal values of applied irrigation water for forage sorghum and cowpea during the two growing seasons of the study are presented in Table 8. It is clear from the presented data that the control water (Trt. A) has the highest values compared to water deficit treatments B and C. In this regard, mean values of I.W. for sorghum could be arranged in descending order as 83.44 > 75.25 > 69.47 cm. for treatments A (100%), B (90%) and C (80%), while the corresponding values for cowpea are 78.51 > 72.93 > 68.30 cm.

Therefore, by comparing I.W. for water deficit treatments with the control (Trt. A), mean water saving for sorghum counted with 9.8 and 16.7% under 90 and 80% water deficit, respectively. For cowpea crop, the corresponding values are 7.1 and 13.0%.

It should be notified that choosing the suitable level of water deficit to be executed under surface irrigation in the clayey soils should be linked with the decreasing in crop yield. In other words, factors should be taken into consideration; water saving and crop productivity as well as net return per m³ of water.

The obtained findings are in the same direction with that reported by {38}, {29} and {6}

Table 8. Seasonal irrigation water (IW, cm) for Sorghum and Cowpea as affected with different irrigation treatments.

Treatment	Crop	1 st season cm.	2 nd season cm.	Mean	
				Cm	m ³ /fed
A (100%)	Sorghum	84.75	82.13	83.44	3504.5
	Cowpea	82.63	74.38	78.51	3297.4
B (90%)	Sorghum	78.29	72.20	75.25	3160.3
	Cowpea	75.12	70.73	72.93	3062.9
C (80%)	Sorghum	71.65	67.29	69.47	2917.7
	Cowpea	70.59	66.00	68.30	2868.6

3.c.2. Consumptive use (CU)

Seasonal values of crop consumed water (CU) for different treatments in the two growing seasons are presented in Table 9. Data show that control treatment A (100%) has the highest values of seasonal CU, while the lowest ones are for 80% water deficit (Trt. C).

In this regard, mean values of seasonal CU for sorghum crop are arranged in descending order as 66.8 > 61.7 > 59.1 cm, while the corresponding ones for cowpea are 62.8 > 59.8 > 58.1 cm for treatments A, B and C, respectively.

This finding is owing to the less applied irrigation water to water deficit irrigation treatments which caused a decreasing in consumed water by the growing plants.

The obtained results are in a good agreement with that obtained by {38}, {29}, {39} and {6}.

Table 9. Seasonal consumptive use (CU, cm) for Sorghum and Cowpea as affected with different irrigation treatments.

Treatment	Crop	1 st season cm.	2 nd season cm.	Mean	
				Cm	m ³ /fed
A (100%)	Sorghum	67.80	65.70	66.75	2803.5
	Cowpea	66.10	59.50	62.80	2637.6
B (90%)	Sorghum	64.20	59.20	61.70	2591.4
	Cowpea	61.60	58.00	59.80	2511.6
C (80%)	Sorghum	60.90	57.20	59.05	2480.1
	Cowpea	60.00	56.10	58.05	2438.1

3.c.3. Crop-water functions

Crop- water functions reflect the capability of consumed and/or irrigation water in water productivity, herewith:

3.c.3.1. Water productivity (WP)

The obtained data of WP in kg/m³ consumed water for different treatments are presented in Table 10. It is clear that the 90% water level (10% deficit) of treatment B has the highest values and vice versa for the non-deficit irrigation treatment A of full watering. This could

be attributed to that CU as the dominator of crop-water function is the highest for full irrigation. In other words, there is a reverse relation between CU and WP.

3.c.3.2. Productivity of Irrigation water (PIW)

It is clear that PIW in kg/m³ irrigation water took the same trend of WP (Table 8). The highest applied irrigation water, the lowest PIW taking into consideration the obtained yield.

Therefore, the obtained results of WP and PIW are in the same trend with that reported by {40}, {41}, {42}, {43}, {29} and {39}.

3.c.3.4. Maximizing productivity of water unit

As mentioned before, the water level of 90% (Trt. B) produced nearly the highest yield as full irrigation (Trt. A, 100%) with slight decrease in fresh and dry yield with 2.0 and 4.0% for sorghum and 3.7 and 4.9%, for cowpea (Tables 6&7).

In that regard, mean values of WP and PIW under treatment B with 10% deficit for fresh and dry yield of sorghum and cowpea are the highest as presented in Table 10.

Table 10. Mean values of fresh, dry yield, consumptive use (CU), irrigation water (IW), water productivity (WP) and productivity of irrigation water (PIW) for sorghum and cowpea under different treatment.

Treatment	Crop	Yield, Kg/fed		CU, m ³ /fed	IW, m ³ /fed	Productivity, Kg/m ³ water			
		Fresh	Dry			WP		PIW	
						Fresh	Dry	Fresh	Dry
A (100%)	Sorghum	88385.1	11562.5	2803.5	3504.5	31.5	4.1	25.2	3.3
	Cowpea	45525.8	7681.1	2637.6	3297.4	17.3	2.9	13.8	2.3
B (90%)	Sorghum	86582.3	11144.3	2591.4	3160.3	33.4	4.3	27.4	3.5
	Cowpea	43844.1	7302.3	2511.6	3062.9	17.5	2.9	14.3	2.4
C (80%)	Sorghum	77598.3	9683.9	2480.1	2917.7	31.3	3.9	26.6	3.3
	Cowpea	40041.5	6526.6	2438.1	2868.6	16.4	2.7	14.0	2.3

Therefore, one m³ consumed water produced 33.4 and/or 4.3 kg as fresh and dry yield for sorghum, and 17.5 and/or 2.9 kg for cowpea. Meanwhile, that one m³ of applied

irrigation water produced 27.4 and/or 3.5 kg as fresh and dry yield of sorghum and 14.3 and/or 2.4 kg for cowpea.

In other words, to produce 1 kg fresh and dry yield of sorghum it consumed 29.9 liter and 232.6 liter, respectively. For cowpea, the values were 57.1 and 344.8 liter, respectively.

Meanwhile, to produce 1kg fresh sorghum, it needs 36.5 liter as irrigation water, while it is 285.7 liter for dry yield. For cowpea, the values were 69.9 and 416.7 liter for fresh and dry yield, respectively.

4- Percent of decrease:

Percent of decrease of sorghum and cowpea in 2018 and 2019 seasons and combined analysis are presented in Table (11). Water stress fewer than 80% irrigation, reduced total fresh and dry forage yield production for sorghum and cowpea {44}, {24}. In this regard, {45} found that drought stress led to reduced average yield for most major crops by more than 50%. Although, cowpea was tolerant to decreased water of irrigation {46}, {47}, {48}.

Table (11) Percent of decrease of summer forage

Sorghum 2018 season													
Irrigation levels	Fresh forage yield				Dry forage yield				\bar{x}	Plant height cm			Average
	Cut 1	Cut 2	Cut 3	\bar{x}	Cut 1	Cut 2	Cut 3	\bar{x}		Cut 1	Cut 2	Cut 3	
100%	100	100	100	100	100	100	100	100	100	100	100	100	
90%	1.2	2.0	2.2	1.7	2.9	4.2	3.3	3.4	3.9	10.8	4.3	6.1	
80%	9.8	10.2	11.3	10.3	12.0	13.6	13.0	12.8	6.2	4.7	7.0	6.0	
Sorghum 2019 season													
Irrigation levels	Fresh forage yield				Dry forage yield				\bar{x}	Plant height cm			Average
	Cut 1	Cut 2	Cut 3	\bar{x}	Cut 1	Cut 2	Cut 3	\bar{x}		Cut 1	Cut 2	Cut 3	
100%	100	100	100	100	100	100	100	100	100	100	100	100	
90%	1.7	2.6	3.0	2.3	3.1	4.2	4.2	3.7	2.3	1.9	2.8	2.3	
80%	10.0	10.7	11.3	10.5	13.5	21.0	12.8	13.3	6.0	9.0	12.4	8.6	
Sorghum combined													
Irrigation levels	Fresh forage yield				Dry forage yield				\bar{x}	Plant height cm			Average
	Cut 1	Cut 2	Cut 3	\bar{x}	Cut 1	Cut 2	Cut 3	\bar{x}		Cut 1	Cut 2	Cut 3	
100%	100	100	100	100	100	100	100	100	100	100	100	100	
90%	1.5	2.4	2.6	2.0	3.0	4.2	3.8	3.6	3.1	6.6	3.6	4.3	
80%	9.9	10.5	11.3	10.4	12.9	13.5	12.9	13.1	6.2	6.8	9.6	7.3	
Cowpea 2018 season													
Irrigation levels	Fresh forage yield				Dry forage yield				\bar{x}	Plant height cm			Average
	Cut 1	Cut 2	Cut 3	\bar{x}	Cut 1	Cut 2	Cut 3	\bar{x}		Cut 1	Cut 2	Cut 3	
100%	100	100	100	100	100	100	100	100	100	100	100	100	
90%	2.3	3.8	4.4	3.5	3.3	5.4	5.9	4.9	1.9	3.0	3.9	2.9	
80%	6.5	9.8	10.0	8.7	8.1	11.3	12.0	10.5	4.4	9.7	9.5	7.6	
Cowpea 2019 season													

	Fresh forage yield				Dry forage yield				Plant height cm			
Irrigation levels	Cut 1	Cut 2	Cut 3	\bar{x}	Cut 1	Cut 2	Cut 3	\bar{x}	Cut 1	Cut 2	Cut 3	Average
100%	100	100	100	100	100	100	100	100	100	100	100	100
90%	3.0	4.2	5.0	3.9	4.2	5.2	5.8	5.0	2.3	3.7	4.4	3.5
80%	6.6	10.1	10.2	8.7	8.9	11.9	11.8	10.7	6.3	9.6	9.8	8.6
Cowpea combined												
	Fresh forage yield				Dry forage yield				Plant height cm			
Irrigation levels	Cut 1	Cut 2	Cut 3	\bar{x}	Cut 1	Cut 2	Cut 3	\bar{x}	Cut 1	Cut 2	Cut 3	Average
100%	100	100	100	100	100	100	100	100	100	100	100	100
90%	2.7	4.0	4.7	3.7	3.9	6.0	5.9	4.9	2.1	3.4	4.3	3.2
80%	6.6	10.0	10.1	8.7	8.6	10.9	11.9	10.6	5.8	9.6	9.6	8.2

CONCLUSIONS

- Water productivity (WP) for m³ of sorghum and cowpea yield (fresh and dry) under treatment B of 90% level (10% deficit) were 33.4, 4.3, 17.5 and 2.9 kg, respectively.
- Productivity of irrigation water (PIW) for sorghum and cowpea yield (fresh and dry) under treatment B of 90% level (10% deficit) were 27.4, 3.5; 14.3, 2.4 kg, respectively
- By implementing irrigation of 90% water needs i.e., 10% deficit for sorghum and cowpea forage crops, several advantages could be achieved such as; nearly highest yield, 10% water saving, the highest productivity of each unit of either consumed or applied irrigation water.
- More research should be done regarding the performance of crops and their water relations under deficit irrigation with considering the net return per consumption of m³ of water term.

REFERENCES

- 1-**Abu-Zeid, M. and A. Hamdy 2002.** Water crisis and food security in the Arab world: where we are and where do we go. Second Regional Conference on Arab Water, Cairo, Egypt.
- 2-**Seckler, D.U., U. Amarasinghe, D. Molden, R. Da Silva and R. Barker 1998.** World water demand and supply, 1990-2025. Scenarios and issues. Research report 19. IWMI, Colombia, Sri Lanka.
- 3-**Hamdy, A. and C. Lacirignola (1999).** Mediterranean water resources: Major challenges towards the 21st century. Mediterranean Agronomic Institute of Bari, Italy.
- 4-**Corgrove, J.W. and R. F. Rijsberman 2000.** World Water Vision. World Water Council Earthscan Publication, Ltd, UK.
- 5-**IWMI- International water management institute 2000.** Water supply and demand in 2025. Colombia, Sri Lanka.

- 6-Steduto, P., T. C. Hisao, E. Fereres and D. Raes 2012.** Crop yield response to water, FAO Irrigation and Drainage Paper 66.
- 7-Water in a Changing World (2009).** The United Nations World Water Development Report 3–WWDR3 - UNESCO, pp 1–313.
- 8-Duvick,D.N.(1999).**Commercial strategies for exploitation of heterosis.The Genetics and Exploitation of Heterosis in Crops.Wisconsin, USA.P.19-29.
- 9-Karigoudar, A. S and S.S. Angadi, (2005).** Influence of seed rate, row spacing and fertility levels on yield and quality of field bean for forage production. Karnataka J.Agric.Sci., 18(2):489-491.
- 10-Singh B.B;H.A Ajeigbe;S.A Tarawali ;S.Fernandez – Rivera;M.Abubakar (2003).**Improving the production and utilization of cowpea as food and fodder .Field Crops Research ,84:169-177.
- 11-Heuze V.; G. Tran;D.Bastianelli ;F.lebas and P.Noziere (2013).** Cowpea (*Vigna unguiculata*) forage. [http://www.feedipedia.org/node/ 23](http://www.feedipedia.org/node/23)September 2013, 20-42.
- 12-Turk, K.J and Hall, A.E (1980b).** Drought adaptation of cowpea. IV: Influence of drought on water use and relation with growth and seed yield. Agron J 72: 440- 448.
- 13-Handy, A.; R. Ragab and Elisa Scarascia-Mugnozza 2002.** Coping with water scarcity: water saving and increasing water productivity. International Commission on Irrigation and Drainage, Montreal.
- 14-Sambo,B.E.;E.C.Odion ;L.Aiyu and D.A.Labe (2013).**Cowpea *Vigna unguiculata* (L.) Walp) clipping management technology.1: Apotential for fodder Production, sustained growth and food security in the Savannah regions of Nigria.J.Agriculture and Crop Research 1(3):36-45.
- 15-James,L.G. (1988).**Principls of farm irrigation system design.John Willey and Sona Inc.,New York,543.
- 16-Klute,A.C(1986)**Water retention laboratory Methods .In:A.koute(ed)Methods of Soil Analysis,part 1-2nd (ed) Agronomy Monogr.9,ASA,Madison,WI U.S.A,pp .635-660.
- 17-Jackson,M.I, (1973).**Soil chemical Analysis Prentice Hall of India Private,LTD new Delhi.
- 18-Steel, R.G.D. and J.H. Torrie. 1980.** Principles and Procedures of Statistics. A Biometry: cal Approach 2nd Ed: McGraw Hill, New York. PP.232- 349.
- 19-Mstat-C, 1986.**A micro computer program for the design of experiment Michigan State Univ, USA.
- 20-Bartelett, M. S. 1937.** Properties of sufficiency and statistical test. Proc. Roy. Soc., A160: 268-282.
- 21-Hansen, V.;Israelsen,W.,and Stringharm,Q.E(1979).**Irrigation principle and practices,4th (ed.),john Willey and Sons,.USA.

- 22-Bos,M.G.1980.**Irrigation efficiencies at crop production level .ICID.Bulletin 29,2:189-260 New Delhi.
- 23-Blum A (2009)** Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress. *Field Crops Res* 112:119–123.
- 24-Hall, A.E, Thiaw, S, Ismail, A.M and Ehlers, J.D (1997).**Water-use efficiency and drought adaptation of cowpea. Pp. 87-98. In: *Advances in cowpea research*, B.B. Singh et al. (Eds). IITA/JIRCAS, IITA, Ibadan, Nigeria.
- 25-Ludlow, M. M and Muchow, R. C (1990).** A critical evaluation of traits for improving crop yields in water limited environments. *Advances in Agronomy* 43: 107-153.
- 26-Assaeed, A.M. (1994)** Evaluation of some forage sorghum varieties under the condition of central region, Saudi Arabia. *Annals Agric, Sci., Ain shams Univ., Cairo*, 39 (2), 649-654.
- 27-Bakheit, B. R. (1990).** Performance, phenotypic and genotypic stability for green fodder and leafiness in sorghum under different nitrogen fertilization levels. *Assiut Journal of Agricultural Sciences* 1990 Vol.21 No.1 pp.65-77.
- 28-Carrow RN (1996)** Drought avoidance characteristic of diverse tall fescue cultivars. *Crop Sci* 36:371–377.
- 29-Abdel-Fattah, I. M. 2011.** Climate change impacts on maize under surface irrigation with gated pipes in North Nile Delta. M.Sc., Fac. Of Agric., Mansoura Univ.
- 30-Mariola Staniak and Anna Kocon (2015).** Forage grasses under drought stress in conditions of Poland. *Acta Physiol Plant* 37:116 DOI 10.1007/s11738-015-1864-1.
- 31-El-Shamarka, sh. A, M. E. Ibrahiem, F. M.Ali , T.G.EL-Gaafarey (2012).** Studies on yield stability of new forage sorghum genotypes in different environments. *Minufiya j.Agric.Res.Vol.37 No.4 (1):871-880.*
- 32- Ghasemi A, Karim MH, Karim Koshteh and MM Ghasemi 2012.** Green fodder yield performance of different varieties of sorghum grown in an arid region. *International Journal of Agriculture and Crop Sciences* Vol., 4(13), 839-843.
- 33-Hayatu,M. and F.B.Mukhtar (2010).**Physiological responses of drought resistant cowpea genotypes (*Vigna unguiculata(L)Walp*) to water stress .*Bayero j.pure and Applied Sci.,3(2):69-75.*
- 34-Mullen, C. (1999).** Summer legume forage crops: cowpeas, lablab, soybeans. SW Department of primary Industries. Broad acre crops. Agfact P4.2.16.
- 35-IITA, (1989).** Cowpea Research at international Institute for Tropical Agriculture (IITA) GLIP Research Monograph, No. (1) Ibadan Nigeria.
- 36-Madamba, R.; G.J.H.Grubben; I.K.Asante and R. Akromah (2006).** *Vigna unguiculata (L) Walp.* Brink, M. and Belay, G (Eds.), PROTA (Plant Resources of Tropical Africa), Wageningen, Netherlands.

- 37-Zeinab M.Khalil ;Azza Kh.Salem and Fadia M.Sultan(2015).** Water Stress Tolerance of Fodder Cowpea As influenced by various Added levels of Potassium Sluphate.J.soil Sci. and Agric.Eng., Mansura Univ.,Vol.6(2):113-131.
- 38-Doorenbos and Kassam (1979)** stated that surface irrigation is commonly used in alfalfa production. Crop water requirements (ET_m) are between 800 and 1600 mm/ growing period depending on climate and length of growing period.
- 39-Kassab, M. M.; M. A. M. Ibrahim N. J. Talha 2012.** Effect of cultivation method and irrigation rate on productivity and water efficiency of berseem crop grown in North Nile Delta. Minufiya J. Agric. Res. Vol.37 No.4(2):1019-1033.
- 40-Lazridou, Martha and S. D. Koutroubas 2004.** Drought effect on water use efficiency of berseem clover at various growth dtages. 4th International Crop Science Congress. Brisbane, Australia.
- 41-Ali, M. H. and M. S. U. Talukder 2008.** Increasing water productivity in crop production. A Syntheses Agriculture Water Management, pp: 1201-1213
- 42-Abdulaziz, A. M.; M. Saleh and Robin Wardlaw 2009.** Modelling irrigation water management under water shortage and salinity conditions: 1-Evaluation of the current irrigation and drainage management practices in South Kazakhstan. Alexandria Science Exchange J 30 (3), September, pp: 350-371.
- 43-Ouda, Samiha A., R. Abou Elenin and M. A. Shreif 2012.** Simulating the effect of deficit irrigation on Egyptian clover and water productivity. 14th International Water Technology IWTC 14 2010, Cairo, Egypt.
- 44-Mai-Kodomi, Y, Singh, B.B, Myers, O., Yopp, J.H, Gibson, P.J and Terao, T (1999a).** Two mechanisms of drought tolerance in cowpea. Indian J.Genet. 59:309-316.
- 45-Wang et, W.; B.Vinocur and A.Altman (2003).**Plant response to drought, salinity and extreme temperature: towards genetic engineering for stress tolerance.Planta.218:1-14.
- 46-Hall, A.E. (2004).** Breeding for adaptation to drought and heat in cowpea. Eur J Agron 21:447–454
- 47- Dadson RB, Hashem FM, Javaid I, Allen AL and Devine TE 2005.** Effect of water stress on yield of cowpea (*Vigna unguiculata* L. Walp.) genotypes in the Delmarva region of the United States. J. Agron. Crop Sci. 191:210-217.
- 48- Ehlers JD and Hall AE 1997.** Cowpea (*Vigna unguiculata* L. Walp.). Field Crops Res. 53:187-204.