

Original Research Article
**Effect of Canopy Gaps on Growth and Water Use
Efficiency of **Seedlings** of Vulnerable Species, *Hopea
sangal* Korth**

ABSTRACT

Aims: *Hopea sangal* Korth. is listed as vulnerable species and recently its remnant habitat was rediscovered in the degraded forest near the springs area in East Java. In the forest, the regeneration of the seedlings and saplings understory is affected by the heterogeneity in environmental factors especially various light levels due to the canopy gaps. *H. sangal* is considered a shade-tolerant species, hence the establishment of the **seedlings** in its natural habitat occur under the closed canopy. This study aimed to understand the Water Use Efficiency (WUE) and Relative Growth Rate (RGR) of *Hopea sangal* Korth seedlings grown in two different levels of tree canopy shades.

Study design This research was using a completely randomized design, with 9 replications.

Place and Duration of Study: This study was conducted in the Purwodadi Botanic Garden, East Java, between March – July.

Methodology: We used seedlings of *H. sangal* collected from the field in Pasuruan which were planted in plastic pots and acclimatized to obtain seedlings **of** similar size. The microclimate was measured weekly using solarimetri and sling psychrometer at 08.00; 10:00; 12:00; 14:00 and 16:00. RGR was measured by harvesting the seedling and whole plant WUE was measured using the gravimetric method.

Result: We observed the light level during the periodic opening of the canopy gap was significantly different at a specific time (8-12 am), with the highest light intensity was at 10.00 of 333.57 $\mu\text{mol photon m}^{-2}\text{s}^{-1}$. The canopy shade differences did not affect the WUE of seedlings ($P = 0.333$), meanwhile, there was a significant difference in the seedling RGR ($P = 0.025$). The seedlings that were grown under a higher gap and received higher light intensity periodically during a day had higher RGR than those under a closed canopy. The WUE has a positive relationship with the RGR of seedlings ($R^2 = 0.5$; $P < 0.05$).

Conclusions: This study suggested that the *H. sangal* is one of the shade-tolerant species capable of responding to gap opening sunlight. The study also showed RGR of the seedling positively correlated with the WUE of plants, indicating that the seedling could use the water supply efficiently to grow rapidly.

Keywords: **seedlings**, canopy gaps, *Hopea sangal*, Water use efficiency, Relative growth rate

1. INTRODUCTION

One of the plant species collected around the water spring in the district of Pasuruan, East Java by the exploration team of Purwodadi Botanic Garden was the seedlings of a rare plant species, *Hopea sangal*. It belongs to the Dipterocarpaceae family. The name *H. Sangal* was obtained from the identification process based on literature and discussion. The identification results have been verified by experts on Dipterocarpaceae, Prof. Dr Peter S. Ashton, of the Royal Botanic Gardens Kew, Richmond, England. The conservation status of *H. sangal* based on the IUCN (the International Union for Conservation of Nature) is vulnerable A2cd ver 3.1 and current population trend is decreasing [1]. The remaining habitat of this plant species was also declared as a new habitat in the region of Pasuruan (rediscovery) [2]. Based on information obtained from the Herbarium Bogoriense, Biology Center-LIPI, *H. sangal* was also found in the Banyuwangi district in 1898, Malang district in 1934 and Blitar regency in

1935 [3]. The remaining habitat *H. sangal* was newly discovered in the district of Pasuruan, located close to the northern coast of Java, while the three locations where it was first discovered were located closer to the southern coast of the island of Java [2]. In terms of ecology, this information is likely to give us new knowledge about the habitat of *H. sangal* in East Java.



**Figure 1. The tree and canopy of *H.sangal* in its natural habitat
Left: The tree stand; Right: The tallest tree among others, in Pasuruan, East Java**

Dipterocarpaceae in Indonesia is not evenly distributed on each island with most found west of the Wallace Line and northwest of the Indonesian archipelago [4]. The Dipterocarpaceae family consists of more than 500 species in 14 genera and mostly grows well in Asia, especially in Sumatra and Kalimantan [4,5]. Generally, In Java, there are 10 species of Dipterocarpaceae, two of them are from the genus Hopea: *H. celebica* and *H. sangal*. However, the population of *H. sangal* has been declared as a rare species [6,7]. The next editions of the publication by Ashton Wong in 1982 and 1994 also did not mention the existence of *H. Sangal* in East Java.

The new habitat where *H. sangal* was rediscovered is a degraded forest near the spring area in East Java. This remaining habitat becomes a great concern because only one individual tree was found in that area [2]. Moreover, the flowering timing and masting periods of Dipterocarpaceae is seasonal and does not occur throughout the year. Previous studies proposed that the mass flowering of Dipterocarps is associated with the El Nino phases and it evolved as a response to the seasonal tropic dry climate. This reproductive strategy of Dipterocarps is important for seedling survival in tropical forests [8,9].

The tropical forest has heterogeneous canopy structures that affect environmental factors, especially light availability and microclimate which would affect the forest regeneration. The canopy gap has an important role in providing the habitat that is suitable for the seedlings and saplings to grow, hence it affects the species richness and diversity in the forest [10]. The light availability is also different in the type of forest. For example, the primary forest consists of larger trees and higher stratification of canopy layers than the regeneration forest. Therefore the light availability in regenerating forests is higher than in the primary forest. The disturbed and degraded forest has a larger canopy gap and lower stratification layer hence this type of forest would allow the higher radiation of sunlight to the forest floor [11].

The functional adaptation of the plant varies among species. Several plants species could tolerate the low intensity of light and are able to compete with another plants species to survive in the forest and some species only can grow under high light intensity [12,13]. The response of Dipterocarpaceae seedlings to the light intensity is usually different depending on the species. For instance, the responses of two species of Dipterocarpaceae seedlings: *H. helferi* . and *H. odorata* were different to various light conditions[14]. A previous study of *Shorea selanica* also stated that the high percentage of shading would reduce the intensity of sunlight received by plants and it would inhibit growth

processes [15]. *H. sangal* is considered a shade-tolerant species, hence the establishment of the seedling in their natural habitat occur under the closed canopy. An earlier study suggested that *H. sangal* has the highest rank in size-specific height, diameter, and aboveground biomass growth rates performance under low light conditions than 21 species of Dipterocarpaceae, but it had lower growth performance under medium and high light treatment [16].

Earlier studies measuring the growth performance of the *H. sangal* seedling were using the shade house, hence there is still a lack of information about its growth performance under variable or non-constant light intensity. In this study, we used two different levels of canopy shades to replace the shade house. We also measured the microclimate differences between two levels of canopy shades and investigated the relative growth rate (RGR) of the seedlings. The RGR has relation to plants' ability to adapt and compete to survive and it would give information on how plants respond to differences in abiotic factors [17].

Besides investigating the RGR we also observed the water use efficiency of the seedling. The plant's distribution in the natural habitat is related to their various responses to their environment [18]. The response of the plants to water availability is related to their functional strategies. The extreme drought, for instance, changes the plant biomass allocation to reproductive organs [19]. The water deficit would disrupt the balance of chemicals in the plant which results in reduced photosynthesis results or all physiological processes running abnormally. Water crucially impacts the dynamics of plant growth, productivity and reproduction [20–22]. The *Hopea* is distributed mostly in lowland forests that have dry climates or in the dry season of less than 4 months [4,5,23]. This suggested that plants cope with the seasonal water limitation. The new remnant habitat of *H. sangal* was found close to the spring area, in which the presence of the water is abundant. Hence it is important to understand the water use efficiency of this species. Moreover, the previous study proposed that the water use of each species of the Dipterocarpaceae family varied. For instance, *Shorea contorta*, *Parashorea malaanonan*, *Hopea maliboto* and *Hopea plagata*, the species with the lowest water use was *H. plagata* with a transpiration rate was 3.3mm day^{-1} and its water use was 4 kg day^{-1} with the trunk's diameter of 6.6 cm [24]. Besides genetic factors, the water use efficiency for plant growth is also influenced by ecological factors such as temperature, humidity, wind speed, weather, and sunlight. Water use efficiency is used as a link between timber production and management of water use by plants.

Our objective in this study was to understand the physiological performance of *H. sangal* seedlings in two different levels of canopy gaps and microclimate. We tested these hypotheses: (1) there will be significant differences in the RGR and WUE of the seedlings under different canopy gaps and, (2) does the WUE has a relationship with the seedling RGR?

2. MATERIALS AND METHODS

2.1. Materials and Location

The research materials were seedlings of *H. sangal* collected from the field in Pasuruan which were transplanted in plastic pots. Before the research began, the seedlings were acclimatized to obtain seedlings of similar size. Since the seedlings of this vulnerable species were limited, we used 18 seedlings in this study. The dimension of the plastic pots used was 29x20x19 cm and pots were filled with the same medium consisted of soil and compost mixture with a ratio mixture of 3:1.

2.2. Experiment establishment

The selected seedlings were established beneath closed-canopy trees. The canopy gaps were divided into two groups. One group was placed under a closed canopy tree with 94% of shade and the other group was placed at a location slightly more open in a shade of 88%. The percentage of shading (94% and 88%) was obtained from the comparison of the average measurement results with the open or without shade results. Each group consisted of 9 replications.

The research activities started with weighing all the pots and seedlings. The soil from each pot was taken to measure the water content of the soil by the gravimetric method, then all pots were covered with transparent plastic and tied with a rubber band so that the unintended water will not enter the pot and

and prevent evaporation from the soil. The pots were weighed every five days and then re-watered to field capacity.

2.3. Microclimate Measurement

To know the differences in microclimate under the two shading canopy gaps, we collected data on energy light - using solarimetri, temperature, and humidity using a sling psychrometer. The observations were performed weekly with the frequency of observation five times at 08:00; 10:00; 12:00; 14:00 and 16:00.

2.4. Relative Growth Rate

The method used was a completely randomized design with two treatments: 94% shading and 88% shading treatments. The parameters measures were: relative growth rate/RGR ($\text{g g}^{-1} \text{week}^{-1}$), water use (kg month^{-1}), and Water Use Efficiency (WUE) ($\text{g DM kg}^{-1} \text{H}_2\text{O}$). The RGR obtained by the harvesting method and the rate of relative growth was calculated using the following equation [25];

$$\text{RGR} = (\ln W_2 - \ln W_1) / (t_2 - t_1)$$

Where:

\ln = natural logarithm

t_1 = time one

t_2 = time two

W_1 = dry weight of plant in time 1

W_2 = dry weight of plant in time 2

2.5. Transpiration and Water Use Efficiency

The WUE was quantified at whole plant level by using gravimetric technique. The analysis of plants' water use efficiency was calculated for each plant as the ratio of biomass production during a definite growth period to water transpired during the same period. The average value increment of biomass was obtained by comparing the measurement results with the amount of the sample's biomass. Meanwhile, the amount of water that transpired during the experiment was calculated by measuring evaporative water loss from the pots experiment.

2.6. Statistical Analysis

The experiment was analyzed as a completely randomized design with two different shading conditions and 9 replicate seedlings per treatment. The following model was applied in this study:

$$Y_{ij} = \mu + a_i + e_{ij}$$

where Y_{ij} is the measured values for seedling j under the treatment i ; μ = overall mean ; a_i = effect of shading treatment and e_{ij} = error term. The normality and randomness of data were tested using Kolmogorov-Smirnov and Run tests. The normal data were analyzed using variance analysis (ANOVA) with a confidence level of 95%. The relationship between RGR and WUE was analyzed with regression using Statistica 64 software version 13.3.

3. RESULTS AND DISCUSSION

3.1. Microclimate under different canopy gaps

The light intensity, temperature, and relative humidity of both canopy-shaded areas were measured from 08:00 until 16:00. Even though the level of shading under the tree canopy did not have large differences (88% and 94%), interestingly the light level during the periodic opening of the gap was significantly different. It occurred at a specific time (08:00 – 12:00). Under the 88% shaded area, the light intensity increased from 08:00 and reached its highest point at 10:00 and then decreased until 16:00, while in 94% canopy shade, the light increased from 08:00 and reached its highest point at 14:00 and then decreased until 16:00 (Fig 2a). The highest light intensity was at 10:00 of $333.57 \mu\text{mol photon m}^{-2}\text{s}^{-1}$ while at same

time light intensity that entered 94% shade was $17.67 \mu\text{mol photon m}^{-2}\text{s}^{-1}$. The differences in gap size at canopy shades increase in the duration of periods of full sunlight in the understory. Hence, at 8 – 12 am the seedlings under 88% shading received maximum solar energy.





Fig 2. (a) Mean value of light intensity between 08.00 -16.00, while (b) mean value of temperature and relative humidity between 08.00-16.00

Temperature and relative humidity have a negative correlation. If the temperature increases, the humidity will decrease and vice versa. Temperature, humidity, and light intensity affect seedling growth and transpiration affects seedling water use. In both different canopy shading conditions between 88% shade and 94% shade, temperature and humidity did not show a significant difference. The temperature at 08.00 would gradually increase and reached the highest value by 10:00 in both shaded areas. Temperatures descended from 12:00 until 16:00. The highest humidity occurred at noon in the 94% shade area (Fig 2b).

3.2. The seedling growth, Water Use, and Water Use Efficiency

The Relative Growth Rate (RGR) is the plant's ability to produce biomass as a result of assimilation per unit of dry weight per time expressed in $\text{g g}^{-1} \text{week}^{-1}$. The results showed that *H. sangal* seedlings placed in two different amounts of shading have significantly different RGRs with $P = 0.025$. The RGR of *H. sangal* seedlings placed in 88% shade were higher than those placed in 94% shade (Fig 3). The

seedlings grown under 88% shade had RGR $0.02 \text{ g g}^{-1} \text{ week}^{-1}$, while those under 94% shade was $0.01 \text{ g g}^{-1} \text{ week}^{-1}$.






Fig 3. Mean value of RGR of seedlings of *H. sangal* growth in two shading treatments T94= 94% shading and T88 = 88% shading, $P < 0.05$

The significant differences in RGR were due to the difference in the light intensity received by seedlings. Our result showed that seedling RGR was higher under 88% shades canopy than under 94%. A higher RGR value indicates that the seedlings grows faster at the given size. Under the canopy gap in the forest, the responses of shade-tolerant species toward the light level were different. Some shade-tolerant species could not tolerate the higher light intensity during the periodic openings [13]. In the right amount of light intensity, plants grow properly and the growth will increase concurrently as much as sunlight received because assimilates produced during the process of photosynthesis are greatly influenced by the intensity of the sun-lighting [16]. Based on our result, *H. sangal* is one of the shade-tolerant species that had a great response toward the higher light levels during the periodic opening of the canopy gap. It has the capability to respond to the higher sunlight with the higher RGR under 88% of the canopy gaps area.



(a)

(b)

Fig 4. (a) Water use of *H. sangal* seedling under two different treatments at $P > 0.05$, while (b) Water Use Efficiency of *H. sangal* seedlings at $P > 0.05$

The amount of water required by the plants is different for each individual, depending on the plant

Figure 4 : Water Use Efficiency of *H. sangal* seedlings

species and external factors. The data showed that the amount of water use of *H. sangal* seedlings in both areas were not significantly different at $P = 0.333$. The seedlings under 88% shade treatment used a lower amount of water ($0.83 \text{ kg month}^{-1}$) compare to the 94% shade treatment which used $0.93 \text{ kg month}^{-1}$ (Fig 4a). Water use efficiency (WUE) is expressed by the number of grams of the dry weight of plants per kg of water transpired. The seedling with the highest WUE was those grown under 88% shades about $7.6 \text{ g DM kg}^{-1} \text{ H}_2\text{O}$ and 94% was $5.8 \text{ g DM kg}^{-1} \text{ H}_2\text{O}$ (fig 4b). Based on our statistical analysis, the WUE of seedlings grown under different shading was not significantly different ($p > 0.05$).




Fig 5. Relationship between RGR and WUE of *H. sangal* seedlings -the model was significant at $P < 0.05$ and the slope indicated a positive relationship.

The model show that WUE has a positive relationship with the RGR of seedlings ($R^2 = 0.5$; $P < 0.05$). The seedlings that had higher WUE grew faster as showed by the RGR values (fig 5). Water use efficiency is the effectiveness of water use by plants in the transpiration process for their growth or biomass addition, in comparison to the amount of water that is absorbed. Hence it will affect plant growth or biomass. Based on this study, the seedlings could use the water supply efficiently to grow rapidly.

Water in the plants play a large role in physiological processes. Water composes protoplasm of 85-90% of the plant's weight acts as a reagent in the process of photosynthesis and also has a function as a salt and substances solvent for cell growth. Water in the plant's system will be released into the air by the transpiration process. Under the 88% canopy shading, the *H. sangal* seedlings produced more biomass than those that were placed in the 94% shading. Interestingly, this study also found that at the higher RGR, the water use of the seedling was lower compared to 94% shade. Thereby the seedlings of this species could optimize carbon assimilation and reduce water losses. The earlier study also proposed that the amount of Dipterocarpaceae water use is lower than 7 other families of plants with the same diameter and in the same climate [26].

The seedlings of *H. sangal* had faster biomass increment under the higher level of light during a day, illustrating that seedlings require the higher light intensity to maximize carbon assimilation. This study also suggested that the seedlings may be able to compete to grow under lower stratification layer

forest or the larger canopy gaps. The high WUE of *H.sangal* was contributed to its growth and it would benefit the seedlings for surviving the dry month season. Understanding the light requirement and also the WUE of the seedlings will give information on the environmental suitability for the seedlings to grow properly. This information would be needed for further conservation action of this vulnerable species.

4. CONCLUSION

The differences in light intensity during periodic opening under the canopies significantly affected the growth but not the water use and water use efficiency of *H. sangal* seedlings. Based on our study, *H. sangal* is considered one of the shade-tolerant species that had a great response toward the higher light levels during the periodic opening of the canopy gap. High WUE of seedlings positively affected plant growth. It suggested that the seedlings of *H. sangal* are able to optimize the carbon assimilation and reduce water losses, hence seedlings could use the water supply efficiently to grow rapidly.

REFERENCES

1. Pooma, R.; Barstow, M.; Newman, M.F. Hopea Sangal. The IUCN Red List of Threatened Species 2017: E.T31314A2804189.
2. Soejono. Short Communication: Rediscovery of a Remnant Habitat of the Critically Endangered Species, Hopea Sangal, in Pasuruan District, East Java, Indonesia. Biodiversitas 2014, 15, doi:10.13057/biodiv/d150217.
3. Kusuma, Y.W.C.; Wihermanto; Risna, R.A.; Ashton, P.S. Rediscovery of the Supposedly Extinct Dipterocarpus Cinereus . Oryx 2013, 47, doi:10.1017/s0030605313000756.
4. Ashton, P. Dipterocarpaceae. Flora Malesiana, Series 1; Netherlands: Martinus Nijhoff Publishers, 1982.
5. Purwaningsih, P. Review: Ecological Distribution of Dipterocarpaceae Species in Indonesia. Biodiversitas Journal of Biological Diversity 2004, 5, doi:10.13057/BIODIV/D050210.
6. Heyne, K. De Nuttige Planten van Indonesie; Druk. NV Uitgeverij W Van Hoeve 's-Gravenhage/ Bandung, 1950.
7. Backer, C.; Bakhuizen van den Brink Jr., R.. Dipterocarpaceae in Flora of Java I; NVP Noordhoff, Groningen, The Netherlands., 1963.
8. Kurten, E.L.; Bunyavejchewin, S.; Davies, S.J. Phenology of a Dipterocarp Forest with Seasonal Drought: Insights into the Origin of General Flowering. Journal of Ecology 2018, 106, 126–136, doi:10.1111/1365-2745.12858.
9. Chechina, M.; Hamann, A. Climatic Drivers of Dipterocarp Mass-Flowering in South-East Asia. Journal of Tropical Ecology 2019, 35, 108–117, doi:10.1017/S0266467419000087.
10. Hubbell, S.P.; Foster, R.B. Biology, Chance, and History and the Structure of Tropical Rain Forest Tree Communities. In Community Ecology; Diamond, J.M. and C.T.J., Ed.; Harper and Row Publishers: New York, 1986; pp. 314–329.
11. Numata, S.; Yasuda M; Okuda T; Kachi N; Nur Supardi M.N Canopy Gap Dynamics Of Two Different Forest Stands In A Malaysian Lowland Rain Forest. Journal of Tropical Forest Science 2006, 18, 109–116.
12. Yamamoto, S.-I. Forest Gap Dynamics and Tree Regeneration. <http://dx.doi.org/10.1007/BF02767114> 2005, 5, 223–229, doi:10.1007/BF02767114.
13. Canham, C.D. Different Responses to Gaps Among Shade-Tolerant Tree Species. Ecology 1989, 70, 548–550, doi:10.2307/1940200.
14. Lee, D.W.; Oberbauer, S.F.; Johnson, P.; Krishnapilay, B.; Mansor, M.; Mohamad, H.; Yap, S.K. Effects of Irradiance and Spectral Quality on Leaf Structure and Function in Seedlings of Two Southeast Asian Hopea (Dipterocarpaceae) Species. American Journal of Botany 2000, 87, 447–455, doi:10.2307/2656588.
15. Panjaitan, S.; Wahyuningtyas Dan/, R.S.; Ambarwati, D.; By, O./; Abstract, ;; Madya, P.; Penelitian, B.; Banjarbaru, K.; Selatan, K.; Km, J.A.Y.; et al. Pengaruh Naungan Terhadap Proses Ekofisiologi Dan Pertumbuhan Semai *Shorea selanica* (Dc.) Blume di Persemaian. Jurnal Penelitian Ekosistem Dipterokarpa 2011, 5, 73–82, doi:10.20886/JPED.2011.5.2.73-82. Indonesian.

16. Philipson, C.D.; Saner, P.; Marthews, T.R.; Nilus, R.; Reynolds, G.; Turnbull, L.A.; Hector, A. Light-Based Regeneration Niches: Evidence from 21 Dipterocarp Species Using Size-Specific RGRs. *Biotropica* 2012, 44, doi:10.1111/j.1744-7429.2011.00833.x.
17. Dodd, G.L.; Donovan, L.A. Water Potential and Ionic Effects on Germination and Seedling Growth of Two Cold Desert Shrubs. *American Journal of Botany* 1999, 86, 1146–1153, doi:10.2307/2656978.
18. Hernández, E.I.; Vilagrosa, A.; Pausas, J.G.; Bellot, J. Morphological Traits and Water Use Strategies in Seedlings of Mediterranean Coexisting Species. *Plant Ecology* 2010, 207, 233–244, doi:10.1007/S11258-009-9668-2/FIGURES/3.
19. Liu, Y.; Li, Z. Effects of Water Addition on Reproductive Allocation of Dominant Plant Species in Inner Mongolia Steppe. *Frontiers in Plant Science* 2020, 11, 1797, doi:10.3389/FPLS.2020.555743/BIBTEX.
20. Lindroth, A.; Cienciala, E. Water Use Efficiency of Short-Rotation *Salix viminalis* at Leaf, Tree and Stand Scales. *Tree Physiology* 1996, 16, 257–262, doi:10.1093/treephys/16.1-2.257.
21. Medrano, H.; Flexas, J.; Galmés, J. Variability in Water Use Efficiency at the Leaf Level among Mediterranean Plants with Different Growth Forms. *Plant and Soil* 2009, 317, 17–29, doi:10.1007/s11104-008-9785-z.
22. Quan, Q.; Zhang, F.; Meng, C.; Ma, F.; Zhou, Q.; Sun, F.; Niu, S. Shifting Biomass Allocation Determines Community Water Use Efficiency under Climate Warming. *Environmental Research Letters* 2020, 15, 094041, doi:10.1088/1748-9326/ABA472.
23. Rahayu, E.M.. Ex Situ Conservation of Dipterocarpaceae in Bogor Botanic Garden. *The Indonesian Botanic Gardens Bulletin* 2009, 12, 69–77.
24. Dierick, D.; Hölscher, D. Species-Specific Tree Water Use Characteristics in Reforestation Stands in the Philippines. *Agricultural and Forest Meteorology* 2009, 149, 1317–1326, doi:10.1016/J.AGRFORMET.2009.03.003.
25. Hoffmann, W.A.; Poorter, H. Avoiding Bias in Calculations of Relative Growth Rate. *Annals of Botany* 2002, 90, 37–42, doi:10.1093/AOB/MCF140.
26. Kitahashi, Y.; Ichie, T.; Maruyama, Y.; Kenzo, T.; Kitaoka, S.; Matsuki, S.; Chong, L.; Nakashizuka, T.; Koike, T. Photosynthetic Water Use Efficiency in Tree Crowns of *Shorea beccariana* and *Dryobalanops aromatica* in a Tropical Rain Forest in Sarawak, East Malaysia. *Photosynthetica* 2008 46:1 2008, 46, 151–155, doi:10.1007/S11099-008-0025-9.