

## **Original Research Article**

### **Characterization of *Brassica napus* (Canola) germplasm based on Microsatellite Markers**

#### **Abstract**

*Brassica napus* L. is a major oilseed crop all over the world. The aim of this study was to investigate the genetic diversity of *B. napus* germplasm by using simple sequence repeats (SSR) markers. In the current study, ten SSR markers were used for studying genetic diversity of ten *Brassica* cultivars. Out of 110 total bands, 68 bands were polymorphic with 52.11% average polymorphism. Mean value of Nei's genetic diversity and Polymorphism Information Content was 1.7, and 0.2630, respectively. These mean values show that there are moderate allelic differences between *Brassica* cultivars. The Nei's genetic distance among various cultivars was 0.3281 and 0.125 which showed that germplasm of *Brassica* cultivars are different from each other, which is probably due to anthropogenic interventions and environmental factors. Thus, genetically

different lines identified in this study could be employed in breeding programmes to develop higher-quality canola inbred varieties in future.

**Keywords:** *Brassica napus*, Genetic diversity, Simple sequence repeats, microsatellite

## 1. Introduction

Mustard (*B. juncea*) and Rapeseed (*Brassica napus* and *B. rapa*) are major oilseed crop around the world. In Pakistan, the crop is cultivated on an area of 272,100 ha, yielding 230,000 tons per year with 812 kg/ha average yield, which account for more than 17% of Pakistan's total domestic edible oil production [1]. Among the genus, genetic diversity provides a significant variation source that may be exploited to change *Brassica* crop cultivars using a variety of ways. Analyses of genetic relations are critical for germplasm management, crop improvement programs, and conservation strategy development. Local cultivars of oil crops usually contain good flavour and quality, as well as exhibit high level of disease and pest resistance, and could be preferable than foreign

materials. Therefore, plant breeding requires the development of genetically heterogeneous gene pools [2].

However, modern plant-breeders face the challenge of developing higher yielding, more nutritional, and ecologically friendly cultivars that increase our life quality without requiring the use of new habitats for agriculture [3]. Plant breeders cannot produce varieties that fulfil changing needs in terms of adaptability to growing circumstances, yield, tolerance to abiotic or biotic challenges, or specific requirements of quality unless they access huge database of heterogeneous material of plant [4]. Therefore, accessibility to the diverse genetic diversity pool is the most effective approach to increase the performance of different cultivars. Furthermore, information regarding the genetic diversity collections of *B. napus* germplasm can provide geneticists and breeders with valuable information about the diversity of alleles found in *B. napus* genotypes, as well as aid in the identification of genetically distinct pools to be used in cross-combinations for improving major agronomic traits and exploit heterosis [5].

For the introduction of new diversity into breeding lines of oilseed rape, morphological, agronomical, and phenological parameters have traditionally been used. Since the emergence of modern molecular biology technologies, molecular markers are being used to investigate phylogenetic relationships among as well as within species of *Brassica*. These markers consist of randomly amplified polymorphic DNA [6], restriction fragment length polymorphism (RFLP) [7], and simple sequence repeat (SSR) [8] *etc.* When compared to other molecular marker approaches, SSR markers are abundant, informative, highly polymorphic, co-dominant, reproducible, technically simple, and reasonably inexpensive when information of primer is available. Moreover, SSR markers are frequently found in the gene-rich genomic areas, implying that they could be useful for association studies of allele-trait in well-characterized regions of genome with quantitative-trait loci.

SSR markers have been frequently employed in tomato, maize, and rice for diversity studies [9-11]. SSR markers have been proved beneficial for studying genetic structure and diversity of *Brassica*. Using eighteen pairs of SSR

primers, 112 polymorphic loci were discovered in Australian canola varieties, revealing their genetic diversity [12]. By evaluating the polymorphic alleles of eighteen SSR markers, Soengas et al. [13] was able to determine the genetic link between eight groups and investigate the genetic structure. Based on this, the present study was aimed to investigate the genetic diversity or polymorphism of different local varieties of *B. napus*, which would be useful for breeding program of *Brassica* in the future.

## 2. Material and Methods

### 2.1. Plant material

Seeds of ten different genotypes of *B. napus* (ACC-23633, ACC-24210, ACC-24214, ACC-24217, ACC-24254, ACC-24875, ACC-24880, ACC-24889, ACC-27396, and ACC-27405) were collected from the Pakistan Agriculture Research Council (PARC), Islamabad, Pakistan. The research was conducted at the Molecular Markers Application Laboratory at the Institute of Molecular Biology and Biotechnology, Bahauddin Zakariya University, Multan, Pakistan. To explore the genetic diversity of these genotypes, the following techniques were used.

## 2.2. Extraction of genomic DNA

Four week old seedlings were collected for extraction of DNA following the protocol of Doyle and Doyle [14] with minor modifications. 2mg leaf sample was homogenized in 500 $\mu$ l of extraction buffer [NaCl (100 mM), SDS (1%), Tris-base (100 mM), and Na<sub>2</sub>EDTA (100 mM)] following incubation in water bath for thirty minutes. After that, 500 $\mu$ l of phenol (25): chloroform (24): isoamyl alcohol (1) solution was added following centrifugation at 9000 rpm for 15 min. After centrifugation, aqueous layer was separated and 500 $\mu$ l chilled isopropanol was added for DNA precipitation following centrifugation at 9000rpm for 2 min. The pellet was washed with 70% ethanol after discarding the supernatant, and was dried for half hour at room temperature. After that, 50  $\mu$ l TE-Buffer was added in the tubes and was stored at 4°C. The quality of DNA was checked using agarose gel electrophoresis, while its quantity was measured using UV-spectrophotometer at 260nm wavelength.

### 2.3. SSR assays

Ten pairs of SSR markers i.e. BRASO-I-I, BRAS014, BRAS072A, BRAS078, BRAS084, CB-10026, CB-I-0028, CB-10092B, CBI-0-1-43, and CBI-0369 (Table 1) were used for the genetic diversity analysis of *B. napus*. SSR primers were obtained from <http://www.brassica.info/resource/markers.php> and were synthesized by MacroGen chill.co.pk (Table 1). PCR amplification of SSR loci was performed in 25 µL reaction mixture containing genomic DNA (20 ng/µL), 10 X PCR-reaction buffer, dNTPs (0.2mM), 0.5pmol of forward and reverse primer, Taq DNA polymerase (0.5U) and MgCl<sub>2</sub> (15mM). PCR was performed in PTC-100 thermo-cycler with the following temperature profile: 95 °C for 5 minutes (initial denaturation), following 35 cycles with 94 °C denaturation for 45 seconds, annealing for 60 seconds, primer elongation for 2 minutes at 72 °C and final elongation for 10 minutes at same temperature. The products of PCR were analysed on polyacrylamide gel electrophoresis containing 10% polyacrylamide using Bio-rad electrophoresis cell. The gel was stained by silver staining following protocol of Zhang et al., [15].

**Table 1 SSR primer for genetic diversity analysis of *Brassica* cultivars**

Sr. No.	Primer Information	
	Name	Sequence
1	BRASO-I-I F	TGG GAC GTA GTC AGT CAA CAA
	BRASO-I-I R	CCA AGT GCG AGA AGA GGA AG
2	BRAS014F	CCC AIT GAC AAC TCT TCT CIT
	BRAS014R	CTG TGT CGC CCA ITA TG
3	BRAS072AF	GCC ATC TAC ACA TTT ATC CC
	BRAS072AR	CAC TAA CCT TCT TGC TAC CGT
4	BRAS078F	ATT GGG TTC TGA CCI TIT CTC
	BRAS078R	CTT ITC CTC ATC GCT ACC AC

5	BRAS084F	ATT GGG TTC TGA CCT IIT CTC
	BRAS084R	TTT TCC TTC ATC GCT ACC AC
6	CB-10026F	TCG TTC TGA CCT GTC GIT AT
	CB-10026R	GGA AAT GGC TGC TCA TGC T
7	CB-I-0028F	CTG CAC ATT TGA AAT IGG TC
	CB-I-0028R	AAA TCA ACG CTT ACC CACT
8	CB-10092BF	TTG ATC CGA AAT CTC TGG
	CB-10092BR	AGG CAA GCA ATA GAT AAA GG
9	CBI-0-1-43F	CAT GGG AGG CTG TCT AAA
	CBI-0-1-43R	TTG CAC CCA TAC GTT TC
10	CBI-0369F	CAT CAC AGG ACC AGA GC
	CBI-0369R	CAA AGC CAA GAC ACC CAT

## 2.4. Data analysis

Bands expressing specific alleles on microsatellite loci were manually scored on gel pictures. The genotype of a specific locus was scored based on the allelic number having specific size at a certain locus. A band present on gel indicating the presence of a certain allele at one locus was labelled 1, whereas its absence was labelled 0. To create a single matrix, genomic data encompassing 10 SSR markers and 10 *Brassica* cultivars were entered into a Microsoft Excel spreadsheet. The number of effective alleles ( $N_E$ ) was derived from this dataset for each SSR marker and polymorphism percentage was also estimated to frame genetic-profile of *Brassica* crops. Nei's genetic diversity ( $H_E$ ) analogous to heterozygosity was also calculated because it estimates genetic diversity of randomly breeding populations. POPGENE version 1.32 (<http://www.ulaberta.ca/fyeh>) was used to calculate Pair wise Nei's genetic-distance. The genetic relatedness of all ten *Brassica* cultivars was assessed by computing genetic distances for all samples using Nei's coefficient. Polymorphism Information Content (PIC) value was calculated using  $PIC_i = 1 -$


$\sum_{j=1}^n (PIJ)^2$  formula,  $i$ =marker, where  $n$  and  $p_{ij}$  = the number and frequency of alleles, respectively for that marker. Un-weighted pair group method with arithmetic average (UPGMA) was used to do cluster analysis [16].

### 3. Results and Discussion

The study of genetic variation is beneficial to the evolution and conservation of species [17]. Molecular markers are frequently utilised in the genetic study of plants species due to their different properties such as high polymorphism, co-dominance, and reproducibility. Due to their multi-allelic character, these markers can easily detect a wide range of comparative allelic differences in germplasm [18].

In this study, SSR-DNA profile was created for investigating genetic relationship between 10 cultivars of *B. napus*. After amplification of PCR (Figure 1), out of 110 total detected loci 68 appeared to be polymorphic with 52.11% average of polymorphism (Table 2). Out of ten successfully amplified SSR markers, eight markers showed polymorphism and created potentially score-able banding array.

However, only one marker (BRAS072A) showed 100% polymorphism.  $N_E$  (no. effective alleles) ranged from 1.0000-2.000 with 1.7 mean values. Out of ten markers, BRAS084, CBI-0-1-43, and CBI-0369 showed 1.000  $N_E$  values than all other markers. The Nei's genetic diversity or  $H_E$  value ranged from 0.000 to 0.48 with an average of 0.298, while the polymorphism information content value (PIC) ranges between 0.0000 to 0.3750 with an average of 0.2360 (Table 2).






Figure 1 Amplification results of the primer (A) BRASO-I-I; (B) BRAS014; (C) BRAS072A; (D) BRAS078

The PIC value predicts the efficacy of any marker in genetic evaluation, linkage mapping, and molecular lineage. Allelic differences are more likely to be shown by markers with greater PIC values [19]. In this study, maximum value of PIC value was 0.3750 using CB-10092B marker, while the minimum was 0.000 with BRAS084 and CBI-0369 (Table 2), this could be attributable to inadequate coverage of genome or a low degree of genetic heterogeneity among cultivars. Since no marker in this study has a PIC value of 1, this indicates that there is no specific gene in the genome of *Brassica* cultivars. Average  $H_E$  and PIC

scores in the moderate range imply that there is moderate allelic variation across all the tested *Brassica* cultivars.

**Table 2 Parameters of genetic diversity of 10 *B. napus* samples with 10**

**SSR markers**

Primers	No. of loci		Polymorphism %	Allele frequency	N <sub>E</sub>	H <sub>E</sub>	PIC
	Total	Polymorphic					
BRASO-I-I	10	7	70%	0.7	2.000	0.42	0.3318
BRAS014	5	1	20%	0.9	2.000	0.18	0.1638
BRAS072	20	20	100%	0.8	2.000	0.32	0.2688
A							
BRAS078	8	4	50%	0.6	2.000	0.48	0.3648
BRAS084	4	0	0%	1.000	1.000	0.000	0.000

<b>CB-10026</b>	6	3	50%	0.6	2.000	0.48	0.3648
<b>CB-I-0028</b>	17	14	82.35%	0.7	2.000	0.42	0.3318
<b>CB-10092B</b>	19	18	94.73%	0.52	2.00 0	0.50 0	0.375 0
<b>CBI-0-1-43</b>	11	1	9.09%	0.9	1.000	0.18	0.1638
<b>CBI-0369</b>	10	0	0%	1.000	1.000	0.000	0.000
<b>Total</b>	110	68	52.11%	0.772	1.7	0.29 8	0.236

Genetic diversity assessment is fundamental knowledge that allows us to not only identify similarities and differences among plant species, but also help us to recognize environmental challenges on populations. Microsatellite markers are employed in genotyping of several species due to their co dominant, highly polymorphic nature [20], and their multi-allelic character allows to detect allelic

variants across a wide variety of genetic resources [21]. In the current study, 10 SSR markers on 10 *Brassica* cultivars were utilized in an efficient way for studying genetic diversity. All amplified markers, resulted in 68 polymorphic bands out of 110 clear bands. These results showed high efficacy of microsatellite or SSR markers and thus, can be an efficient and suitable tool for investigation of genetic variations of many species of plant [22, 23]. Previous researchers have examined the differences between winter and spring oilseed of Europe and China accessions using SSR markers [24, 25]. Several other studies investigated genetic diversity of rapeseed germplasm and cultivars which showed similar results as observed in the current study [26-28]. Furthermore, understanding the genetic relationships and diversity of germplasm among local varieties and primary breeding lines might be an effective tool in agricultural improvement methods.

On the basis of Nei's genetic distance, cluster analysis was done using UPGMA that divided all cultivars into three clusters (Table 3). Cultivars within the same cluster share genetic characteristics, indicating that they are near

relatives. Cultivars in one cluster are genetically more similar than cultivars in another. Each cluster is sub-divided into smaller groups called sub-clusters. There are two cultivars in Cluster I (ACC-27405 and ACC-23633) and Cluster II (ACC-27396 and ACC-24889). However, in Cluster III the germplasm ACC-24217 is sub-divided into two cultivars (ACC-24214 and ACC-24880). Moreover, the three germplasm; *i.e.* ACC-24875, ACC-24210 and ACC-24254 form out boundaries from all the other cultivars. Thus, the results obtained in this study showed that SSR markers are beneficial and effective for evaluating the genetic variation of *B. napus* germplasm. Several other researchers have come to similar results about using SSR markers in rapeseed breeding [27, 29].

**Table 3 Frequency based Nei's genetic distance**

	ACC-2363	ACC-2421	ACC-2421	ACC-2421	ACC-2425	ACC-2487	ACC-2488	ACC-2488	ACC-2739	ACC-2740
<b>Cultivars</b>	<b>3</b>	<b>0</b>	<b>4</b>	<b>7</b>	<b>4</b>	<b>5</b>	<b>0</b>	<b>9</b>	<b>6</b>	<b>5</b>

ACC- 23633	0									
ACC- 24210	0.328 1	0								
ACC- 24214	0.25	0.234 4	0							
ACC- 24217	0.265 6	0.25	0.171 9	0						
ACC- 24254	0.218 8	0.312 5	0.265 6	0.312 5	0					
ACC- 24875	0.359 4	0.328 1	0.218 8	0.25	0.375	0				
ACC- 24880	0.195 3	0.289 1	0.148 4	0.210 9	0.210 9	0.289 1	0			
ACC- 24889	0.132 8	0.320 3	0.226 6	0.210 9	0.273 4	0.335 9	0.203 1	0		
ACC-	0.179	0.335	0.164	0.257	0.289	0.289	0.187	0.140	0	

27396	7	9	1	8	1	1	5	6		
ACC-		0.343	0.234	0.281	0.171	0.328	0.179	0.148	0.195	
27405	0.125	8	4	3	9	1	7	4	3	0

## Conclusion

The present study is the first preliminary research that utilized microsatellite or SSR markers for reporting the diversity of *B. napus* using a small number of cultivars. For further genetic variations and phylogenetic relationships of this species in the future, a detailed study with a high sample size is recommended.

## References

1. Rashid A. Pakistan Agriculture Research Council: Rapeseed 2021  
[Available from: <http://www.parc.gov.pk/index.php/en/csi/137-narc/crop-sciences-institute/726-rapeseed>.
2. Bailey-Serres J, Parker JE, Ainsworth EA, Oldroyd GE, Schroeder JI. Genetic strategies for improving crop yields. Nat. 2019;575(7781):109-18.

3. Zamir D. Improving plant breeding with exotic genetic libraries. *Nat Rev Genet.* 2001;2(12):983-9.
4. Friedt W, Snowdon R, Ordon F, Ahlemeyer J. Plant breeding: assessment of genetic diversity in crop plants and its exploitation in breeding. *Progress in botany: Springer;* 2007. p. 151-78.
5. Diers B, Osborn T. Genetic diversity of oilseed *Brassica napus* germ plasm based on restriction fragment length polymorphisms. *Theor Appl Genet.* 1994;88(6-7):662-8.
6. Demeke T, Adams R, Chibbar R. Potential taxonomic use of random amplified polymorphic DNA (RAPD): a case study in *Brassica*. *Theor Appl Genet.* 1992;84(7-8):990-4.
7. McGrath JM, Quiros CF. Genetic diversity at isozyme and RFLP loci in *Brassica campestris* as related to crop type and geographical origin. *Theor Appl Genet.* 1992;83(6):783-90.

8. Plieske J, Struss D. Microsatellite markers for genome analysis in Brassica. I. Development in *Brassica napus* and abundance in Brassicaceae species. Theor Appl Genet. 2001;102(5):689-94.
9. Pathaichindachote W, Panyawut N, Sikaewtung K, Patarapuwadol S, Muangprom A. Genetic diversity and allelic frequency of selected Thai and exotic rice germplasm using SSR markers. Rice Sci. 2019;26(6):393-403.
10. Kumari A, Sinha S, Rashmi K, Mandal S, Sahay S. Genetic diversity analysis in maize (*Zea mays* L.) using SSR markers. J Pharm Phytochem. 2018;7:1116-20.
11. Abdulazeez SD, Kakarash SA, Ismael NB. Genetic Diversity Among Ten Maize Genotypes Using Simple Sequence Repeat DNA Markers. Polytech J. 2021;11(1):32-7.
12. Wang J, Kaur S, Cogan N, Dobrowolski MP, Salisbury P, Burton W, et al. Assessment of genetic diversity in Australian canola (*Brassica napus* L.) cultivars using SSR markers. Crop Pasture Sci. 2009;60(12):1193-201.

13. Soengas P, Cartea M, Francisco M, Lema M, Velasco P. Genetic structure and diversity of a collection of *Brassica rapa* subsp. *rapa* L. revealed by simple sequence repeat markers. *The J Agri Sci.* 2011;149(5):617-24.
14. Doyle J. DNA protocols for plants. *Molecular techniques in taxonomy*: Springer; 1991. p. 283-93.
15. Zhang J, Guo W, Zhang T. Molecular linkage map of allotetraploid (*Gossypium hirsutum* L. × *Gossypium barbadense* L.) with a haploid population 2002 24. *Theor Appl Genet.* (105):1166-74.
16. Sneath PH, Sokal RR. *Numerical taxonomy. The principles and practice of numerical classification* 1973.
17. Nielsen ES, Begger M, Henriques R, von der Heyden S. A comparison of genetic and genomic approaches to represent evolutionary potential in conservation planning. *Biol Conserv.* 2020;251:108770.
18. Bharti R, Chaudhary N, Gupta S, Sharma M. Molecular characterization of candidate gene in glucosinolate and erucic acid using SSR markers. *Int J Ad Res Ideas Innov Tech.* 2018;4(1):24-54.

19. Spandana B, Reddy VP, Prasanna GJ, Anuradha G, Sivaramakrishnan S. Development and characterization of microsatellite markers (SSR) in *Sesamum* (*Sesamum indicum* L.) species. *Appl Biochem Biotechnol.* 2012;168(6):1594-607.
20. Kumar S, Palve AS, Patel SK, Selvanayagam S, Sharma R, Rathore A. Development of genomic microsatellite markers in cluster bean using next-generation DNA sequencing and their utility in diversity analysis. *Curr Plant Biol.* 2020;21:100134.
21. Guang-Xin E, Hong QH, Zhao YJ, Ma YH, Chu MX, Zhu L, et al. Genetic diversity estimation of Yunnan indigenous goat breeds using microsatellite markers. *Eco Evol.* 2019;9(10):5916.
22. Lu J, Zhang Y, Diao X, Yu K, Dai X, Qu P, et al. Evaluation of genetic diversity and population structure of *Fragaria nilgerrensis* using EST-SSR markers. *Gene.* 2021;796:145791.
23. Sadiyah H, Ashari S, Waluyo B, Soegiarto A. Genetic diversity and relationship of husk tomato (*Physalis* spp.) from East Java Province revealed by SSR markers. *Biodiversitas.* 2021;22(1):184-92.

24. Shengwu H, Ovesna J, Kucera L, Kucera V, Vyvadilova M. Evaluation of genetic diversity of *Brassica napus* germplasm from China and Europe assessed by RAPD markers. *Plant Soil Environ.* 2003;49(3):106-13.
25. Hasan M, Seyis F, Badani A, Pons-Kühnemann J, Friedt W, Lühs W, et al. Analysis of genetic diversity in the *Brassica napus* L. gene pool using SSR markers. *Genet Res Crop Evol.* 2006;53(4):793-802.
26. Verma K, Tripathi MK, Tiwari S, Tripathi N. Analysis of genetic diversity among *Brassica juncea* genotypes using morphophysiological and SSR markers. *Int J Curr Microbiol App Sci.* 2021;10(01):1108-17.
27. Thakur A, Singh K, Parmar N, Sharma D, Mishra DC, Singh L, et al. Population structure and genetic diversity as revealed by SSR markers in Ethiopian mustard (*Brassica carinata* A. Braun): a potential edible and industrially important oilseed crop. *Genet Res Crop Evol.* 2021;68(1):321-33.
28. Singh V, Avtar R, Kumari N, Kumar R. Assessment of genetic diversity and population structure in Indian mustard (*Brassica juncea* L.) using SSR markers. *J Environ Biol.* 2021;42:396-405.

29. Singh M, Avtar R, Singh VK, MAHAVIR M, KUMARI N. Genetic diversity in Indian mustard (*Brassica juncea*) genotypes for Sclerotinia stem rot (*Seclerotinia sclerotiorum*) using SSR markers. India J Agric Sci. 2021;91(6).

UNDER PEER REVIEW