

Response of quality attributes of banana (*Musa paradisiaca* L.) cv. Williamto post shooting spray and bunch covering material

ABSTRACT

Aims: To study the post harvest shooting namely; control, humic acid 2 %, 2, 4-D 30 mg/l, gibberellic acid (GA₃) 100 mg/l, CPPU 4 mg/l and sulphate of potash (SOP) 2 % and covering materials viz., non- woven material bag and blue colour polyethylene sleeve (6% perforated) bag on bunch of banana cv. William

Study Design: Experiment was laid out in a Completely Randomized Design (Factorial) with three repetitions.

Place and Duration of Study: Experiment was carried out at the Horticultural Research Farm, Department of Horticulture, B. A. College of Agriculture, AAU, Anand during the year 2017-18 and 2018-19

Methodology: The experimental plot was prepared by deep ploughing, harrowing and levelling. The pits of 30 x 30 x 30 cm were dug out at a spacing of 1.8 x 1.8 m² and well decomposed fine textured Farm Yard Manure (FYM) at the rate of 10 kg per pit was applied at planting. Well hardened, healthy, uniform tissue cultured tissue culture plants of willium banana having 5-6 leaves were used for planting. Post shooting sprays were given twice i.e. 1st spray after complete opening of inflorescence and 2nd spray after 30 days of first spray with covering the bunch immediately after second spray. The fruits which were used for recording the weight loss during ripening, were used to calculating pulp: peel ratio. Pulp to peel ratio was calculated by dividing respective pulp weight by respective peel weight. The TSS value of the fruit was recorded by using hand refractometer having range of 0-32 ° Brix. Acidity (%) was calculated by the method described by Rangana (1977) was adopted for estimation of titrable acidity.

Results: The results revealed that the banana bunches spraying of CPPU 4 mg/l with bunch covering with non woven materials were recorded significantly lower physiological loss in weight (12.80, 13.60 and 13.20 %) and (13.26, 14.14 and 13.70 %), respectively. Whereas, post shooting spray of SOP 2 % with non-woven material bag covering were recorded significantly maximum TSS (21.42, 21.05 and 21.17 °Brix; 21.39, 20.93 and 21.16 °Brix), reducing sugar (12.55, 12.34 and 12.45 %; 12.21, 12.16 and 12.18 %) and total sugar (21.42, 20.82 and 21.12 %; 20.84, 20.50 and 20.67 %) as compared to rest of the treatments of post shooting sprays and bunch covering materials on banana quality.

Conclusion: From the two years of field study, it can be concluded that the post shooting spraying of CPPU 4 mg/l and SOP 2 % with non woven bunch covering bags were improve fruit quality of banana

KEYWORDS: Banana, post shooting spray, bunch covering materials and quality

INTRODUCTION:

Banana (*Musa paradisiaca* L.) belongs to family Musaceae, is the cheapest, plentiful and most nourishing fruit crop of the world. It is a premier fruit having great socio-economic significance in India. Indeed many consider the banana to be one of man's first foods. It is also a dessert fruit for millions, apart from a staple food owing to its rich and easily digestible carbohydrates with a calorific value of

67-137/100 g fruit. It is a good source of Vitamin A (190 IU per 100 g of edible portion) and Vitamin C (100 mg/100g) and fair source of Vitamin B1 and B2. Fruits are also rich source of minerals like magnesium, sodium, potassium, phosphorus and a fair source of calcium and iron. As a diet, it is highly satisfying, easy to digest, nearly fat free, rich source of carbohydrate with calorific value of 375 kJ per 100 g. It contains various vitamins and has therapeutic values for the treatment of many diseases (Singh, 2007)

Many reports have indicated banana post shooting spray of CPPU and GA₃ increases fruit size with stored under ambient condition found ripening inhibition, shelf life extension, quality maintenance of banana fruits during storage and delays senescence of fruit (Huanga et al. 2014). Also post shooting foliar spray of SOP on quality parameters of banana (*Musa paradisiaca* L.) cv. Grand Nain and found that the maximum reducing sugar and total sugar were recorded. While, maximum TSS and shelf life (Gamit et al. 2017)

For covering the fruit or bunches, different kinds of materials have been used so far. Brown paper, gunny bag, newsprint, old cloths, dried banana leaves, and polythene bags of different colors and thicknesses have been used (Debnath et al., 2001). The decision of using a bagging material depends on the availability, price, local climate, and type of fruit. All of these materials have some advantages and disadvantages. Blue polyethylene perforated bunch covers have been used elsewhere in the tropics and have been shown to improve fruit visual appeal among other postharvest qualities (Robinson, 1996). But there is very limited information found in India on effect of bunch covering and its impacts quality of banana cv. willium. Therefore, this present research work has been undertaken to find out the effect of post shooting spray and bunch covering materials on fruit quality of banana cv. willium.

MATERIAL AND METHODS:

An experiment entitled Response of quality attributes of banana (*Musa paradisiaca* L.) cv. Willium to post shooting spray and covering material on bunch was conducted at Horticultural Research Farm, Department of Horticulture, B. A. College of Agriculture, Anand Agricultural University, Anand during the years 2017-18 and 2018-19. Experiment was laid out in a Completely Randomized Design (Factorial) with three repetitions. The experimental plot was prepared by deep ploughing, harrowing and levelling. The pits of 30 x 30 x 30cm were dug out at a spacing of 1.8 x 1.8 m² and well decomposed fine textured Farm Yard Manure (FYM) at the rate of 10 kg per pit was applied at planting. Well hardened, healthy, uniform tissue cultured tissue culture plants of willium banana having 5-6 leaves were used for planting. The experiment comprises of involving six levels of post shooting sprays namely; control, humic acid 2%, 2, 4-D 30 mg/l, gibberellic acid (GA₃) 100 mg/l, N-(2-Chloro-4-pyridyl)-N-phenylurea (CPPU) 4 mg/l and sulphate of potash (SOP) 2% with two levels of bunch covering material viz., non- woven material bag and blue colour polyethylene sleeve (6% perforated) bag. Post shooting sprays were given twice i.e. 1st spray after complete opening of inflorescence and 2nd spray after 30 days of first spray with covering the bunch immediately after second spray. Observations were recorded daily for quality characters i.e. Pulp :peel ratio, TSS (° Brix), Acidity (%), Reducing sugars (%), Total sugars (%) and Physiological loss in weight (%) were recorded. The fruits which were used for recording the weight loss during ripening, were used to calculating pulp: peel ratio. Pulp to peel ratio was calculated by dividing respective pulp weight by respective peel weight. The TSS value of the fruit was recorded by using hand refractometer having range of 0-32 ° Brix. Acidity (%) was calculated by the method described by Rangana (1977) was adopted for estimation of titrable acidity. The titrametric method of Lane and Eynon described by Ranganna (1979) was adopted for estimation of reducing sugar (%) and total sugars (%). The data recorded during the course of investigation were subjected to statistical analysis following standard procedure described by Gomez and Gomez (1984).

RESULTS AND DISCUSSION:

Physiological loss in weight (%):

Effect of post shooting spray

The result revealed that physiological loss in weight of banana was influenced significant by various post shooting sprays during the both experimental years 2017-18, 2018-19 and in pooled basis. The post shooting spray of CPPU 4 mg/l (S_5) recorded significantly lower physiological loss in weight (12.80, 13.60 and 13.20 %) and which was found at par with treatment SOP 2% (13.02, 13.53 and 13.28 %) and GA_3 100 mg/l (13.52, 14.03 and 13.77 %). Whereas, significantly higher physiological loss in weight was found under control (17.52, 18.33 and 17.93 %) during the years 2017-18, 2018-19 and in pooled analysis, respectively. This might be due to the reason that CPPU plays vital role in enhancing the physiological activities in **delayed color change, suppressed fruit softening and transformation of starch to sucrose, and maintained high contents of ascorbic acid and total phenolic compounds and the inhibition of the peaks of ethylene production rate in banana fruit during storage at ambient temperature** reported by Huang *et al.*, 2014 and Rajan, 2017. Similar results were obtained by Marzouk and Kassem (2011) in grapes.

Effect of bunch covering material

It is evident from the observations on bunch covering materials exerted significant effect on physiological loss in weight. Bunch covering with non woven materials gave significantly minimum physiological loss in weight (13.26, 14.14 and 13.70 %) as compared to blue colour polythene sleeve (15.30, 15.85 and 15.58 %) during 2017-18, 2018-19 and in pooled, respectively.

Decreased in fruit weight during the storage period could be due to physiological process such as higher rate of respiration and transpiration from fruit surface degradation of reserve carbohydrate with release of water and transpiration through fruit skin. It could be also due to decrease in peel thickness with passage of ripening period which accelerates the weight loss of fruit (Patil and Shanmugasundaram, 2015; Santosh *et al.*, 2017). The decline in fruit weight has faster rate in control as compare to non-woven bags. These results are accordance with Pathak *et al.* (2017) in banana

Pulp: peel ratio:

Effect of post shooting spray

Data (Table 1) pertaining to various post shooting sprays exerted significantly effect on pulp: peel ratio. The post shooting spray with CPPU 4mg/l (S_5) recorded significantly maximum pulp: peel ratio with numerical value 3.22, 3.17 and 3.19 which was statistically found at par with treatments S_6 i.e. SOP 2% (3.18, 3.17 and 3.18) and S_4 i.e. GA_3 100mg/l (3.15, 3.04 and 3.10) as compared to rest of the treatments. Whereas, significantly minimum pulp: peel ratio was found under control (2.58, 2.64 and 2.61) in the years 2017-18, 2018-19 and in pooled analysis, respectively. This might be due to the reason that CPPU plays vital role in enhancing the physiological activities in suppressed fruit softening in association with the delayed peaks of respiration and the inhibition of the peaks of ethylene production rate in banana (Rajan 2017). Similar results were obtained by Curry and Greene (1993) in apple, Marzouk and Kassem (2011) in grapes.

Effect of bunch covering material

It is evident from observation that bunch covering materials significant effect on pulp: peel ratio. The significantly increasing the pulp: peel ratio was observed with non-woven material bag (3.18, 3.13 and 3.16) as compared to blue colour polythene sleeve (2.84, 2.81 and 2.83) during both experimental years as well as in pooled data, respectively. Pulp: peel ratio of banana fruit tends to increase with the advancement from harvest to ripening. In banana fruit moisture content of peel reduced gradually during ripening while that of the pulp increased with ripening hence pulp: peel ratio increased gradually (Burdon *et al.*, 1994). Pulp portion continues to grow even in later stage of maturation and bunch protection material enhances the fruit quality (Nakasone and Paul, 1998) in banana. Similar results were also noted by Sarkar (2014) and Santosh *et al.* (2017).

Total soluble solids (°brix):

Effect of post shooting spray

The result revealed that total soluble solids of banana fruit was significantly influenced by various post shooting sprays. Post shooting spray with SOP 2% (S₆) recorded significantly higher total soluble solids (21.42°Brix) and which was found at par with all treatments except Control in the year 2017-18. Whereas, significantly higher total soluble solids was recorded with post shooting spray of SOP 2% (21.23 and 21.33 °Brix) and which was found at par with treatments S₅ i.e. CPPU 4mg/l (21.05 and 21.17°Brix) and S₄ i.e. GA₃100 mg/l (20.45 and 20.60°Brix) in the years 2018-19 and in pooled analysis, respectively. The maximum TSS was noted in bunch spray with SOP 2% treatment. This might be due to post-shooting application of K favours the conservation of starch into simple sugars during ripening by activating sucrose synthase enzyme, resulting in high total soluble solid content in fruits. Similar results were also noted by Kumar *et al.* (2008), Kumar and Kumar (2010), Gamit *et al.* (2017) and Kachhadia *et al.* (2017) in banana.

Effect of bunch covering material

It is evident from the observations (Table 2), bunch covering materials significant effect on total soluble solids. The significantly the maximum total soluble solids was recorded under non-woven material bag covering (B₁) with numerical value 21.39, 20.93 and 21.16°Brix as compared to blue colour polythene sleeve (B₂) with numerical value 19.24, 19.18 and 19.21°Brix during individual years as well as in pooled, respectively. It might be due to during the climacteric stage, the accumulated polysaccharide is rapidly degraded and most of it is converted into soluble sugars which form a large proportion of TSS in the banana (Seymour *et al.* 1993). The present result of non-woven bag is coincide with findings of Sarkar (2014) and Santosh *et al.* (2017) in banana.

Acidity (%):

Effect of post shooting spray

All the treatments of post shooting sprays, bunch covering material and all interaction effect were found non-significant in regarding of acidity during the years 2017-18, 2018-19 and in pooled analysis (Table 2).

Reducing sugar (%):

Effect of post shooting spray

Results revealed (Table 3) that various treatments of post shooting sprays was found significant in respect to reducing sugar of banana fruits. The significantly maximum reducing sugar was recorded with SOP 2% (S₆) i.e. 12.55 % and which was at par with all the treatments except control in year 2017-18. Whereas, significantly maximum reducing sugar was recorded with spray of SOP 2% (12.34 and 12.45 %) and which was at par with treatments S₄ (GA₃100 mg/l) and CPPU 4mg/l (S₅) in the years 2017-18, 2018-19 and in pooled analysis, respectively.

Effect of bunch covering material

The result revealed that reducing sugar of banana fruits was influenced significantly by different bunch covering materials. The reducing sugar was recorded higher with non-woven material bag covering (B₁) i.e. 12.21, 12.16 and 12.18 % as compared to blue colour polythene sleeve (B₂) i.e. 11.13, 10.87 and 11.00 % during the year 2017-18, 2018-19 and in pooled, respectively.

Total sugar (%):

Effect of post shooting spray

The results revealed that various post shooting sprays was found significant for total sugar of banana fruits. The significantly maximum total sugar was recorded with spraying of SOP 2% (21.42 %) and it was found at par with all the treatments except control (S₁) in first year of experiment. Whereas, significantly maximum reducing sugar was recorded with treatment SOP 2% (20.82 and 21.12 %) and which was statistically found at par with treatments S₄ i.e. GA₃100 mg/l (20.33 and 20.40 %) and S₅ i.e. CPPU 4mg/l (20.10 and 20.35 %) in the year 2018-19 and in pooled analysis, respectively. This might be due to potassium is involved in carbohydrate synthesis, breakdown and translocation and synthesis of protein and neutralization of physiologically important organic acids (Tisdale and Nelson, 1966). Potassium is responsible for energy production in the form of ATP and NADPH in chloroplast by maintaining balanced electric charges. Besides, K is involved in phloem loading and unloading of

sucrose and amino acids and storage inform of starch in developing fruits by activating the enzymes starch synthase (Mengel and Kirkby, 1987). In plants well supplied with K, the osmotic potential of the phloem sap and the volume flow rate are higher than untreated plants and as a result, sucrose concentration in the phloem sap is increased. Similar results were also recorded by Kumar *et al.*(2008), Kumar and Kumar (2010), Kachhadia *et al.*(2017) and Pebbudiet *al.* (2017) in banana.

Effect of bunch covering material

The result revealed that total sugar of banana fruit was influenced by different bunch covering materials. The significantly higher total sugar was recorded with non-woven material bag covering (20.84, 20.50 and 20.67 %) as compared to blue colour polythene sleeve (19.00, 18.34 and 18.67 %) during the years 2017-18, 2018-19 and in pooled, respectively. It might be due to significantly positive association with temperature. The covered bunches had more total sugar, probably higher temperature inside bunch which is favorable for conversion of starch into sugar. A similar result was reported by Sarkar (2014) in banana.

Interaction effect

All interaction effects of post shooting sprays x bunch covering materials were found non-significant with respect to physiological loss in weight, pulp: peel ratio, total soluble solid, acidity, reducing sugar and total sugar of banana during both experimental years as well as in pooled data.

CONCLUSION:

From the two years of field study, it can be concluded that the post shooting bunches spraying of CPPU 4 mg/l was obtained minimum physiological weight of fruit and higher pulp: peel ratio. Whereas, post shooting spray of SOP 2 % was recorded maximum TSS, reducing sugar and total sugar with bunches covering of non-woven material bag was significantly better for improving the quality of banana.

REFERENCES:

- Burdon JN, Dori S, Lomaniec E, Marinansky R, Pesis E. The post-harvest ripening of water stressed banana fruits. *The Journal of Horticultural Sciences*, 1994;69(5): 799-804.
- Curry EA, Greene DW. CPPU influences fruit quality, fruit set, return blossom and preharvest drop of apples. *Horticultural Science*, 1993;28:115-119.
- Debnath U, Suresh CP, Hasan MA. Bunch management for profitable production in winter developing banana bunches. *Indian Journal of Horticulture*, 2001;58(3): 202-207.
- Gamit S, Patil SJ, Prajapti D. Effect of post shooting foliar spray of fertilizers on quality parameters of banana (*Musa paradisiacal L.*) cv. Grand Nain. *International Journal of Chemical Studies*, 2017;5(4):959-960.
- Gamit, S., Patil, S. J. & Prajapti, D. Effect of post shooting foliar spray of fertilizers on quality parameters of banana (*Musa paradisiacal L.*) cv. Grand Nain. *International Journal of Chemical Studies*, 2017: 5(4), 959-960.
- Gomez KA, Gomez AA. Statistical procedures for agricultural research (2nd ed.), John Wiley and Sons, New York. 1984.
- Huanga H, Jimh G, Wanga, Duana X, Qua H, Jianga Y. The combined effects of phenylurea and gibberellins on quality maintenance and shelf life extension of banana fruit during storage. *Scientia Horticulturae*, 2014;167:36-42.
- Huanga, H., Jimh, G., Wanga, H., Duana, X., Qua, H. & Jianga, Y. The combined effects of phenylurea and gibberellins on quality maintenance and shelf life extension of banana fruit during storage. *Scientia Horticulturae*, 2014:167, 36-42
- Kachhadia P, Tank RV, Bhanderi DR, Vagadia PS. Response of post-shooting bunch spray of chemicals on yield of ratoon banana (*Musa paradisiaca L.*) cv. Grand Nain. *Trends in Biosciences*, 2017;10(16):2951-2953.
- Kumar AR, Kumar N, Jeyakumar P. Effect of post-shooting spray of Sulphate of Potash (SOP) on yield and quality of banana cv. Robusta. *Research Journal of Agriculture and Biological Sciences*, 2008;4(6):655-659.
- Kumar, A. R. and Kumar, N. 2010. Effect of post-shooting spray of certain nutrients on yield and quality of banana cv. Ney Poovan (AB). *Agricultural Science Digest*, 30(2), 125-128.

- Mahalakshmi M, Sathiyamoorthy S. Effect of post shoot application of potassium on yield characters in banana cv. Rasthali (AAB). *South Indian Horticulture*, 1999;47(1/6):155-157.
- Marzouk HA, Kassem HA. Improving yield, quality, and shelf life of Thompson seedless grapevine by preharvest foliar applications. *Scientia Horticulturae*, 2011;130:425–430.
- Mengel K, Kirkby EA. Principles of Plant Nutrition, *International Potash Institute, Bern*, 1987; 436-437.
- Nakasone HY, Paul RE. Tropical fruits. CAB International New York. 1998.
- Pathak P, Baruah K, Bhattacharyya RK, Kalita P, Baishya BK. Influence of bunch covers on yield of banana cv. Jahaji (AAA) under high density planting system. *International journal of pure and applied bioscience*, 2017;5 (6):1488-1493.
- Patil SK, Shanmugasundaram S. Physicochemical changes during ripening of Monthan banana. *International journal of Technology Enhancements and Emerging Engineering research*, 2015;3(2):18-21.
- Pebbudi A, Bauri FK, Sarkar SK. Effect of potassium spray on fruit quality and shelf life of banana cv. Grand Naine (AAA). *Environment and Ecology*, 2017;35(1A):368-371.
- Rajan R. Effect of post shooting bunch spray of chemicals on bunch characters and yield of banana (*Musa paradisiaca*L.) cv. Grand Naine. M. Sc. (Horti.) in Fruit Science thesis submitted to Navsari Agricultural University, Navsari, Gujarat; 2017.
- Ranganna S. Manual of analysis of fruit and vegetable products. Tata McGraw Hill Publishing Company Ltd., New Delhi; 1979.
- Robinson JC. Bananas and Plantains. Institute for Tropical and Subtropical Crops, University Press, Cambridge, UK. 1996;172-174.
- Santosh DT, Tiwari KN, Reddy RG. Banana bunch covers for quality banana production – A Review. *International Journal of Current Microbiology and Applied Sciences*, 2017;6(7):1275-1291.
- Sarkar S. Shooting-harvest interval and physico-chemical properties of banana (*Musa* AAA cv. Grand Naine) in relation to micro climate inside the bunch cover. *The Asian Journal of Horticulture*, 2014;9(1):40-42.
- Seymour GB, Taylor JE, Tucker GA. Biochemistry of fruit ripening Seymour G.B., Ed. Chapman and Hall, London, 1993;83-106.
- Singh HP. National and International scenario of Banana and Plantain. "Banana: Technological Advancement". Ed. Singh, HP. Uma, S. 2007; 1-19.
- Swietlik D, Faust M. Foliar nutrition of fruit crops. *Hort. Rev.*, 1984;6:287-355.
- Tisdale SL, Nelson WL. *Soil Fertility and Fertilizers*, Macmillan Co., London, 1966;81.

Table 1: Effect of post shooting sprays and bunchcovering materialson physiological loss in weight and pulp: peel ratio

Treatments	Physiological loss in weight (%)			Pulp: peel ratio		
	2017-18	2018-19	Pooled	2017-18	2018-19	Pooled
Post shooting spray (S)						
S ₁ : Control	17.52	18.33	17.93	2.58	2.64	2.61
S ₂ : Humic acid @ 2%	14.57	15.23	14.90	2.96	2.95	2.95
S ₃ : 2,4-D @ 30 mg/l	14.25	15.27	14.76	2.96	2.87	2.92
S ₄ : GA3 @ 100 mg/l	13.52	14.03	13.77	3.15	3.04	3.10
S ₅ : CPPU @ 4 mg/l	12.80	13.60	13.20	3.22	3.18	3.19
S ₆ : SOP @ 2%	13.02	13.53	13.28	3.18	3.17	3.18
S.Em ±	0.31	0.35	0.23	0.08	0.06	0.05
CD at 5%	0.89	1.02	0.66	0.23	0.18	0.14
Bunch Covering Material (B)						
B ₁ : Non- woven material bag covering	13.26	14.14	13.70	3.18	3.13	3.16
B ₂ : Blue colour polyethylene sleeve	15.30	15.85	15.58	2.84	2.81	2.83
S.Em ±	0.18	0.20	0.13	0.05	0.04	0.03
CD at 5%	0.52	0.59	0.66	0.13	0.10	0.14
Interaction effect (S X B)						
S.Em ±	0.43	0.49	0.47	0.11	0.09	0.12
CD at 5%	NS	NS	NS	NS	NS	0.14
Pooled Interaction						
Source	Y x S	Y x B	YxSxB	Y x S	Y x B	YxSxB
S.Em ±	0.33	0.19	0.46	0.07	0.04	0.10
CD at 5%	NS	NS	NS	NS	NS	NS
CV %	5.25	5.69	5.48	6.39	5.07	5.78

Table 2: Effect of post shooting sprays and bunch covering materials on total soluble solids and acidity

Treatments	Total soluble solids (°Brix)			Acidity (%)		
	2017-18	2018-19	Pooled	2017-18	2018-19	Pooled
Post shooting spray (S)						
S ₁ : Control	18.37	18.28	18.32	0.294	0.305	0.299
S ₂ : Humic acid @ 2%	19.78	19.58	19.68	0.280	0.292	0.286
S ₃ : 2,4-D @ 30 mg/l	20.28	19.75	20.02	0.280	0.295	0.287
S ₄ : GA3 @ 100 mg/l	20.75	20.45	20.60	0.275	0.290	0.282
S ₅ : CPPU @ 4 mg/l	21.28	21.05	21.17	0.278	0.286	0.282
S ₆ : SOP @ 2%	21.42	21.23	21.33	0.276	0.283	0.279
S.Em ±	0.58	0.47	0.37	0.010	0.006	0.006
CD at 5%	1.71	1.37	1.07	NS	NS	NS
Bunch Covering Material (B)						
B ₁ : Non- woven material bag covering	21.39	20.93	21.16	0.281	0.292	0.287
B ₂ : Blue colour polyethylene sleeve	19.24	19.18	19.21	0.280	0.291	0.286
S.Em ±	0.34	0.27	0.22	0.006	0.004	0.003
CD at 5%	0.99	0.79	1.07	NS	NS	NS
Interaction effect (S X B)						
S.Em ±	0.83	0.66	0.66	0.015	0.009	0.004
CD at 5%	NS	NS	NS	NS	NS	NS
Pooled Interaction						
Source	Y x S	Y x B	YxSxB	Y x S	Y x B	YxSxB
S.Em ±	0.53	0.31	0.75	0.008	0.005	0.012
CD at 5%	NS	NS	NS	NS	NS	NS
CV %	7.05	5.71	6.42	8.960	5.210	7.260

Table 3: Effect of post shooting sprays and bunch covering materials on reducing sugar and total sugar (%)

Treatments	Reducing sugar (%)			Total sugar (%)		
	2017-18	2018-19	Pooled	2017-18	2018-19	Pooled
Post shooting spray (S)						
S ₁ : Control	10.52	10.19	10.35	17.95	17.18	17.57
S ₂ : Humic acid @ 2%	11.42	11.31	11.36	19.48	19.07	19.28
S ₃ : 2,4-D @ 30 mg/l	11.49	11.29	11.39	19.60	19.00	19.30
S ₄ : GA3 @ 100 mg/l	11.99	12.05	12.02	20.47	20.33	20.40
S ₅ : CPPU @ 4 mg/l	12.07	11.92	12.00	20.60	20.10	20.35
S ₆ : SOP @ 2%	12.55	12.34	12.45	21.42	20.82	21.12
S.Em ±	0.39	0.33	0.25	0.67	0.55	0.43
CD at 5%	1.13	0.95	0.72	1.94	1.60	1.23
Bunch Covering Material (B)						
B ₁ : Non- woven material bag covering	12.21	12.16	12.18	20.84	20.50	20.67
B ₂ : Blue colour polyethylene sleeve	11.13	10.87	11.00	19.00	18.34	18.67
S.Em ±	0.22	0.19	0.15	0.38	0.32	0.25
CD at 5%	0.65	0.55	0.72	1.12	0.92	1.23
Interaction effect (S X B)						
S.Em ±	0.55	0.46	0.46	0.94	0.78	0.78
CD at 5%	NS	NS	NS	NS	NS	NS
Pooled Interaction						
Source	Y x S	Y x B	YxSxB	Y x S	Y x B	YxSxB
S.Em ±	0.36	0.21	0.50	0.61	0.35	0.86
CD at 5%	NS	NS	NS	NS	NS	NS
CV %	8.10	6.92	7.54	8.19	6.92	7.60